



low\_Egamma L proton: max  $C_{\mu\gamma}$  regions



# L proton characteristic max $C_{\mu\gamma}$ configuration

3D momenta




c\_e\_gamma\_low\_egamma\_l\_proton\_region\_0  $C_{\mu\gamma}=0.239545$   
 region: low\_Egamma\_L\_proton\_region\_0 final-pair delta\_xy=0 rad  
 outgoing-state purity: 0.724714  
 kinematic point: high\_s\_high\_theta\_in\_low\_Egamma  
 incoming state:  $\frac{1}{2}\sum_{h_\mu}\rho(h_\mu, L_p)$


$s=21.5158$ ,  $\sqrt{s}=4.63852$ ,  $|\vec{P}|=2.22311$ ,  $|\vec{P}'|=2.22295$   
 $\theta_{in}=1.57079$ ,  $\phi_l=3.04342$ ,  $\phi_p=6.18501$   
 $E_\gamma=0.25$ ,  $\phi_\gamma=4.71239$   
 $Q^2=9.63214$ ,  $x_B=0.806038$ ,  $t=-10.8525$ ,  $W^2=3.19769$ ,  $y=0.579398$   
 $\theta(l', \gamma)=0$  deg

four-momenta  $[E, p_x, p_y, p_z]$  GeV:  
 $l$   $E= 2.2256$   $p_x= -2.2124$   $p_y= 0.2179$   $p_z= -0.0000$   
 $P$   $E= 2.4129$   $p_x= 2.2124$   $p_y= -0.2179$   $p_z= 0.0000$   
 $l'$   $E= 1.9758$   $p_x= 0.0000$   $p_y= -1.9729$   $p_z= 0.0000$   
 $P'$   $E= 2.4127$   $p_x= 0.0000$   $p_y= 2.2229$   $p_z= 0.0000$   
 $q_\gamma$   $E= 0.2500$   $p_x= -0.0000$   $p_y= -0.2500$   $p_z= 0.0000$

Transverse plane




Leading initial-ensemble/final-state components (N=6, retained=98.8%)



# L proton characteristic max $C_{\mu\gamma}$ configuration

3D momenta




c\_e\_gamma\_low\_egamma\_l\_proton\_region\_1  $C_{\mu\gamma}=0.239545$   
 region: low\_Egamma\_L\_proton\_region\_1 final-pair delta\_xy=0 rad  
 outgoing-state purity: 0.724714  
 kinematic point: high\_s\_high\_theta\_in\_low\_Egamma  
 incoming state:  $\frac{1}{2}\sum_{h_\mu} \rho(h_\mu, L_p)$


$s=21.5158$ ,  $\sqrt{s}=4.63852$ ,  $|\vec{P}|=2.22311$ ,  $|\vec{P}'|=2.22295$   
 $\theta_{in}=1.57079$ ,  $\phi_l=0.0981748$ ,  $\phi_p=3.23977$   
 $E_\gamma=0.25$ ,  $\phi_\gamma=4.71239$   
 $Q^2=9.63214$ ,  $x_B=0.806038$ ,  $t=-10.8525$ ,  $W^2=3.19769$ ,  $y=0.579398$   
 $\theta(l', \gamma)=0$  deg

four-momenta  $[E, p_x, p_y, p_z]$  GeV:  
 $l$   $E= 2.2256$   $p_x= 2.2124$   $p_y= 0.2179$   $p_z= -0.0000$   
 $P$   $E= 2.4129$   $p_x= -2.2124$   $p_y= -0.2179$   $p_z= 0.0000$   
 $l'$   $E= 1.9758$   $p_x= 0.0000$   $p_y= -1.9729$   $p_z= 0.0000$   
 $P'$   $E= 2.4127$   $p_x= 0.0000$   $p_y= 2.2229$   $p_z= 0.0000$   
 $q_\gamma$   $E= 0.2500$   $p_x= -0.0000$   $p_y= -0.2500$   $p_z= 0.0000$

Transverse plane




Leading initial-ensemble/final-state components (N=6, retained=98.8%)



# L proton characteristic max $C_{\mu\gamma}$ configuration

3D momenta




c\_e\_gamma\_low\_egamma\_l\_proton\_region\_2  $C_{\mu\gamma}=0.05397$   
region: low\_Egamma\_L\_proton\_region\_2 final-pair delta\_xy=0.110591 rad  
outgoing-state purity: 0.999727  
kinematic point: high\_s\_high\_theta\_in\_low\_Egamma  
incoming state:  $\frac{1}{2}\sum_{h_\mu}$

$s=21.5158$ ,  $\sqrt{s}=4.63852$ ,  $|\vec{P}|=2.22311$ ,  $|\vec{P}'|=2.22224$   
 $\theta_{in}=1.57079$ ,  $\phi_\ell=1.66897$ ,  $\phi_P=4.81056$   
 $E_\gamma=0.25$ ,  $\phi_\gamma=4.61421$   
 $Q^2=17.518$ ,  $x_B=0.883417$ ,  $t=-19.7136$ ,  $W^2=3.19166$ ,  $y=0.961454$   
 $\theta(\ell', \gamma)=6.33641$  deg

four-momenta  $[E, p_x, p_y, p_z]$  GeV:  
 $\ell$   $E= 2.2256$   $p_x= -0.2179$   $p_y= 2.2124$   $p_z= -0.0000$   
 $P$   $E= 2.4129$   $p_x= 0.2179$   $p_y= -2.2124$   $p_z= 0.0000$   
 $\ell'$   $E= 1.9764$   $p_x= 0.0245$   $p_y= -1.9734$   $p_z= 0.0000$   
 $P'$   $E= 2.4121$   $p_x= 0.0000$   $p_y= 2.2222$   $p_z= 0.0000$   
 $q_\gamma$   $E= 0.2500$   $p_x= -0.0245$   $p_y= -0.2488$   $p_z= 0.0000$

Transverse plane



Leading initial-ensemble/final-state components (N=2, retained=99.8%)

