



high\_Egamma L muon + Ty proton: max  $C_{\mu\gamma}$  regions



# L muon + Ty proton characteristic max $C_{\mu\gamma}$ configuration

3D momenta




c\_e\_gamma\_high\_egamma\_lty\_region\_0  $C_{\mu\gamma}=0.556072$   
 region: high\_Egamma\_LTy\_region\_0 final-pair delta\_xy=0.510997 rad  
 outgoing-state purity: 1  
 kinematic point: high\_s\_high\_theta\_in\_high\_Egamma  
 incoming state:  $|L_\mu\rangle \otimes |T_{\gamma,p}\rangle$

$s=21.5158$ ,  $\sqrt{s}=4.63852$ ,  $|\vec{P}|=2.22311$ ,  $|\vec{P}'|=2.19415$   
 $\theta_{in}=1.57079$ ,  $\phi_l=4.71239$ ,  $\phi_p=1.5708$   
 $E_\gamma=1.8$ ,  $\phi_\gamma=4.61421$   
 $Q^2=0.199845$ ,  $x_B=0.0120018$ ,  $t=-0.000128163$ ,  $W^2=17.3312$ ,  $y=0.807338$   
 $\theta(l', \gamma)=29.278$  deg


four-momenta  $[E, p_x, p_y, p_z]$  GeV:

$l$   $E= 2.2256$   $p_x= -0.0000$   $p_y= -2.2231$   $p_z= -0.0000$   
 $P$   $E= 2.4129$   $p_x= 0.0000$   $p_y= 2.2231$   $p_z= 0.0000$   
 $l'$   $E= 0.4523$   $p_x= 0.1764$   $p_y= -0.4028$   $p_z= 0.0000$   
 $P'$   $E= 2.3862$   $p_x= 0.0000$   $p_y= 2.1942$   $p_z= 0.0000$   
 $q_\gamma$   $E= 1.8000$   $p_x= -0.1764$   $p_y= -1.7913$   $p_z= 0.0000$

Transverse plane




Leading initial-ensemble/final-state components (N=6, retained=100.0%)



# L muon + Ty proton characteristic max $C_{\mu\gamma}$ configuration

3D momenta




c\_e\_gamma\_high\_egamma\_lty\_region\_1 C\_{\mu\gamma}=0.136216  
 region: high Egamma\_LTy\_region\_1 final-pair delta\_xy=0.510997 rad  
 outgoing-state purity: 1  
 kinematic point: high s\_high\_theta\_in\_high\_Egamma  
 incoming state:  $|L_\mu\rangle \otimes |\bar{T}_{y,p}\rangle$


$s=21.5158$ ,  $\sqrt{s}=4.63852$ ,  $|\vec{P}|=2.22311$ ,  $|\vec{P}'|=2.19415$   
 $\theta_{in}=1.57079$ ,  $\phi_l=0.687223$ ,  $\phi_P=3.82882$   
 $E_\gamma=1.8$ ,  $\phi_\gamma=4.81056$   
 $Q^2=3.73347$ ,  $x_B=0.184964$ ,  $t=-15.9448$ ,  $W^2=17.3312$ ,  $y=0.978667$   
 $\theta(l', \gamma)=29.278$  deg

four-momenta  $[E, p_x, p_y, p_z]$  GeV:  
 $l$   $E= 2.2256$   $p_x= 1.7185$   $p_y= 1.4103$   $p_z= -0.0000$   
 $P$   $E= 2.4129$   $p_x= -1.7185$   $p_y= -1.4103$   $p_z= 0.0000$   
 $l'$   $E= 0.4523$   $p_x= -0.1764$   $p_y= -0.4028$   $p_z= 0.0000$   
 $P'$   $E= 2.3862$   $p_x= 0.0000$   $p_y= 2.1942$   $p_z= 0.0000$   
 $q_\gamma$   $E= 1.8000$   $p_x= 0.1764$   $p_y= -1.7913$   $p_z= 0.0000$

Transverse plane




Leading initial-ensemble/final-state components (N=4, retained=99.2%)



# L muon + Ty proton characteristic max $C_{\mu\gamma}$ configuration

3D momenta




c\_e\_gamma\_high\_egamma\_lty\_region\_2  $C_{\mu\gamma}=0.136216$   
 region: high\_Egamma\_LTy\_region\_2 final-pair delta\_xy=0.510997 rad  
 outgoing-state purity: 1  
 kinematic point: high\_s\_high\_theta\_in\_high\_Egamma  
 incoming state:  $|L_\mu\rangle \otimes |T_{\gamma,p}\rangle$

$s=21.5158$ ,  $\sqrt{s}=4.63852$ ,  $|\vec{P}|=2.22311$ ,  $|\vec{P}'|=2.19415$   
 $\theta_{in}=1.57079$ ,  $\phi_l=2.45437$ ,  $\phi_p=5.59596$   
 $E_\gamma=1.8$ ,  $\phi_\gamma=4.61421$   
 $Q^2=3.73347$ ,  $x_B=0.184964$ ,  $t=-15.9448$ ,  $W^2=17.3312$ ,  $y=0.978667$   
 $\theta(l', \gamma)=29.278$  deg

four-momenta  $[E, p_x, p_y, p_z]$  GeV:

$l$   $E= 2.2256$   $p_x= -1.7185$   $p_y= 1.4103$   $p_z= -0.0000$   
 $P$   $E= 2.4129$   $p_x= 1.7185$   $p_y= -1.4103$   $p_z= 0.0000$   
 $l'$   $E= 0.4523$   $p_x= 0.1764$   $p_y= -0.4028$   $p_z= 0.0000$   
 $P'$   $E= 2.3862$   $p_x= 0.0000$   $p_y= 2.1942$   $p_z= 0.0000$   
 $q_\gamma$   $E= 1.8000$   $p_x= -0.1764$   $p_y= -1.7913$   $p_z= 0.0000$

Transverse plane



Leading initial-ensemble/final-state components (N=4, retained=99.2%)

