



low\_Egamma Tx proton: max  $C_{\mu\gamma}$  regions



# Tx proton characteristic max $C_{\mu\gamma}$ configuration

3D momenta




c\_e\_gamma\_low\_egamma\_tx\_proton\_region\_0  $C_{\mu\gamma}=0.0663755$   
 region: low\_Egamma\_Tx\_proton\_region\_0 final-pair delta\_xy=0.331208 rad  
 outgoing-state purity: 0.53875  
 kinematic point: high\_s\_high\_theta\_in\_low\_Egamma  
 incoming state:  $\frac{1}{2}\sum_{h_\mu} \rho(h_\mu, T_{x,p})$


$s=21.5158$ ,  $\sqrt{s}=4.63852$ ,  $|\vec{P}|=2.22311$ ,  $|\vec{P}'|=2.21665$   
 $\theta_{in}=1.57079$ ,  $\phi_l=4.71239$ ,  $\phi_p=1.5708$   
 $E_\gamma=0.25$ ,  $\phi_\gamma=5.00691$   
 $Q^2=0.0060701$ ,  $x_B=0.00267388$ ,  $t=-6.31712e-06$ ,  $W^2=3.14392$ ,  $y=0.110069$   
 $\theta(l', \gamma)=18.9768$  deg

four-momenta  $[E, p_x, p_y, p_z]$  GeV:  
 $l$   $E= 2.2256$   $p_x= -0.0000$   $p_y= -2.2231$   $p_z= -0.0000$   
 $P$   $E= 2.4129$   $p_x= 0.0000$   $p_y= 2.2231$   $p_z= 0.0000$   
 $l'$   $E= 1.9816$   $p_x= -0.0726$   $p_y= -1.9774$   $p_z= 0.0000$   
 $P'$   $E= 2.4069$   $p_x= 0.0000$   $p_y= 2.2167$   $p_z= 0.0000$   
 $q_\gamma$   $E= 0.2500$   $p_x= 0.0726$   $p_y= -0.2392$   $p_z= 0.0000$

Transverse plane




Leading initial-ensemble/final-state components (N=5, retained=92.2%)



# Tx proton characteristic max $C_{\mu\gamma}$ configuration

3D momenta




c\_e\_gamma\_low\_egamma\_tx\_proton\_region\_1 C\_{\mu\gamma}=0.0251841  
region: low\_Egamma\_Tx\_proton\_region\_1 final-pair delta\_xy=0 rad  
outgoing-state purity: 0.506527  
kinematic point: high\_s\_high\_theta\_in\_low\_Egamma  
incoming state:  $\frac{1}{2}\sum_{h_\mu}\rho(h_\mu, T_{x,p})$


$s=21.5158$ ,  $\sqrt{s}=4.63852$ ,  $|\vec{P}|=2.22311$ ,  $|\vec{P}'|=2.22295$   
 $\theta_{in}=1.57079$ ,  $\phi_\ell=2.55254$ ,  $\phi_P=5.69414$   
 $E_\gamma=0.25$ ,  $\phi_\gamma=4.71239$   
 $Q^2=13.6459$ ,  $x_B=0.854805$ ,  $t=-15.3748$ ,  $W^2=3.19769$ ,  $y=0.774004$   
 $\theta(\ell', \gamma)=0$  deg

four-momenta  $[E, p_x, p_y, p_z]$  GeV:  
 $\ell$   $E= 2.2256$   $p_x= -1.8484$   $p_y= 1.2351$   $p_z= -0.0000$   
 $P$   $E= 2.4129$   $p_x= 1.8484$   $p_y= -1.2351$   $p_z= 0.0000$   
 $\ell'$   $E= 1.9758$   $p_x= 0.0000$   $p_y= -1.9729$   $p_z= 0.0000$   
 $P'$   $E= 2.4127$   $p_x= 0.0000$   $p_y= 2.2229$   $p_z= 0.0000$   
 $q_\gamma$   $E= 0.2500$   $p_x= -0.0000$   $p_y= -0.2500$   $p_z= 0.0000$

Transverse plane




Leading initial-ensemble/final-state components (N=3, retained=96.4%)



# Tx proton characteristic max $C_{\mu\gamma}$ configuration

3D momenta




c\_e\_gamma\_low\_egamma\_tx\_proton\_region\_2  $C_{\mu\gamma}=0.0251841$   
 region: low\_Egamma\_Tx\_proton\_region\_2 final-pair delta\_xy=0 rad  
 outgoing-state purity: 0.506527  
 kinematic point: high\_s\_high\_theta\_in\_low\_Egamma  
 incoming state:  $\frac{1}{2}\sum_{h_\mu}\rho(h_\mu, T_{x,p})$

$s=21.5158$ ,  $\sqrt{s}=4.63852$ ,  $|\vec{P}|=2.22311$ ,  $|\vec{P}'|=2.22295$   
 $\theta_{in}=1.57079$ ,  $\phi_l=0.589049$ ,  $\phi_p=3.73064$   
 $E_\gamma=0.25$ ,  $\phi_\gamma=4.71239$   
 $Q^2=13.6459$ ,  $x_B=0.854805$ ,  $t=-15.3748$ ,  $W^2=3.19769$ ,  $y=0.774004$   
 $\theta(l', \gamma)=0$  deg

four-momenta  $[E, p_x, p_y, p_z]$  GeV:  
 $l$   $E= 2.2256$   $p_x= 1.8484$   $p_y= 1.2351$   $p_z= -0.0000$   
 $P$   $E= 2.4129$   $p_x= -1.8484$   $p_y= -1.2351$   $p_z= 0.0000$   
 $l'$   $E= 1.9758$   $p_x= 0.0000$   $p_y= -1.9729$   $p_z= 0.0000$   
 $P'$   $E= 2.4127$   $p_x= 0.0000$   $p_y= 2.2229$   $p_z= 0.0000$   
 $q_\gamma$   $E= 0.2500$   $p_x= -0.0000$   $p_y= -0.2500$   $p_z= 0.0000$

Transverse plane



Leading initial-ensemble/final-state components (N=3, retained=96.4%)

