


# L massless lepton: $C_{\ell_0 p}$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 0.25 \text{ GeV}$ ,  $\max C_{\ell_0 p} = 0.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1 \text{ GeV}$ ,  $\max C_{\ell_0 p} = 0.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1.8 \text{ GeV}$ ,  $\max C_{\ell_0 p} = 0.000$




# L massless lepton: $C_{\ell_0\gamma}$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 0.25 \text{ GeV}$ ,  $\max C_{\ell_0\gamma} = 0.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1 \text{ GeV}$ ,  $\max C_{\ell_0\gamma} = 0.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1.8 \text{ GeV}$ ,  $\max C_{\ell_0\gamma} = 0.000$




# L massless lepton: $C_{p\gamma}$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 0.25 \text{ GeV}$ ,  $\max C_{p\gamma} = 0.495$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1 \text{ GeV}$ ,  $\max C_{p\gamma} = 0.617$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1.8 \text{ GeV}$ ,  $\max C_{p\gamma} = 0.681$




# L massless lepton: $C_{\ell_0|p\gamma}$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 0.25 \text{ GeV}$ ,  $\max C_{\ell_0|p\gamma} = 0.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1 \text{ GeV}$ ,  $\max C_{\ell_0|p\gamma} = 0.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1.8 \text{ GeV}$ ,  $\max C_{\ell_0|p\gamma} = 0.000$




# L massless lepton: $C_{p|\ell_0\gamma}$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 0.25 \text{ GeV}$ ,  $\max C_{p|\ell_0\gamma} = 0.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1 \text{ GeV}$ ,  $\max C_{p|\ell_0\gamma} = 0.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1.8 \text{ GeV}$ ,  $\max C_{p|\ell_0\gamma} = 0.000$




# L massless lepton: $C_{\gamma|\ell_0 p}$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 0.25 \text{ GeV}$ ,  $\max C_{\gamma|\ell_0 p} = 0.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1 \text{ GeV}$ ,  $\max C_{\gamma|\ell_0 p} = 0.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1.8 \text{ GeV}$ ,  $\max C_{\gamma|\ell_0 p} = 0.000$




# L massless lepton: $M_{\ell_0}$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 0.25 \text{ GeV}$ ,  $\max M_{\ell_0} = -0.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 1 \text{ GeV}$ ,  $\max M_{\ell_0} = -0.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 1.8 \text{ GeV}$ ,  $\max M_{\ell_0} = 0.000$




# L massless lepton: $M_p$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 0.25 \text{ GeV}$ ,  $\max M_p = -0.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1 \text{ GeV}$ ,  $\max M_p = -0.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1.8 \text{ GeV}$ ,  $\max M_p = 0.000$




# L massless lepton: $M_Y$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 0.25 \text{ GeV}$ ,  $\max M_Y = -0.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 1 \text{ GeV}$ ,  $\max M_Y = -0.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 1.8 \text{ GeV}$ ,  $\max M_Y = 0.000$




# L massless lepton: $F_3$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 0.25 \text{ GeV}$ ,  $\max F_3 = 0.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1 \text{ GeV}$ ,  $\max F_3 = 0.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1.8 \text{ GeV}$ ,  $\max F_3 = 0.000$




# L massless lepton: $P_{\text{GHZ}+}$ + two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 0.25 \text{ GeV}$ ,  $\max P_{\text{GHZ}+} = 0.280$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 1 \text{ GeV}$ ,  $\max P_{\text{GHZ}+} = 0.384$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 1.8 \text{ GeV}$ ,  $\max P_{\text{GHZ}+} = 0.483$



# L massless lepton: $P_W$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 0.25 \text{ GeV}$ ,  $\max P_W = 0.104$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 1 \text{ GeV}$ ,  $\max P_W = 0.132$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 1.8 \text{ GeV}$ ,  $\max P_W = 0.108$

