


# L muon + Ty proton: $C_{\mu p}$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 0.25 \text{ GeV}$ ,  $\max C_{\mu p} = 0.187$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1 \text{ GeV}$ ,  $\max C_{\mu p} = 0.366$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1.8 \text{ GeV}$ ,  $\max C_{\mu p} = 0.702$




# L muon + Ty proton: $C_{\mu\gamma}$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 0.25 \text{ GeV}$ ,  $\max C_{\mu\gamma} = 0.664$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1 \text{ GeV}$ ,  $\max C_{\mu\gamma} = 0.611$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1.8 \text{ GeV}$ ,  $\max C_{\mu\gamma} = 0.556$




# L muon + Ty proton: $C_{p\gamma}$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 0.25 \text{ GeV}$ ,  $\max C_{p\gamma} = 1.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1 \text{ GeV}$ ,  $\max C_{p\gamma} = 0.999$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1.8 \text{ GeV}$ ,  $\max C_{p\gamma} = 0.991$




# L muon + Ty proton: $C_{\mu|p\gamma}$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 0.25 \text{ GeV}$ ,  $\max C_{\mu|p\gamma} = 0.664$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1 \text{ GeV}$ ,  $\max C_{\mu|p\gamma} = 0.635$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1.8 \text{ GeV}$ ,  $\max C_{\mu|p\gamma} = 0.774$




# L muon + Ty proton: $C_{p|\mu\gamma}$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 0.25 \text{ GeV}$ ,  $\max C_{p|\mu\gamma} = 1.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1 \text{ GeV}$ ,  $\max C_{p|\mu\gamma} = 0.999$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1.8 \text{ GeV}$ ,  $\max C_{p|\mu\gamma} = 0.991$




# L muon + Ty proton: $C_{\gamma|\mu p}$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 0.25 \text{ GeV}$ ,  $\max C_{\gamma|\mu p} = 1.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1 \text{ GeV}$ ,  $\max C_{\gamma|\mu p} = 0.999$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1.8 \text{ GeV}$ ,  $\max C_{\gamma|\mu p} = 0.991$




# L muon + Ty proton: $M_\mu$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 0.25 \text{ GeV}$ ,  $\max M_\mu = 0.035$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1 \text{ GeV}$ ,  $\max M_\mu = 0.041$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1.8 \text{ GeV}$ ,  $\max M_\mu = 0.018$




# L muon + Ty proton: $M_p$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 0.25 \text{ GeV}$ ,  $\max M_p = 0.035$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 1 \text{ GeV}$ ,  $\max M_p = 0.041$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 1.8 \text{ GeV}$ ,  $\max M_p = 0.018$






# L muon + Ty proton: $M_Y$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 0.25 \text{ GeV}$ ,  $\max M_Y = 0.035$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 1 \text{ GeV}$ ,  $\max M_Y = 0.041$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 1.8 \text{ GeV}$ ,  $\max M_Y = 0.018$




# L muon + Ty proton: $F_3$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 0.25 \text{ GeV}$ ,  $\max F_3 = 0.231$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1 \text{ GeV}$ ,  $\max F_3 = 0.445$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1.8 \text{ GeV}$ ,  $\max F_3 = 0.658$




# L muon + Ty proton: $P_{\text{GHZ}+}$ + two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 0.25 \text{ GeV}$ ,  $\max P_{\text{GHZ}+} = 0.279$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 1 \text{ GeV}$ ,  $\max P_{\text{GHZ}+} = 0.383$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 1.8 \text{ GeV}$ ,  $\max P_{\text{GHZ}+} = 0.480$



# L muon + Ty proton: $P_W$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 0.25 \text{ GeV}$ ,  $\max P_W = 0.409$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 1 \text{ GeV}$ ,  $\max P_W = 0.342$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 1.8 \text{ GeV}$ ,  $\max P_W = 0.333$

