



high_Egamma L electron + Tx proton: max C_{ey} regions



L electron + Tx proton characteristic max $C_{e\gamma}$ configuration

3D momenta




c_e_gamma_high_egamma_ltx_region_0 $C_{e\gamma}=0.00350422$
 region: high_Egamma_LTx_region_0 final-pair delta_xy=0.504869 rad
 outgoing-state purity: 1
 kinematic point: high_s_high_theta_in_high_Egamma
 incoming state: $|L_e\rangle \otimes |T_{x,p}\rangle$

$s=21.5158$, $\sqrt{s}=4.63852$, $|\vec{P}|=2.22442$, $|\vec{P}'|=2.20096$
 $\theta_{in}=1.57079$, $\phi_l=4.71239$, $\phi_p=1.5708$
 $E_\gamma=1.8$, $\phi_\gamma=4.61421$
 $Q^2=0.161847$, $x_B=0.00971461$, $t=-8.37957e-05$, $W^2=17.3782$, $y=0.807336$
 $\theta(l', \gamma)=28.9269$ deg

four-momenta $[E, p_x, p_y, p_z]$ GeV:
 l $E= 2.2244$ $p_x= -0.0000$ $p_y= -2.2244$ $p_z= -0.0000$
 P $E= 2.4141$ $p_x= 0.0000$ $p_y= 2.2244$ $p_z= 0.0000$
 l' $E= 0.4460$ $p_x= 0.1764$ $p_y= -0.4096$ $p_z= 0.0000$
 P' $E= 2.3925$ $p_x= 0.0000$ $p_y= 2.2010$ $p_z= 0.0000$
 q_γ $E= 1.8000$ $p_x= -0.1764$ $p_y= -1.7913$ $p_z= 0.0000$

Transverse plane




$h_e=+1, h_p=-1, h_\gamma=+1$
 electron L, proton Tx

$h_e=+1, h_p=+1, h_\gamma=-1$
 electron L, proton Tx


$h_e=+1, h_p=-1, h_\gamma=-1$
 electron L, proton Tx

Leading initial-ensemble/final-state components (N=3, retained=100.0%)



L electron + Tx proton characteristic max $C_{e\gamma}$ configuration

3D momenta




c_e_gamma_high_egamma_ltx_region_1 C_{e\gamma}=0.00121272
region: high Egamma LTx region_1 final-pair delta_xy=2.48366 rad
outgoing-state purity: 1
kinematic point: high_s_high_theta_in_high_Egamma
incoming state: $|L_e\rangle \otimes |T_{x,p}\rangle$

$s=21.5158$, $\sqrt{s}=4.63852$, $|\vec{P}|=2.22442$, $|\vec{P}'|=1.10115$
 $\theta_{in}=1.57079$, $\phi_l=5.59596$, $\phi_p=2.45437$
 $E_\gamma=1.8$, $\phi_\gamma=5.59596$
 $Q^2=11.093$, $x_B=0.589575$, $t=-2.11653$, $W^2=8.60206$, $y=0.911766$
 $\theta(l', \gamma)=142.303$ deg

four-momenta $[E, p_x, p_y, p_z]$ GeV:
 l $E= 2.2244$ $p_x= 1.7195$ $p_y= -1.4112$ $p_z= -0.0000$
 P $E= 2.4141$ $p_x= -1.7195$ $p_y= 1.4112$ $p_z= 0.0000$
 l' $E= 1.3920$ $p_x= -1.3914$ $p_y= 0.0408$ $p_z= 0.0000$
 P' $E= 1.4465$ $p_x= 0.0000$ $p_y= 1.1011$ $p_z= 0.0000$
 q_γ $E= 1.8000$ $p_x= 1.3914$ $p_y= -1.1419$ $p_z= 0.0000$


Transverse plane



$h_e=-1, h_p=-1, h_\gamma=+1$
electron L, proton Tx


$h_e=-1, h_p=+1, h_\gamma=+1$
electron L, proton Tx

Leading initial-ensemble/final-state components (N=2, retained=100.0%)



L electron + Tx proton characteristic max $C_{e\gamma}$ configuration

3D momenta




c_e_gamma_high_egamma_ltx_region_2 C_{e\gamma}=0.00120734
region: high_Egamma_LTx_region_2 final-pair delta_xy=2.5801 rad
outgoing-state purity: 1
kinematic point: high_s_high_theta_in_high_Egamma
incoming state: $|L_e\rangle \otimes |T_{x,p}\rangle$

$s=21.5158$, $\sqrt{s}=4.63852$, $|\vec{P}|=2.22442$, $|\vec{P}'|=0.958776$
 $\theta_{in}=1.57079$, $\phi_l=3.73064$, $\phi_p=0.589049$
 $E_\gamma=1.8$, $\phi_\gamma=3.73064$
 $Q^2=12.299$, $x_B=0.645776$, $t=-2.34664$, $W^2=7.62614$, $y=0.922917$
 $\theta(l', \gamma)=147.829$ deg

four-momenta $[E, p_x, p_y, p_z]$ GeV:
 l $E= 2.2244$ $p_x= -1.8495$ $p_y= -1.2358$ $p_z= -0.0000$
 P $E= 2.4141$ $p_x= 1.8495$ $p_y= 1.2358$ $p_z= 0.0000$
 l' $E= 1.4972$ $p_x= 1.4966$ $p_y= 0.0413$ $p_z= 0.0000$
 P' $E= 1.3413$ $p_x= 0.0000$ $p_y= 0.9588$ $p_z= 0.0000$
 q_γ $E= 1.8000$ $p_x= -1.4966$ $p_y= -1.0000$ $p_z= 0.0000$


Transverse plane



$h_e=-1, h_p=-1, h_\gamma=+1$
electron L, proton Tx


$h_e=-1, h_p=+1, h_\gamma=+1$
electron L, proton Tx

Leading initial-ensemble/final-state components (N=2, retained=100.0%)



L electron + Tx proton characteristic max $C_{e\gamma}$ configuration

3D momenta




c_e_gamma_high_egamma_ltx_region_3 $C_{e\gamma}=0.000755772$
 region: high_Egamma_LTx_region_3 final-pair delta_xy=0.504869 rad
 outgoing-state purity: 1
 kinematic point: high_s_high_theta_in_high_Egamma
 incoming state: $|L_e\rangle \otimes |\bar{T}_{x,p}\rangle$

$s=21.5158$, $\sqrt{s}=4.63852$, $|\vec{P}|=2.22442$, $|\vec{P}'|=2.20096$
 $\theta_{in}=1.57079$, $\phi_l=0.490874$, $\phi_P=3.63247$
 $E_\gamma=1.8$, $\phi_\gamma=4.81056$
 $Q^2=3.53553$, $x_B=0.176477$, $t=-14.4076$, $W^2=17.3782$, $y=0.970821$
 $\theta(l', \gamma)=28.9269$ deg

four-momenta $[E, p_x, p_y, p_z]$ GeV:
 l $E= 2.2244$ $p_x= 1.9618$ $p_y= 1.0486$ $p_z= -0.0000$
 P $E= 2.4141$ $p_x= -1.9618$ $p_y= -1.0486$ $p_z= 0.0000$
 l' $E= 0.4460$ $p_x= -0.1764$ $p_y= -0.4096$ $p_z= 0.0000$
 P' $E= 2.3925$ $p_x= 0.0000$ $p_y= 2.2010$ $p_z= 0.0000$
 q_γ $E= 1.8000$ $p_x= 0.1764$ $p_y= -1.7913$ $p_z= 0.0000$

Transverse plane



Leading initial-ensemble/final-state components (N=3, retained=100.0%)

