



high\_Egamma L electron + Ty proton: max  $C_{ep}$  regions



L electron + Ty proton characteristic max  $C_{ep}$  configuration

3D momenta




c\_ep\_high\_egamma\_lty\_region\_0  $C_{ep}$ =0.0142742  
region: high\_Egamma\_LTy\_region\_0 final-pair delta\_xy=3.14159 rad  
outgoing-state purity: 1  
kinematic point: high\_s\_high\_theta\_in\_high\_Egamma  
incoming state:  $|L_e\rangle \otimes |T_{y,p}\rangle$

$s=21.5158$ ,  $\sqrt{s}=4.63852$ ,  $|\vec{P}|=2.22442$ ,  $|\vec{P}'|=0.0956529$   
 $\theta_{in}=1.57079$ ,  $\phi_\ell=1.5708$ ,  $\phi_p=4.71239$   
 $E_\gamma=1.8$ ,  $\phi_\gamma=1.5708$   
 $Q^2=16.8669$ ,  $x_B=0.846865$ ,  $t=-3.21819$ ,  $W^2=3.92981$ ,  $y=0.965151$   
 $\theta(\ell',\gamma)=180$  deg


four-momenta  $[E, p_x, p_y, p_z]$  GeV:  
 $\ell$   $E= 2.2244$   $p_x= 0.0000$   $p_y= 2.2244$   $p_z= -0.0000$   
 $P$   $E= 2.4141$   $p_x= -0.0000$   $p_y= -2.2244$   $p_z= 0.0000$   
 $\ell'$   $E= 1.8957$   $p_x= -0.0000$   $p_y= -1.8957$   $p_z= 0.0000$   
 $P'$   $E= 0.9429$   $p_x= 0.0000$   $p_y= 0.0957$   $p_z= 0.0000$   
 $q_\gamma$   $E= 1.8000$   $p_x= 0.0000$   $p_y= 1.8000$   $p_z= 0.0000$

Transverse plane




$h_e=-1$ ,  $h_p=-1$ ,  $h_\gamma=+1$   
electron L, proton Ty

Leading initial-ensemble/final-state components (N=1, retained=100.0%)



# L electron + Ty proton characteristic max $C_{ep}$ configuration

3D momenta




c\_ep\_high\_egamma\_lty\_region\_1 C\_{ep}=0.0139936  
 region: high\_Egamma\_LTy\_region\_1 final-pair delta\_xy=2.22659 rad  
 outgoing-state purity: 1  
 kinematic point: high\_s\_high\_theta\_in\_high\_Egamma  
 incoming state:  $|L_e\rangle \otimes |T_{y,p}\rangle$

$s=21.5158$ ,  $\sqrt{s}=4.63852$ ,  $|\vec{P}|=2.22442$ ,  $|\vec{P}'|=0.151482$   
 $\theta_{in}=1.57079$ ,  $\phi_l=0.589049$ ,  $\phi_p=3.73064$   
 $E_Y=1.8$ ,  $\phi_Y=0.589049$   
 $Q^2=16.7833$ ,  $x_B=0.843345$ ,  $t=-3.20225$ ,  $W^2=3.99742$ ,  $y=0.964378$   
 $\theta(l', \gamma)=176.176$  deg

four-momenta  $[E, p_x, p_y, p_z]$  GeV:  
 $l$   $E= 2.2244$   $p_x= 1.8495$   $p_y= 1.2358$   $p_z= -0.0000$   
 $P$   $E= 2.4141$   $p_x= -1.8495$   $p_y= -1.2358$   $p_z= 0.0000$   
 $l'$   $E= 1.8884$   $p_x= -1.4966$   $p_y= -1.1515$   $p_z= 0.0000$   
 $P'$   $E= 0.9502$   $p_x= 0.0000$   $p_y= 0.1515$   $p_z= 0.0000$   
 $q_Y$   $E= 1.8000$   $p_x= 1.4966$   $p_y= 1.0000$   $p_z= 0.0000$


Transverse plane



$h_e=-1, h_p=-1, h_Y=+1$   
 electron L, proton Ty


$h_e=-1, h_p=+1, h_Y=+1$   
 electron L, proton Ty

Leading initial-ensemble/final-state components (N=2, retained=100.0%)



# L electron + Ty proton characteristic max $C_{ep}$ configuration

3D momenta




c\_ep\_high\_egamma\_lty\_region\_2 C\_{ep}=0.0139936  
 region: high\_Egamma\_LTy\_region\_2 final-pair delta\_xy=2.22659 rad  
 outgoing-state purity: 1  
 kinematic point: high\_s\_high\_theta\_in\_high\_Egamma  
 incoming state:  $|L_e\rangle \otimes |T_{y,p}\rangle$

$s=21.5158$ ,  $\sqrt{s}=4.63852$ ,  $|\vec{P}|=2.22442$ ,  $|\vec{P}'|=0.151482$   
 $\theta_{in}=1.57079$ ,  $\phi_l=2.55254$ ,  $\phi_p=5.69414$   
 $E_\gamma=1.8$ ,  $\phi_\gamma=2.55254$   
 $Q^2=16.7833$ ,  $x_B=0.843345$ ,  $t=-3.20225$ ,  $W^2=3.99742$ ,  $y=0.964378$   
 $\theta(l', \gamma)=176.176$  deg

four-momenta  $[E, p_x, p_y, p_z]$  GeV:

$l$   $E= 2.2244$   $p_x= -1.8495$   $p_y= 1.2358$   $p_z= -0.0000$   
 $P$   $E= 2.4141$   $p_x= 1.8495$   $p_y= -1.2358$   $p_z= 0.0000$   
 $l'$   $E= 1.8884$   $p_x= 1.4966$   $p_y= -1.1515$   $p_z= 0.0000$   
 $P'$   $E= 0.9502$   $p_x= 0.0000$   $p_y= 0.1515$   $p_z= 0.0000$   
 $q_\gamma$   $E= 1.8000$   $p_x= -1.4966$   $p_y= 1.0000$   $p_z= 0.0000$


Transverse plane



$h_e=-1, h_p=-1, h_\gamma=+1$   
 electron L, proton Ty


$h_e=-1, h_p=+1, h_\gamma=+1$   
 electron L, proton Ty

Leading initial-ensemble/final-state components (N=2, retained=100.0%)



# L electron + Ty proton characteristic max $C_{ep}$ configuration

3D momenta




c\_ep\_high\_egamma\_lty\_region\_3  $C_{ep}$ =0.0122091  
 region: high\_Egamma\_LTy\_region\_3 final-pair delta\_xy=1.58254 rad  
 outgoing-state purity: 1  
 kinematic point: high\_s\_high\_theta\_in\_high\_Egamma  
 incoming state:  $|L_e\rangle \otimes |T_{\gamma,p}\rangle$

$s=21.5158$ ,  $\sqrt{s}=4.63852$ ,  $|\vec{P}|=2.22442$ ,  $|\vec{P}'|=0.708355$   
 $\theta_{in}=1.57079$ ,  $\phi_l=5.89049$ ,  $\phi_p=2.74889$   
 $E_\gamma=1.8$ ,  $\phi_\gamma=5.89049$   
 $Q^2=14.2008$ ,  $x_B=0.731691$ ,  $t=-2.7095$ ,  $W^2=6.08723$ ,  $y=0.9405$   
 $\theta(l', \gamma)=156.827$  deg

four-momenta  $[E, p_x, p_y, p_z]$  GeV:  
 $l$   $E= 2.2244$   $p_x= 2.0551$   $p_y= -0.8512$   $p_z= -0.0000$   
 $P$   $E= 2.4141$   $p_x= -2.0551$   $p_y= 0.8512$   $p_z= 0.0000$   
 $l'$   $E= 1.6631$   $p_x= -1.6630$   $p_y= -0.0195$   $p_z= 0.0000$   
 $P'$   $E= 1.1754$   $p_x= 0.0000$   $p_y= 0.7084$   $p_z= 0.0000$   
 $q_\gamma$   $E= 1.8000$   $p_x= 1.6630$   $p_y= -0.6888$   $p_z= 0.0000$

Transverse plane



$h_e=-1, h_p=-1, h_\gamma=+1$   
 electron L, proton Ty

$h_e=-1, h_p=+1, h_\gamma=+1$   
 electron L, proton Ty

Leading initial-ensemble/final-state components (N=2, retained=100.0%)

