



high_Egamma Tx electron + Tx proton: max C_{ey} regions



Tx electron + Tx proton characteristic max $C_{e\gamma}$ configuration

3D momenta




c_e_gamma_high_egamma_txtx_region_0 $C_{e\gamma}=0.999979$
 region: high_Egamma_TxTx_region_0 final-pair delta_xy=8.88178e-16 rad
 outgoing-state purity: 1
 kinematic point: high_s_high_theta_in_high_Egamma
 incoming state: $|T_{x,e}\rangle \otimes |T_{x,p}\rangle$

$s=21.5158$, $\sqrt{s}=4.63852$, $|\vec{P}|=2.22442$, $|\vec{P}'|=2.22442$
 $\theta_{in}=1.57079$, $\phi_l=4.61421$, $\phi_P=1.47262$
 $E_\gamma=1.8$, $\phi_\gamma=4.71239$
 $Q^2=0.00909292$, $x_B=0.000544233$, $t=-0.0476523$, $W^2=17.5785$, $y=0.809641$
 $\theta(l', \gamma)=0$ deg

four-momenta $[E, p_x, p_y, p_z]$ GeV:
 l $E= 2.2244$ $p_x= -0.2180$ $p_y= -2.2137$ $p_z= -0.0000$
 P $E= 2.4141$ $p_x= 0.2180$ $p_y= 2.2137$ $p_z= 0.0000$
 l' $E= 0.4244$ $p_x= 0.0000$ $p_y= -0.4244$ $p_z= 0.0000$
 P' $E= 2.4141$ $p_x= 0.0000$ $p_y= 2.2244$ $p_z= 0.0000$
 q_γ $E= 1.8000$ $p_x= -0.0000$ $p_y= -1.8000$ $p_z= 0.0000$

Transverse plane



$h_e=-1, h_p=+1, h_\gamma=+1$
 electron Tx, proton Tx

$h_e=+1, h_p=+1, h_\gamma=-1$
 electron Tx, proton Tx

$h_e=+1, h_p=-1, h_\gamma=-1$
 electron Tx, proton Tx

$h_e=-1, h_p=-1, h_\gamma=+1$
 electron Tx, proton Tx

Leading initial-ensemble/final-state components (N=4, retained=100.0%)

