



gamma L auxiliary lepton + Ty proton: max  $C_{\ell p}$  regions



# L auxiliary lepton + Ty proton characteristic max $C_{lp}$ configuration

3D momenta




c\_aux\_p\_low\_egamma\_lty\_region\_0 C\_{l\_auxp}=0.724665  
 region: low\_Egamma\_LTy\_region\_0 final-pair delta\_xy=3.03969 rad  
 outgoing-state purity: 1  
 kinematic point: high\_s\_high\_theta\_in\_low\_Egamma  
 incoming state:  $|L_\ell\rangle \otimes |\bar{T}_{y,p}\rangle$

$s=21.5158$ ,  $\sqrt{s}=4.63852$ ,  $|\vec{P}|=2.10694$ ,  $|\vec{P}'|=1.89395$   
 $\theta_{in}=1.57079$ ,  $\phi_\ell=2.65072$ ,  $\phi_P=5.79231$   
 $E_\gamma=0.25$ ,  $\phi_\gamma=2.55254$   
 $Q^2=13.4222$ ,  $x_B=0.961969$ ,  $t=-11.7513$ ,  $W^2=1.41048$ ,  $y=0.710574$   
 $\theta(\ell', \gamma)=129.588$  deg


four-momenta  $[E, p_x, p_y, p_z]$  GeV:

|            |             |                |                |                |
|------------|-------------|----------------|----------------|----------------|
| $\ell$     | $E= 2.3322$ | $p_x= -1.8582$ | $p_y= 0.9932$  | $p_z= -0.0000$ |
| $P$        | $E= 2.3063$ | $p_x= 1.8582$  | $p_y= -0.9932$ | $p_z= 0.0000$  |
| $\ell'$    | $E= 2.2750$ | $p_x= 0.2079$  | $p_y= -2.0328$ | $p_z= 0.0000$  |
| $P'$       | $E= 2.1135$ | $p_x= 0.0000$  | $p_y= 1.8940$  | $p_z= 0.0000$  |
| $q_\gamma$ | $E= 0.2500$ | $p_x= -0.2079$ | $p_y= 0.1389$  | $p_z= 0.0000$  |

Transverse plane




Leading initial-ensemble/final-state components (N=4, retained=97.8%)



# L auxiliary lepton + Ty proton characteristic max $C_{lp}$ configuration

3D momenta




c\_aux\_p\_low\_egamma\_lty\_region\_1  $C_{\{l\_aux\}}=0.724665$   
 region: low\_Egamma\_LTy\_region\_1 final-pair delta\_xy=3.03969 rad  
 outgoing-state purity: 1  
 kinematic point: high\_s\_high\_theta\_in\_low\_Egamma  
 incoming state:  $|L_l\rangle \otimes |\bar{T}_{y,p}\rangle$


$s=21.5158$ ,  $\sqrt{s}=4.63852$ ,  $|\vec{P}|=2.10694$ ,  $|\vec{P}'|=1.89395$   
 $\theta_{in}=1.57079$ ,  $\phi_l=0.490874$ ,  $\phi_P=3.63247$   
 $E_\gamma=0.25$ ,  $\phi_\gamma=0.589049$   
 $Q^2=13.4222$ ,  $x_B=0.961969$ ,  $t=-11.7513$ ,  $W^2=1.41048$ ,  $y=0.710574$   
 $\theta(l', \gamma)=129.588$  deg

four-momenta  $[E, p_x, p_y, p_z]$  GeV:  
 $l$   $E= 2.3322$   $p_x= 1.8582$   $p_y= 0.9932$   $p_z= -0.0000$   
 $P$   $E= 2.3063$   $p_x= -1.8582$   $p_y= -0.9932$   $p_z= 0.0000$   
 $l'$   $E= 2.2750$   $p_x= -0.2079$   $p_y= -2.0328$   $p_z= 0.0000$   
 $P'$   $E= 2.1135$   $p_x= 0.0000$   $p_y= 1.8940$   $p_z= 0.0000$   
 $q_\gamma$   $E= 0.2500$   $p_x= 0.2079$   $p_y= 0.1389$   $p_z= 0.0000$

Transverse plane




Leading initial-ensemble/final-state components (N=4, retained=97.8%)



# L auxiliary lepton + Ty proton characteristic max $C_{lp}$ configuration

3D momenta




c\_aux\_p\_low\_egamma\_lty\_region\_2 C\_{l\_auxp}=0.53125  
 region: low\_Egamma\_LTy\_region\_2 final-pair delta\_xy=3.14159 rad  
 outgoing-state purity: 1  
 kinematic point: high\_s\_high\_theta\_in\_low\_Egamma  
 incoming state:  $|L_\ell\rangle \otimes |\bar{T}_{y,p}\rangle$

$s=21.5158$ ,  $\sqrt{s}=4.63852$ ,  $|\vec{P}|=2.10694$ ,  $|\vec{P}'|=1.84352$   
 $\theta_{in}=1.57079$ ,  $\phi_\ell=1.66897$ ,  $\phi_p=4.81056$   
 $E_\gamma=0.25$ ,  $\phi_\gamma=1.5708$   
 $Q^2=17.6012$ ,  $x_B=0.993651$ ,  $t=-15.5121$ ,  $W^2=0.9923$ ,  $y=0.902103$   
 $\theta(\ell', \gamma)=180$  deg

four-momenta  $[E, p_x, p_y, p_z]$  GeV:  
 $l$   $E= 2.3322$   $p_x= -0.2065$   $p_y= 2.0968$   $p_z= -0.0000$   
 $P$   $E= 2.3063$   $p_x= 0.2065$   $p_y= -2.0968$   $p_z= 0.0000$   
 $l'$   $E= 2.3201$   $p_x= -0.0000$   $p_y= -2.0935$   $p_z= 0.0000$   
 $P'$   $E= 2.0684$   $p_x= 0.0000$   $p_y= 1.8435$   $p_z= 0.0000$   
 $q_\gamma$   $E= 0.2500$   $p_x= 0.0000$   $p_y= 0.2500$   $p_z= 0.0000$

Transverse plane



Leading initial-ensemble/final-state components (N=3, retained=100.0%)

