



low_Egamma L massless lepton + Tx proton: max $C_{p\gamma}$ regions



L massless lepton + Tx proton characteristic max $C_{p\gamma}$ configuration

3D momenta




c_p_gamma_low_egamma_ltx_region_0 C_{p\gamma}=0.999987
 region: low_Egamma_LTx_region_0 final-pair delta_xy=0 rad
 outgoing-state purity: 1
 kinematic point: high_s_high_theta_in_low_Egamma
 incoming state: $|L_{\ell_0}\rangle \otimes |T_{x,p}\rangle$

$s=21.5158$, $\sqrt{s}=4.63852$, $|\vec{P}|=2.22442$, $|\vec{P}'|=1.96296$
 $\theta_{in}=1.57079$, $\phi_{\ell}=4.71239$, $\phi_p=1.5708$
 $E_{\gamma}=0.25$, $\phi_{\gamma}=1.5708$
 $Q^2=8.66557e-12$, $x_B=8.15204e-11$, $t=-0.0114583$, $W^2=0.986143$, $y=0.00515116$
 $\theta(\ell', \gamma)=180$ deg

four-momenta $[E, p_x, p_y, p_z]$ GeV:
 ℓ $E= 2.2244$ $p_x= -0.0000$ $p_y= -2.2244$ $p_z= -0.0000$
 P $E= 2.4141$ $p_x= 0.0000$ $p_y= 2.2244$ $p_z= 0.0000$
 ℓ' $E= 2.2130$ $p_x= -0.0000$ $p_y= -2.2130$ $p_z= 0.0000$
 P' $E= 2.1756$ $p_x= 0.0000$ $p_y= 1.9630$ $p_z= 0.0000$
 q_{γ} $E= 0.2500$ $p_x= 0.0000$ $p_y= 0.2500$ $p_z= 0.0000$


Transverse plane



$h_e=+1$, $h_p=-1$, $h_{\gamma}=+1$
 electron L, proton Tx


$h_e=+1$, $h_p=+1$, $h_{\gamma}=-1$
 electron L, proton Tx

Leading initial-ensemble/final-state components (N=2, retained=100.0%)



L massless lepton + Tx proton characteristic max $C_{p\gamma}$ configuration

3D momenta




c_p_gamma_low_egamma_ltx_region_1 C_{\gamma}=0.991282
region: low_Egamma_LTx_region_1 final-pair delta_xy=1.37445 rad
outgoing-state purity: 1
kinematic point: high_s_high_theta_in_low_Egamma
incoming state: $|L_{\ell_0}\rangle \otimes |T_{x,p}\rangle$

$s=21.5158$, $\sqrt{s}=4.63852$, $|\vec{P}|=2.22442$, $|\vec{P}'|=2.06107$
 $\theta_{in}=1.57079$, $\phi_l=4.71239$, $\phi_p=1.5708$
 $E_{\gamma}=0.25$, $\phi_{\gamma}=2.94524$
 $Q^2=0.0631736$, $x_B=0.0635316$, $t=-0.00429535$, $W^2=1.81103$, $y=0.0481859$
 $\theta(\ell', \gamma)=107.879$ deg

four-momenta $[E, p_x, p_y, p_z]$ GeV:
 ℓ $E= 2.2244$ $p_x= -0.0000$ $p_y= -2.2244$ $p_z= -0.0000$
 P $E= 2.4141$ $p_x= 0.0000$ $p_y= 2.2244$ $p_z= 0.0000$
 ℓ' $E= 2.1240$ $p_x= 0.2452$ $p_y= -2.1098$ $p_z= 0.0000$
 P' $E= 2.2645$ $p_x= 0.0000$ $p_y= 2.0611$ $p_z= 0.0000$
 q_{γ} $E= 0.2500$ $p_x= -0.2452$ $p_y= 0.0488$ $p_z= 0.0000$


Transverse plane



$h_e=+1, h_p=+1, h_{\gamma}=-1$
electron L, proton Tx


$h_e=+1, h_p=-1, h_{\gamma}=+1$
electron L, proton Tx

Leading initial-ensemble/final-state components (N=2, retained=99.4%)



L massless lepton + Tx proton characteristic max $C_{p\gamma}$ configuration

3D momenta




c_p_gamma_low_egamma_ltx_region_2 C_{p\gamma}=0.975579
 region: low_Egamma LTx_region_2 final-pair delta_xy=1.37445 rad
 outgoing-state purity: 1
 kinematic point: high_s_high_theta_in_low_Egamma
 incoming state: $|L_{\ell_0}\rangle \otimes |T_{x,p}\rangle$

$s=21.5158$, $\sqrt{s}=4.63852$, $|\vec{P}|=2.22442$, $|\vec{P}'|=2.06107$
 $\theta_{in}=1.57079$, $\phi_{\ell}=4.71239$, $\phi_p=1.5708$
 $E_{\gamma}=0.25$, $\phi_{\gamma}=0.19635$
 $Q^2=0.0631736$, $x_B=0.0635316$, $t=-0.00429535$, $W^2=1.81103$, $y=0.0481859$
 $\theta(\ell', \gamma)=107.879$ deg

four-momenta $[E, p_x, p_y, p_z]$ GeV:

ℓ $E= 2.2244$ $p_x= -0.0000$ $p_y= -2.2244$ $p_z= -0.0000$
 P $E= 2.4141$ $p_x= 0.0000$ $p_y= 2.2244$ $p_z= 0.0000$
 ℓ' $E= 2.1240$ $p_x= -0.2452$ $p_y= -2.1098$ $p_z= 0.0000$
 P' $E= 2.2645$ $p_x= 0.0000$ $p_y= 2.0611$ $p_z= 0.0000$
 q_{γ} $E= 0.2500$ $p_x= 0.2452$ $p_y= 0.0488$ $p_z= 0.0000$


Transverse plane



$h_e=+1, h_p=-1, h_{\gamma}=+1$
 electron L, proton Tx


$h_e=+1, h_p=+1, h_{\gamma}=-1$
 electron L, proton Tx

Leading initial-ensemble/final-state components (N=2, retained=99.2%)



L massless lepton + Tx proton characteristic max $C_{p\gamma}$ configuration

3D momenta




c_p_gamma_low_egamma_ltx_region_3 $C_{\{p\gamma\}}=0.799942$
 region: low_Egamma_LTx_region_3 final-pair delta_xy=2.74889 rad
 outgoing-state purity: 1
 kinematic point: high_s_high_theta_in_low_Egamma
 incoming state: $|L_{\ell_0}\rangle \otimes |T_{X,p}\rangle$

$s=21.5158$, $\sqrt{s}=4.63852$, $|\vec{P}|=2.22442$, $|\vec{P}'|=2.21331$
 $\theta_{in}=1.57079$, $\phi_l=4.71239$, $\phi_p=1.5708$
 $E_\gamma=0.25$, $\phi_\gamma=4.31969$
 $Q^2=0.0102647$, $x_B=0.00459348$, $t=-1.87039e-05$, $W^2=3.10419$, $y=0.108287$
 $\theta(\ell', \gamma)=25.263$ deg

four-momenta $[E, p_x, p_y, p_z]$ GeV:

ℓ $E= 2.2244$ $p_x= -0.0000$ $p_y= -2.2244$ $p_z= -0.0000$
 P $E= 2.4141$ $p_x= 0.0000$ $p_y= 2.2244$ $p_z= 0.0000$
 ℓ' $E= 1.9846$ $p_x= 0.0957$ $p_y= -1.9823$ $p_z= 0.0000$
 P' $E= 2.4039$ $p_x= 0.0000$ $p_y= 2.2133$ $p_z= 0.0000$
 q_γ $E= 0.2500$ $p_x= -0.0957$ $p_y= -0.2310$ $p_z= 0.0000$

Transverse plane



Leading initial-ensemble/final-state components (N=3, retained=99.9%)

