



high\_Egamma Tx proton: max  $C_{\ell_0\gamma}$  regions



# Tx proton characteristic max $C_{\ell_0\gamma}$ configuration

3D momenta




c\_massless\_gamma\_high\_egamma\_tx\_proton\_region\_0  $C_{\ell_0\gamma}=5.52142\text{e-}12$   
 region: high\_Egamma\_Tx\_proton\_region\_0 final-pair delta\_xy=2.07925 rad  
 outgoing-state purity: 0.511112  
 kinematic point: high\_s\_high\_theta\_in\_high\_Egamma  
 incoming state:  $\frac{1}{2}\sum_{h_{\ell_0}}\rho(h_{\ell_0}, T_{X,p})$


$s=21.5158$ ,  $\sqrt{s}=4.63852$ ,  $|\vec{P}|=2.22442$ ,  $|\vec{P}'|=1.57742$   
 $\theta_{\text{in}}=1.57079$ ,  $\phi_l=4.12334$ ,  $\phi_P=0.981748$   
 $E_\gamma=1.8$ ,  $\phi_\gamma=4.12334$   
 $Q^2=6.63637$ ,  $x_B=0.369408$ ,  $t=-1.26622$ ,  $W^2=12.2084$ ,  $y=0.87056$   
 $\theta(\ell', \gamma)=119.132$  deg

four-momenta  $[E, p_x, p_y, p_z]$  GeV:  
 $\ell$   $E= 2.2244$   $p_x= -1.2358$   $p_y= -1.8495$   $p_z= -0.0000$   
 $P$   $E= 2.4141$   $p_x= 1.2358$   $p_y= 1.8495$   $p_z= 0.0000$   
 $\ell'$   $E= 1.0033$   $p_x= 1.0000$   $p_y= -0.0808$   $p_z= 0.0000$   
 $P'$   $E= 1.8352$   $p_x= 0.0000$   $p_y= 1.5774$   $p_z= 0.0000$   
 $q_\gamma$   $E= 1.8000$   $p_x= -1.0000$   $p_y= -1.4966$   $p_z= 0.0000$

Transverse plane




Leading initial-ensemble/final-state components (N=4, retained=96.5%)



# Tx proton characteristic max $C_{\ell_0\gamma}$ configuration

3D momenta




c\_massless\_gamma\_high\_egamma\_tx\_proton\_region\_1  $C_{\ell_0\gamma}=4.9728e-12$   
 region: high\_Egamma\_Tx\_proton\_region\_1 final-pair delta\_xy=2.07925 rad  
 outgoing-state purity: 0.511112  
 kinematic point: high\_s\_high\_theta\_in\_high\_Egamma  
 incoming state:  $\frac{1}{2}\sum_{h_{\ell_0}}\rho(h_{\ell_0}, T_{X,p})$


$s=21.5158$ ,  $\sqrt{s}=4.63852$ ,  $|\vec{P}|=2.22442$ ,  $|\vec{P}'|=1.57742$   
 $\theta_{in}=1.57079$ ,  $\phi_l=5.30144$ ,  $\phi_P=2.15984$   
 $E_\gamma=1.8$ ,  $\phi_\gamma=5.30144$   
 $Q^2=6.63637$ ,  $x_B=0.369408$ ,  $t=-1.26622$ ,  $W^2=12.2084$ ,  $y=0.87056$   
 $\theta(\ell', \gamma)=119.132$  deg

four-momenta  $[E, p_x, p_y, p_z]$  GeV:  
 $\ell$   $E= 2.2244$   $p_x= 1.2358$   $p_y= -1.8495$   $p_z= -0.0000$   
 $P$   $E= 2.4141$   $p_x= -1.2358$   $p_y= 1.8495$   $p_z= 0.0000$   
 $\ell'$   $E= 1.0033$   $p_x= -1.0000$   $p_y= -0.0808$   $p_z= 0.0000$   
 $P'$   $E= 1.8352$   $p_x= 0.0000$   $p_y= 1.5774$   $p_z= 0.0000$   
 $q_\gamma$   $E= 1.8000$   $p_x= 1.0000$   $p_y= -1.4966$   $p_z= 0.0000$

Transverse plane




Leading initial-ensemble/final-state components (N=4, retained=96.5%)



# Tx proton characteristic max $C_{\ell_0\gamma}$ configuration

3D momenta




c\_massless\_gamma\_high\_egamma\_tx\_proton\_region\_2  $C_{\ell_0\gamma}=2.80609\text{e-}12$   
 region: high Egamma Tx proton region\_2 final-pair delta\_xy=3.11511 rad  
 outgoing-state purity: 0.500035  
 kinematic point: high\_s\_high\_theta\_in\_high\_Egamma  
 incoming state:  $\frac{1}{2}\sum_{h_{\ell_0}}\rho(h_{\ell_0},T_{x,p})$


$s=21.5158$ ,  $\sqrt{s}=4.63852$ ,  $|\vec{P}|=2.22442$ ,  $|\vec{P}'|=0.106402$   
 $\theta_{\text{in}}=1.57079$ ,  $\phi_l=2.06167$ ,  $\phi_P=5.20326$   
 $E_\gamma=1.8$ ,  $\phi_\gamma=2.06167$   
 $Q^2=16.8537$ ,  $x_B=0.84631$ ,  $t=-3.21568$ ,  $W^2=3.94048$ ,  $y=0.965029$   
 $\theta(\ell',\gamma)=178.483$  deg

four-momenta  $[E, p_x, p_y, p_z]$  GeV:  
 $\ell$   $E= 2.2244$   $p_x= -1.0486$   $p_y= 1.9618$   $p_z= -0.0000$   
 $P$   $E= 2.4141$   $p_x= 1.0486$   $p_y= -1.9618$   $p_z= 0.0000$   
 $\ell'$   $E= 1.8945$   $p_x= 0.8485$   $p_y= -1.6939$   $p_z= 0.0000$   
 $P'$   $E= 0.9440$   $p_x= 0.0000$   $p_y= 0.1064$   $p_z= 0.0000$   
 $q_\gamma$   $E= 1.8000$   $p_x= -0.8485$   $p_y= 1.5875$   $p_z= 0.0000$

Transverse plane




Leading initial-ensemble/final-state components (N=4, retained=96.5%)



# Tx proton characteristic max $C_{\ell_0\gamma}$ configuration

3D momenta




c\_massless\_gamma\_high\_egamma\_tx\_proton\_region\_3  $C_{\ell_0\gamma}=2.73204\text{e-}12$   
 region: high Egamma Tx\_proton\_region\_3 final-pair delta\_xy=3.10882 rad  
 outgoing-state purity: 0.500047  
 kinematic point: high\_s\_high\_theta\_in\_high\_Egamma  
 incoming state:  $\frac{1}{2}\sum_{h_{\ell_0}}\rho(h_{\ell_0},T_{x,p})$

$s=21.5158$ ,  $\sqrt{s}=4.63852$ ,  $|\vec{P}|=2.22442$ ,  $|\vec{P}'|=0.111698$   
 $\theta_{in}=1.57079$ ,  $\phi_\ell=0.981748$ ,  $\phi_P=4.12334$   
 $E_\gamma=1.8$ ,  $\phi_\gamma=0.981748$   
 $Q^2=16.8467$ ,  $x_B=0.846014$ ,  $t=-3.21434$ ,  $W^2=3.94616$ ,  $y=0.964964$   
 $\theta(\ell',\gamma)=178.122$  deg

four-momenta  $[E, p_x, p_y, p_z]$  GeV:  
 $\ell$   $E= 2.2244$   $p_x= 1.2358$   $p_y= 1.8495$   $p_z= -0.0000$   
 $P$   $E= 2.4141$   $p_x= -1.2358$   $p_y= -1.8495$   $p_z= 0.0000$   
 $\ell'$   $E= 1.8939$   $p_x= -1.0000$   $p_y= -1.6083$   $p_z= 0.0000$   
 $P'$   $E= 0.9446$   $p_x= 0.0000$   $p_y= 0.1117$   $p_z= 0.0000$   
 $q_\gamma$   $E= 1.8000$   $p_x= 1.0000$   $p_y= 1.4966$   $p_z= 0.0000$

Transverse plane



Leading initial-ensemble/final-state components (N=4, retained=96.5%)

