



low_Egamma Tx proton: max $C_{l_{0p}}$ regions



Tx proton characteristic max C_{l_0p} configuration

3D momenta




c_massless_p_low_egamma_tx_proton_region_0 $C_{l_0p}=2.74751e-11$
 region: low_Egamma_Tx_proton_region_0 final-pair delta_xy=3.0259 rad
 outgoing-state purity: 0.501046
 kinematic point: high_s_high_theta_in_low_Egamma
 incoming state: $\frac{1}{2}\sum_{h_{l_0}}\rho(h_{l_0}, T_{x,p})$

$s=21.5158$, $\sqrt{s}=4.63852$, $|\vec{P}|=2.22442$, $|\vec{P}'|=2.06107$
 $\theta_{in}=1.57079$, $\phi_l=0.19635$, $\phi_p=3.33794$
 $E_Y=0.25$, $\phi_Y=0.19635$
 $Q^2=12.3506$, $x_B=0.92989$, $t=-10.9625$, $W^2=1.81103$, $y=0.643621$
 $\theta(l', \gamma)=107.879$ deg

four-momenta $[E, p_x, p_y, p_z]$ GeV:


l $E= 2.2244$ $p_x= 2.1817$ $p_y= 0.4340$ $p_z= -0.0000$
 P $E= 2.4141$ $p_x= -2.1817$ $p_y= -0.4340$ $p_z= 0.0000$
 l' $E= 2.1240$ $p_x= -0.2452$ $p_y= -2.1098$ $p_z= 0.0000$
 P' $E= 2.2645$ $p_x= 0.0000$ $p_y= 2.0611$ $p_z= 0.0000$
 q_Y $E= 0.2500$ $p_x= 0.2452$ $p_y= 0.0488$ $p_z= 0.0000$

Transverse plane




$h_e=+1, h_p=+1, h_Y=+1$
 electron $h=+1$, proton Tx
 $h_e=-1, h_p=-1, h_Y=-1$
 electron $h=-1$, proton Tx
 $h_e=+1, h_p=+1, h_Y=-1$
 electron $h=+1$, proton Tx
 $h_e=-1, h_p=-1, h_Y=+1$
 electron $h=-1$, proton Tx
 $h_e=-1, h_p=+1, h_Y=-1$
 electron $h=-1$, proton Tx
 $h_e=-1, h_p=+1, h_Y=+1$
 electron $h=-1$, proton Tx
 $h_e=+1, h_p=-1, h_Y=+1$
 electron $h=+1$, proton Tx
 $h_e=+1, h_p=-1, h_Y=-1$
 electron $h=+1$, proton Tx

Leading initial-ensemble/final-state components (N=8, retained=100.0%)



Tx proton characteristic max $C_{l_0 p}$ configuration

3D momenta




c_massless_p_low_egamma_tx_proton_region_1 $C_{l_0 p}=2.11015e-11$
 region: low_Egamma_Tx_proton_region_1 final-pair delta_xy=3.10873 rad
 outgoing-state purity: 0.501997
 kinematic point: high_s_high_theta_in_low_Egamma
 incoming state: $\frac{1}{2} \sum_{h_{l_0}} \rho(h_{l_0}, T_{x,p})$


$s=21.5158$, $\sqrt{s}=4.63852$, $|\vec{P}|=2.22442$, $|\vec{P}'|=1.96799$
 $\theta_{in}=1.57079$, $\phi_l=1.27627$, $\phi_p=4.41786$
 $E_Y=0.25$, $\phi_Y=1.27627$
 $Q^2=19.3154$, $x_B=0.992374$, $t=-17.1445$, $W^2=1.02827$, $y=0.943196$
 $\theta(l', \gamma)=165.008$ deg

four-momenta $[E, p_x, p_y, p_z]$ GeV:
 l $E= 2.2244$ $p_x= 0.6457$ $p_y= 2.1286$ $p_z= -0.0000$
 P $E= 2.4141$ $p_x= -0.6457$ $p_y= -2.1286$ $p_z= 0.0000$
 l' $E= 2.2084$ $p_x= -0.0726$ $p_y= -2.2072$ $p_z= 0.0000$
 P' $E= 2.1801$ $p_x= 0.0000$ $p_y= 1.9680$ $p_z= 0.0000$
 q_Y $E= 0.2500$ $p_x= 0.0726$ $p_y= 0.2392$ $p_z= 0.0000$

Transverse plane




Leading initial-ensemble/final-state components (N=4, retained=99.8%)



Tx proton characteristic max $C_{\ell_0 p}$ configuration

3D momenta




c_massless_p_low_egamma_tx_proton_region_2 $C_{\ell_0 p}=5.99276e-12$
 region: low_Egamma_Tx_proton_region_2 final-pair delta_xy=3.05257 rad
 outgoing-state purity: 0.506922
 kinematic point: high_s_high_theta_in_low_Egamma
 incoming state: $\frac{1}{2} \sum_{h_{\ell_0}} \rho(h_{\ell_0}, T_{x,p})$

$s=21.5158$, $\sqrt{s}=4.63852$, $|\vec{P}|=2.22442$, $|\vec{P}'|=2.00644$
 $\theta_{in}=1.57079$, $\phi_{\ell}=2.45437$, $\phi_P=5.59596$
 $E_Y=0.25$, $\phi_Y=2.45437$
 $Q^2=16.4452$, $x_B=0.972157$, $t=-14.5969$, $W^2=1.35084$, $y=0.819743$
 $\theta(\ell', \gamma)=134.476$ deg


four-momenta $[E, p_x, p_y, p_z]$ GeV:

ℓ $E= 2.2244$ $p_x= -1.7195$ $p_y= 1.4112$ $p_z= -0.0000$
 P $E= 2.4141$ $p_x= 1.7195$ $p_y= -1.4112$ $p_z= 0.0000$
 ℓ' $E= 2.1736$ $p_x= 0.1933$ $p_y= -2.1650$ $p_z= 0.0000$
 P' $E= 2.2149$ $p_x= 0.0000$ $p_y= 2.0064$ $p_z= 0.0000$
 q_Y $E= 0.2500$ $p_x= -0.1933$ $p_y= 0.1586$ $p_z= 0.0000$

Transverse plane




Leading initial-ensemble/final-state components (N=4, retained=96.7%)



Tx proton characteristic max $C_{\ell_0 p}$ configuration

3D momenta




c_massless_p_low_egamma_tx_proton_region_3 $C_{\ell_0 p}=4.21072e-12$
 region: low_Egamma_Tx_proton_region_3 final-pair delta_xy=3.03348 rad
 outgoing-state purity: 0.503265
 kinematic point: high_s_high_theta_in_low_Egamma
 incoming state: $\frac{1}{2} \sum_{h_{l_0}} \rho(h_{l_0}, T_{X,p})$

$s=21.5158$, $\sqrt{s}=4.63852$, $|\vec{P}|=2.22442$, $|\vec{P}'|=2.14934$
 $\theta_{in}=1.57079$, $\phi_l=3.63247$, $\phi_P=0.490874$
 $E_\gamma=0.25$, $\phi_\gamma=3.63247$
 $Q^2=5.69552$, $x_B=0.772309$, $t=-5.05541$, $W^2=2.55899$, $y=0.357369$
 $\theta(\ell', \gamma)=68.0691$ deg

four-momenta $[E, p_x, p_y, p_z]$ GeV:

ℓ $E= 2.2244$ $p_x= -1.9618$ $p_y= -1.0486$ $p_z= -0.0000$
 P $E= 2.4141$ $p_x= 1.9618$ $p_y= 1.0486$ $p_z= 0.0000$
 ℓ' $E= 2.0434$ $p_x= 0.2205$ $p_y= -2.0315$ $p_z= 0.0000$
 P' $E= 2.3451$ $p_x= 0.0000$ $p_y= 2.1493$ $p_z= 0.0000$
 q_γ $E= 0.2500$ $p_x= -0.2205$ $p_y= -0.1178$ $p_z= 0.0000$

Transverse plane



Leading initial-ensemble/final-state components (N=8, retained=100.0%)

