



mid_Egamma L muon + Ty proton: max $C_{p\gamma}$ regions



L muon + Ty proton characteristic max $C_{p\gamma}$ configuration

3D momenta




c_p_gamma_mid_egamma_lty_region_0 C_{p\gamma}=0.999459
 region: mid_Egamma LTy_region_0 final-pair delta_xy=0 rad
 outgoing-state purity: 1
 kinematic point: high_s_high_theta_in_mid_Egamma
 incoming state: $|L_\mu\rangle \otimes |T_{\gamma,p}\rangle$

$s=21.5158$, $\sqrt{s}=4.63852$, $|\vec{P}|=2.22311$, $|\vec{P}'|=1.15107$
 $\theta_{in}=1.57079$, $\phi_l=4.71239$, $\phi_p=1.5708$
 $E_\gamma=1$, $\phi_\gamma=1.5708$
 $Q^2=1.20887e-05$, $x_B=1.81079e-05$, $t=-0.288022$, $W^2=1.54742$, $y=0.0323683$
 $\theta(l',\gamma)=180$ deg

four-momenta $[E, p_x, p_y, p_z]$ GeV:
 l $E= 2.2256$ $p_x= -0.0000$ $p_y= -2.2231$ $p_z= -0.0000$
 P $E= 2.4129$ $p_x= 0.0000$ $p_y= 2.2231$ $p_z= 0.0000$
 l' $E= 2.1537$ $p_x= -0.0000$ $p_y= -2.1511$ $p_z= 0.0000$
 P' $E= 1.4849$ $p_x= 0.0000$ $p_y= 1.1511$ $p_z= 0.0000$
 q_γ $E= 1.0000$ $p_x= 0.0000$ $p_y= 1.0000$ $p_z= 0.0000$


Transverse plane



$h_e=+1, h_p=-1, h_\gamma=+1$
 electron L, proton Ty


$h_e=+1, h_p=+1, h_\gamma=-1$
 electron L, proton Ty

Leading initial-ensemble/final-state components (N=2, retained=100.0%)



L muon + Ty proton characteristic max $C_{p\gamma}$ configuration

3D momenta




c_p_gamma_mid_egamma_lty_region_1 $C_{\{p\gamma\}}=0.971144$
 region: mid_Egamma_LTy_region_1 final-pair delta_xy=1.37445 rad
 outgoing-state purity: 1
 kinematic point: high_s_high_theta_in_mid_Egamma
 incoming state: $|L_\mu\rangle \otimes |T_{y,p}\rangle$

$s=21.5158$, $\sqrt{s}=4.63852$, $|\vec{P}|=2.22311$, $|\vec{P}'|=1.4461$
 $\theta_{in}=1.57079$, $\phi_t=4.71239$, $\phi_p=1.5708$
 $E_\gamma=1$, $\phi_\gamma=2.94524$
 $Q^2=1.20398$, $x_B=0.294582$, $t=-0.128717$, $W^2=3.76295$, $y=0.198164$
 $\theta(l', \gamma)=132.113$ deg

four-momenta $[E, p_x, p_y, p_z]$ GeV:

l $E= 2.2256$ $p_x= -0.0000$ $p_y= -2.2231$ $p_z= -0.0000$
 P $E= 2.4129$ $p_x= 0.0000$ $p_y= 2.2231$ $p_z= 0.0000$
 l' $E= 1.9148$ $p_x= 0.9808$ $p_y= -1.6412$ $p_z= 0.0000$
 P' $E= 1.7237$ $p_x= 0.0000$ $p_y= 1.4461$ $p_z= 0.0000$
 q_γ $E= 1.0000$ $p_x= -0.9808$ $p_y= 0.1951$ $p_z= 0.0000$


Transverse plane



$h_e=+1, h_p=-1, h_\gamma=+1$
 electron L, proton Ty


$h_e=+1, h_p=+1, h_\gamma=-1$
 electron L, proton Ty

Leading initial-ensemble/final-state components (N=2, retained=99.4%)



L muon + Ty proton characteristic max $C_{p\gamma}$ configuration

3D momenta




c_p_gamma_mid_egamma_lty_region_2 C_{p\gamma}=0.971144
 region: mid_Egamma LTy_region_2 final-pair delta_xy=1.37445 rad
 outgoing-state purity: 1
 kinematic point: high_s_high_theta_in_mid_Egamma
 incoming state: $|L_\mu\rangle \otimes |T_{\gamma,p}\rangle$

$s=21.5158$, $\sqrt{s}=4.63852$, $|\vec{P}|=2.22311$, $|\vec{P}'|=1.4461$
 $\theta_{in}=1.57079$, $\phi_l=4.71239$, $\phi_p=1.5708$
 $E_\gamma=1$, $\phi_\gamma=0.19635$
 $Q^2=1.20398$, $x_B=0.294582$, $t=-0.128717$, $W^2=3.76295$, $y=0.198164$
 $\theta(l', \gamma)=132.113$ deg

four-momenta $[E, p_x, p_y, p_z]$ GeV:

ℓ $E= 2.2256$ $p_x= -0.0000$ $p_y= -2.2231$ $p_z= -0.0000$
 P $E= 2.4129$ $p_x= 0.0000$ $p_y= 2.2231$ $p_z= 0.0000$
 ℓ' $E= 1.9148$ $p_x= -0.9808$ $p_y= -1.6412$ $p_z= 0.0000$
 P' $E= 1.7237$ $p_x= 0.0000$ $p_y= 1.4461$ $p_z= 0.0000$
 q_γ $E= 1.0000$ $p_x= 0.9808$ $p_y= 0.1951$ $p_z= 0.0000$

Transverse plane



$h_e=+1$, $h_p=-1$, $h_\gamma=+1$
 electron L, proton Ty

$h_e=+1$, $h_p=+1$, $h_\gamma=-1$
 electron L, proton Ty

Leading initial-ensemble/final-state components (N=2, retained=99.4%)

