



high_Egamma Tx muon + Ty proton: max $C_{p\gamma}$ regions



Tx muon + Ty proton characteristic max $C_{p\gamma}$ configuration

3D momenta




c_p_gamma_high_egamma_txtty_region_0 $C_{\{p\}\gamma}=0.98725$
 region: high_Egamma_TxTy_region_0 final-pair delta_xy=0 rad
 outgoing-state purity: 1
 kinematic point: high_s_high_theta_in_high_Egamma
 incoming state: $|T_{x,\mu}\rangle \otimes |\bar{T}_{y,p}\rangle$


$s=21.5158$, $\sqrt{s}=4.63852$, $|\vec{P}|=2.22311$, $|\vec{P}'|=0.0929745$
 $\theta_{in}=1.57079$, $\phi_l=4.71239$, $\phi_P=1.5708$
 $E_\gamma=1.8$, $\phi_\gamma=1.5708$
 $Q^2=0.000288358$, $x_B=9.42678e-05$, $t=-2.3757$, $W^2=3.93848$, $y=0.148313$
 $\theta(l', \gamma)=180$ deg

four-momenta $[E, p_x, p_y, p_z]$ GeV:
 l $E= 2.2256$ $p_x= -0.0000$ $p_y= -2.2231$ $p_z= -0.0000$
 P $E= 2.4129$ $p_x= 0.0000$ $p_y= 2.2231$ $p_z= 0.0000$
 l' $E= 1.8959$ $p_x= -0.0000$ $p_y= -1.8930$ $p_z= 0.0000$
 P' $E= 0.9426$ $p_x= 0.0000$ $p_y= 0.0930$ $p_z= 0.0000$
 q_γ $E= 1.8000$ $p_x= 0.0000$ $p_y= 1.8000$ $p_z= 0.0000$

Transverse plane




Leading initial-ensemble/final-state components (N=4, retained=100.0%)



Tx muon + Ty proton characteristic max $C_{p\gamma}$ configuration

3D momenta




c_p_gamma_high_egamma_txty_region_1 C_{p\gamma}=0.986466
 region: high_Egamma_TxTy_region_1 final-pair delta_xy=0.981748 rad
 outgoing-state purity: 1
 kinematic point: high_s_high_theta_in_high_Egamma
 incoming state: $|T_{x,\mu}\rangle \otimes |T_{y,p}\rangle$

$s=21.5158$, $\sqrt{s}=4.63852$, $|\vec{P}|=2.22311$, $|\vec{P}'|=0.147625$
 $\theta_{in}=1.57079$, $\phi_t=5.59596$, $\phi_p=2.45437$
 $E_\gamma=1.8$, $\phi_\gamma=2.55254$
 $Q^2=0.00488027$, $x_B=0.00156019$, $t=-2.40622$, $W^2=4.00295$, $y=0.151661$
 $\theta(\ell', \gamma)=176.268$ deg

four-momenta $[E, p_x, p_y, p_z]$ GeV:


ℓ $E= 2.2256$ $p_x= 1.7185$ $p_y= -1.4103$ $p_z= -0.0000$
 P $E= 2.4129$ $p_x= -1.7185$ $p_y= 1.4103$ $p_z= 0.0000$
 ℓ' $E= 1.8890$ $p_x= 1.4966$ $p_y= -1.1477$ $p_z= 0.0000$
 P' $E= 0.9495$ $p_x= 0.0000$ $p_y= 0.1476$ $p_z= 0.0000$
 q_γ $E= 1.8000$ $p_x= -1.4966$ $p_y= 1.0000$ $p_z= 0.0000$

Transverse plane




$h_e=-1, h_p=+1, h_\gamma=-1$
 electron Tx, proton Ty
 $h_e=+1, h_p=-1, h_\gamma=+1$
 electron Tx, proton Ty
 $h_e=+1, h_p=+1, h_\gamma=-1$
 electron Tx, proton Ty
 $h_e=-1, h_p=-1, h_\gamma=+1$
 electron Tx, proton Ty
 $h_e=-1, h_p=-1, h_\gamma=-1$
 electron Tx, proton Ty
 $h_e=+1, h_p=+1, h_\gamma=+1$
 electron Tx, proton Ty
 $h_e=+1, h_p=-1, h_\gamma=-1$
 electron Tx, proton Ty
 $h_e=-1, h_p=+1, h_\gamma=+1$
 electron Tx, proton Ty

Leading initial-ensemble/final-state components (N=8, retained=100.0%)



Tx muon + Ty proton characteristic max $C_{p\gamma}$ configuration

3D momenta




c_p_gamma_high_egamma_txtty_region_2 C_{p\gamma}=0.986456
 region: high_Egamma_TxTy_region_2 final-pair delta_xy=0.981748 rad
 outgoing-state purity: 1
 kinematic point: high_s_high_theta_in_high_Egamma
 incoming state: $|T_{x,\mu}\rangle \otimes |\bar{T}_{y,p}\rangle$

$s=21.5158$, $\sqrt{s}=4.63852$, $|\vec{P}|=2.22311$, $|\vec{P}'|=0.147625$
 $\theta_{in}=1.57079$, $\phi_l=3.82882$, $\phi_p=0.687223$
 $E_V=1.8$, $\phi_\gamma=0.589049$
 $Q^2=0.00488027$, $x_B=0.00156019$, $t=-2.40622$, $W^2=4.00295$, $y=0.151661$
 $\theta(l', \gamma)=176.268$ deg

four-momenta $[E, p_x, p_y, p_z]$ GeV:
 l $E= 2.2256$ $p_x= -1.7185$ $p_y= -1.4103$ $p_z= -0.0000$
 P $E= 2.4129$ $p_x= 1.7185$ $p_y= 1.4103$ $p_z= 0.0000$
 l' $E= 1.8890$ $p_x= -1.4966$ $p_y= -1.1477$ $p_z= 0.0000$
 P' $E= 0.9495$ $p_x= 0.0000$ $p_y= 0.1476$ $p_z= 0.0000$
 q_γ $E= 1.8000$ $p_x= 1.4966$ $p_y= 1.0000$ $p_z= 0.0000$

Transverse plane



Leading initial-ensemble/final-state components (N=8, retained=100.0%)

