



high_Egamma Ty proton: max $C_{\mu\gamma}$ regions



Ty proton characteristic max $C_{\mu\gamma}$ configuration

3D momenta




c_mu_gamma_high_egamma_ty_proton_region_0 C_{\mu\gamma}=0.0489763
 region: high_Egamma_Ty_proton_region_0 final-pair delta_xy=0.510997 rad
 outgoing-state purity: 0.582694
 kinematic point: high_s_high_theta_in_high_Egamma
 incoming state: $\frac{1}{2}\sum_{h_\mu}\rho(h_\mu, T_{y,p})$

$s=21.5158$, $\sqrt{s}=4.63852$, $|\vec{P}|=2.22311$, $|\vec{P}'|=2.19415$
 $\theta_{in}=1.57079$, $\phi_l=4.71239$, $\phi_p=1.5708$
 $E_\gamma=1.8$, $\phi_\gamma=4.61421$
 $Q^2=0.199845$, $x_B=0.0120018$, $t=-0.000128163$, $W^2=17.3312$, $y=0.807338$
 $\theta(l', \gamma)=29.278$ deg

four-momenta $[E, p_x, p_y, p_z]$ GeV:


l $E= 2.2256$ $p_x= -0.0000$ $p_y= -2.2231$ $p_z= -0.0000$
 P $E= 2.4129$ $p_x= 0.0000$ $p_y= 2.2231$ $p_z= 0.0000$
 l' $E= 0.4523$ $p_x= 0.1764$ $p_y= -0.4028$ $p_z= 0.0000$
 P' $E= 2.3862$ $p_x= 0.0000$ $p_y= 2.1942$ $p_z= 0.0000$
 q_γ $E= 1.8000$ $p_x= -0.1764$ $p_y= -1.7913$ $p_z= 0.0000$

Transverse plane




$h_e=+1, h_p=-1, h_\gamma=+1$
 electron $h=+1$, proton Ty
 $h_e=-1, h_p=+1, h_\gamma=-1$
 electron $h=-1$, proton Ty
 $h_e=-1, h_p=-1, h_\gamma=+1$
 electron $h=-1$, proton Ty
 $h_e=+1, h_p=+1, h_\gamma=-1$
 electron $h=+1$, proton Ty
 $h_e=+1, h_p=-1, h_\gamma=-1$
 electron $h=+1$, proton Ty
 $h_e=-1, h_p=+1, h_\gamma=+1$
 electron $h=-1$, proton Ty
 $h_e=+1, h_p=-1, h_\gamma=-1$
 electron $h=-1$, proton Ty
 $h_e=-1, h_p=+1, h_\gamma=+1$
 electron $h=+1$, proton Ty

Leading initial-ensemble/final-state components (N=8, retained=83.1%)



Ty proton characteristic max $C_{\mu\gamma}$ configuration

3D momenta




c_mu_gamma_high_egamma_ty_proton_region_1 $C_{\mu\gamma}=0.0340969$
 region: high_Egamma_Ty_proton_region_1 final-pair delta_xy=0.510997 rad
 outgoing-state purity: 0.526907
 kinematic point: high_s_high_theta_in_high_Egamma
 incoming state: $\frac{1}{2}\sum_{h_\mu}\rho(h_\mu, T_{y,p})$


$s=21.5158$, $\sqrt{s}=4.63852$, $|\vec{P}|=2.22311$, $|\vec{P}'|=2.19415$
 $\theta_{in}=1.57079$, $\phi_i=0.0981748$, $\phi_p=3.23977$
 $E_\gamma=1.8$, $\phi_\gamma=4.81056$
 $Q^2=2.94709$, $x_B=0.151924$, $t=-10.712$, $W^2=17.3312$, $y=0.940539$
 $\theta(l', \gamma)=29.278$ deg

four-momenta $[E, p_x, p_y, p_z]$ GeV:
 l $E= 2.2256$ $p_x= 2.2124$ $p_y= 0.2179$ $p_z= -0.0000$
 P $E= 2.4129$ $p_x= -2.2124$ $p_y= -0.2179$ $p_z= 0.0000$
 l' $E= 0.4523$ $p_x= -0.1764$ $p_y= -0.4028$ $p_z= 0.0000$
 P' $E= 2.3862$ $p_x= 0.0000$ $p_y= 2.1942$ $p_z= 0.0000$
 q_γ $E= 1.8000$ $p_x= 0.1764$ $p_y= -1.7913$ $p_z= 0.0000$

Transverse plane




Leading initial-ensemble/final-state components (N=5, retained=94.7%)



Ty proton characteristic max $C_{\mu\gamma}$ configuration

3D momenta




c_mu_gamma_high_egamma_ty_proton_region_2 $C_{\mu\gamma}=0.0340969$
 region: high_Egamma_Ty_proton_region_2 final-pair delta_xy=0.510997 rad
 outgoing-state purity: 0.526907
 kinematic point: high_s_high_theta_in_high_Egamma
 incoming state: $\frac{1}{2}\sum_{h_\mu}\rho(h_\mu, T_{y,p})$

$s=21.5158$, $\sqrt{s}=4.63852$, $|\vec{P}|=2.22311$, $|\vec{P}'|=2.19415$
 $\theta_{in}=1.57079$, $\phi_l=3.04342$, $\phi_P=6.18501$
 $E_\gamma=1.8$, $\phi_\gamma=4.61421$
 $Q^2=2.94709$, $x_B=0.151924$, $t=-10.712$, $W^2=17.3312$, $y=0.940539$
 $\theta(l', \gamma)=29.278$ deg

four-momenta $[E, p_x, p_y, p_z]$ GeV:

l	$E= 2.2256$	$p_x= -2.2124$	$p_y= 0.2179$	$p_z= -0.0000$
P	$E= 2.4129$	$p_x= 2.2124$	$p_y= -0.2179$	$p_z= 0.0000$
l'	$E= 0.4523$	$p_x= 0.1764$	$p_y= -0.4028$	$p_z= 0.0000$
P'	$E= 2.3862$	$p_x= 0.0000$	$p_y= 2.1942$	$p_z= 0.0000$
q_γ	$E= 1.8000$	$p_x= -0.1764$	$p_y= -1.7913$	$p_z= 0.0000$

Transverse plane



Leading initial-ensemble/final-state components (N=5, retained=94.7%)

