



high_Egamma Ty proton: max $C_{\mu p}$ regions



Ty proton characteristic max $C_{\mu p}$ configuration

3D momenta




c_mu_p_high_egamma_ty_proton_region_0 C_{\mu p}=0.139948
 region: high_Egamma_Ty_proton_region_0 final-pair delta_xy=2.72877 rad
 outgoing-state purity: 0.582694
 kinematic point: high_s_high_theta_in_high_Egamma
 incoming state: $\frac{1}{2}\sum_{h_\mu}\rho(h_\mu, T_{y,p})$

$s=21.5158$, $\sqrt{s}=4.63852$, $|\vec{P}|=2.22311$, $|\vec{P}'|=2.19415$
 $\theta_{in}=1.57079$, $\phi_l=4.71239$, $\phi_p=1.5708$
 $E_Y=1.8$, $\phi_Y=4.81056$
 $Q^2=0.199845$, $x_B=0.0120018$, $t=-0.000128163$, $W^2=17.3312$, $y=0.807338$
 $\theta(l', \gamma)=29.278$ deg


four-momenta $[E, p_x, p_y, p_z]$ GeV:
 l $E= 2.2256$ $p_x= -0.0000$ $p_y= -2.2231$ $p_z= -0.0000$
 P $E= 2.4129$ $p_x= 0.0000$ $p_y= 2.2231$ $p_z= 0.0000$
 l' $E= 0.4523$ $p_x= -0.1764$ $p_y= -0.4028$ $p_z= 0.0000$
 P' $E= 2.3862$ $p_x= 0.0000$ $p_y= 2.1942$ $p_z= 0.0000$
 q_Y $E= 1.8000$ $p_x= 0.1764$ $p_y= -1.7913$ $p_z= 0.0000$

Transverse plane




$h_e=+1, h_p=-1, h_\gamma=+1$
 electron $h=+1$, proton Ty
 $h_e=-1, h_p=+1, h_\gamma=-1$
 electron $h=-1$, proton Ty
 $h_e=-1, h_p=-1, h_\gamma=+1$
 electron $h=-1$, proton Ty
 $h_e=+1, h_p=+1, h_\gamma=-1$
 electron $h=+1$, proton Ty
 $h_e=+1, h_p=-1, h_\gamma=-1$
 electron $h=+1$, proton Ty
 $h_e=-1, h_p=+1, h_\gamma=+1$
 electron $h=-1$, proton Ty
 $h_e=+1, h_p=-1, h_\gamma=-1$
 electron $h=-1$, proton Ty
 $h_e=-1, h_p=+1, h_\gamma=+1$
 electron $h=+1$, proton Ty

Leading initial-ensemble/final-state components (N=8, retained=83.1%)



Ty proton characteristic max $C_{\mu p}$ configuration

3D momenta




c_mu_p_high_egamma_ty_proton_region_1 C_{\mu p}=0.00972081
 region: high_Egamma_Ty_proton_region_1 final-pair delta_xy=1.82802 rad
 outgoing-state purity: 0.501149
 kinematic point: high_s_high_theta_in_high_Egamma
 incoming state: $\frac{1}{2}\sum_{h_\mu}\rho(h_\mu, T_{y,p})$


$s=21.5158$, $\sqrt{s}=4.63852$, $|\vec{P}|=2.22311$, $|\vec{P}'|=0.29479$
 $\theta_{in}=1.57079$, $\phi_l=3.23977$, $\phi_P=0.0981748$
 $E_\gamma=1.8$, $\phi_\gamma=3.04342$
 $Q^2=15.9569$, $x_B=0.822839$, $t=-2.85671$, $W^2=4.31546$, $y=0.940253$
 $\theta(l', \gamma)=170.887$ deg

four-momenta $[E, p_x, p_y, p_z]$ GeV:
 l $E= 2.2256$ $p_x= -2.2124$ $p_y= -0.2179$ $p_z= -0.0000$
 P $E= 2.4129$ $p_x= 2.2124$ $p_y= 0.2179$ $p_z= 0.0000$
 l' $E= 1.8553$ $p_x= 1.7913$ $p_y= -0.4712$ $p_z= 0.0000$
 P' $E= 0.9832$ $p_x= 0.0000$ $p_y= 0.2948$ $p_z= 0.0000$
 q_γ $E= 1.8000$ $p_x= -1.7913$ $p_y= 0.1764$ $p_z= 0.0000$

Transverse plane




Leading initial-ensemble/final-state components (N=4, retained=93.2%)



Ty proton characteristic max $C_{\mu p}$ configuration

3D momenta




c_mu_p_high_egamma_ty_proton_region_2 C_{\mu p}=0.00972081
 region: high_Egamma_Ty_proton_region_2 final-pair delta_xy=1.82802 rad
 outgoing-state purity: 0.501149
 kinematic point: high_s_high_theta_in_high_Egamma
 incoming state: $\frac{1}{2}\sum_{h_\mu}\rho(h_\mu, T_{y,p})$


$s=21.5158$, $\sqrt{s}=4.63852$, $|\vec{P}|=2.22311$, $|\vec{P}'|=0.29479$
 $\theta_{in}=1.57079$, $\phi_\ell=6.18501$, $\phi_P=3.04342$
 $E_\gamma=1.8$, $\phi_\gamma=0.0981748$
 $Q^2=15.9569$, $x_B=0.822839$, $t=-2.85671$, $W^2=4.31546$, $y=0.940253$
 $\theta(\ell', \gamma)=170.887$ deg

four-momenta $[E, p_x, p_y, p_z]$ GeV:
 ℓ $E= 2.2256$ $p_x= 2.2124$ $p_y= -0.2179$ $p_z= -0.0000$
 P $E= 2.4129$ $p_x= -2.2124$ $p_y= 0.2179$ $p_z= 0.0000$
 ℓ' $E= 1.8553$ $p_x= -1.7913$ $p_y= -0.4712$ $p_z= 0.0000$
 P' $E= 0.9832$ $p_x= 0.0000$ $p_y= 0.2948$ $p_z= 0.0000$
 q_γ $E= 1.8000$ $p_x= 1.7913$ $p_y= 0.1764$ $p_z= 0.0000$

Transverse plane




Leading initial-ensemble/final-state components (N=4, retained=93.2%)



Ty proton characteristic max $C_{\mu p}$ configuration

3D momenta




c_mu_p_high_egamma_ty_proton_region_3 $C_{\mu p}=0.0093982$
 region: high_Egamma_Ty_proton_region_3 final-pair delta_xy=2.6765 rad
 outgoing-state purity: 0.50004
 kinematic point: high_s_high_theta_in_high_Egamma
 incoming state: $\frac{1}{2}\sum_{h_\mu}\rho(h_\mu, T_{\gamma, p})$

$s=21.5158$, $\sqrt{s}=4.63852$, $|\vec{P}|=2.22311$, $|\vec{P}'|=0.103468$
 $\theta_{in}=1.57079$, $\phi_l=0.883573$, $\phi_p=4.02517$
 $E_\gamma=1.8$, $\phi_\gamma=1.07992$
 $Q^2=16.6171$, $x_B=0.844112$, $t=-3.14998$, $W^2=3.94862$, $y=0.954472$
 $\theta(l', \gamma)=178.523$ deg

four-momenta $[E, p_x, p_y, p_z]$ GeV:

l $E= 2.2256$ $p_x= 1.4103$ $p_y= 1.7185$ $p_z= -0.0000$
 P $E= 2.4129$ $p_x= -1.4103$ $p_y= -1.7185$ $p_z= 0.0000$
 l' $E= 1.8948$ $p_x= -0.8485$ $p_y= -1.6909$ $p_z= 0.0000$
 P' $E= 0.9437$ $p_x= 0.0000$ $p_y= 0.1035$ $p_z= 0.0000$
 q_γ $E= 1.8000$ $p_x= 0.8485$ $p_y= 1.5875$ $p_z= 0.0000$

Transverse plane



Leading initial-ensemble/final-state components (N=4, retained=93.2%)

