


Tx electron: C_{ep} two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 0.5 \text{ GeV}$, $\max C_{ep} = 0.000$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1 \text{ GeV}$, $\max C_{ep} = 0.000$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1.8 \text{ GeV}$, $\max C_{ep} = 0.000$




Tx electron: $C_{e\gamma}$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 0.5 \text{ GeV}$, $\max C_{e\gamma} = 0.548$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1 \text{ GeV}$, $\max C_{e\gamma} = 0.553$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1.8 \text{ GeV}$, $\max C_{e\gamma} = 0.931$




Tx electron: $C_{p\gamma}$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 0.5 \text{ GeV}$, $\max C_{p\gamma} = 0.000$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1 \text{ GeV}$, $\max C_{p\gamma} = 0.000$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1.8 \text{ GeV}$, $\max C_{p\gamma} = 0.000$



Tx electron: $C_{e|p\gamma}$ two-angle scans at characteristic kinematics

Tx electron: $C_{p|e\gamma}$ two-angle scans at characteristic kinematics

Tx electron: $C_{\gamma|ep}$ two-angle scans at characteristic kinematics

Tx electron: M_e two-angle scans at characteristic kinematics

Tx electron: M_p two-angle scans at characteristic kinematics

Tx electron: M_γ two-angle scans at characteristic kinematics

Tx electron: F_3 two-angle scans at characteristic kinematics