


Tx proton: C_{ep} two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 0.5 \text{ GeV}$, max $C_{ep} = 0.000$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1 \text{ GeV}$, max $C_{ep} = 0.000$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1.8 \text{ GeV}$, max $C_{ep} = 0.000$




Tx proton: $C_{e\gamma}$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 0.5 \text{ GeV}$, max $C_{e\gamma} = 0.000$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1 \text{ GeV}$, max $C_{e\gamma} = 0.000$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1.8 \text{ GeV}$, max $C_{e\gamma} = 0.000$




Tx proton: $C_{p\gamma}$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 0.5 \text{ GeV}$, max $C_{p\gamma} = 1.000$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1 \text{ GeV}$, max $C_{p\gamma} = 0.999$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1.8 \text{ GeV}$, max $C_{p\gamma} = 0.987$



Tx proton: $C_{e|p\gamma}$ two-angle scans at characteristic kinematics

Tx proton: $C_{p|e\gamma}$ two-angle scans at characteristic kinematics

Tx proton: $C_{\gamma|ep}$ two-angle scans at characteristic kinematics

Tx proton: M_e two-angle scans at characteristic kinematics

Tx proton: M_p two-angle scans at characteristic kinematics

Tx proton: M_γ two-angle scans at characteristic kinematics

Tx proton: F_3 two-angle scans at characteristic kinematics