



low\_Egamma Tx\_electron: max  $C_{ey}$  regions



# Tx\_electron characteristicmax $C_{e\gamma}$ configuration

3D momenta




c\_e\_gamma\_low\_egamma\_tx\_electron\_region\_0  $C_{e\gamma}=0.547539$   
 region: low\_Egamma\_Tx\_electron\_region\_0 final-pair delta\_xy=0.168697 rad  
 kinematic point: high\_s\_high\_theta\_in\_low\_Egamma  
 incoming state:  $\frac{1}{2}\sum_{h_p}\rho(T_{X,e}, h_p)$

$s=21.5158$ ,  $\sqrt{s}=4.63852$ ,  $|\vec{P}|=2.22442$ ,  $|\vec{P}'|=2.22155$   
 $\theta_{in}=1.57079$ ,  $\phi_l=4.71239$ ,  $\phi_p=1.5708$   
 $E_\gamma=0.5$ ,  $\phi_\gamma=4.58149$   
 $Q^2=0.00548781$ ,  $x_B=0.00118797$ ,  $t=-1.2435e-06$ ,  $W^2=5.49384$ ,  $y=0.223856$   
 $\theta(l', \gamma)=9.66564$  deg

four-momenta  $[E, p_x, p_y, p_z]$  GeV:  
 $l$   $E= 2.2244$   $p_x= -0.0000$   $p_y= -2.2244$   $p_z= -0.0000$   
 $P$   $E= 2.4141$   $p_x= 0.0000$   $p_y= 2.2244$   $p_z= 0.0000$   
 $l'$   $E= 1.7271$   $p_x= 0.0653$   $p_y= -1.7258$   $p_z= 0.0000$   
 $P'$   $E= 2.4115$   $p_x= 0.0000$   $p_y= 2.2215$   $p_z= 0.0000$   
 $q_\gamma$   $E= 0.5000$   $p_x= -0.0653$   $p_y= -0.4957$   $p_z= 0.0000$

Transverse plane



$h_e=-1, h_p=+1, h_\gamma=+1$   
 electron Tx, proton h=+1  
 $h_e=+1, h_p=-1, h_\gamma=-1$   
 electron Tx, proton h=-1  
 $h_e=-1, h_p=-1, h_\gamma=+1$   
 electron Tx, proton h=-1  
 $h_e=+1, h_p=+1, h_\gamma=-1$   
 electron Tx, proton h=+1  
 $h_e=+1, h_p=-1, h_\gamma=+1$   
 electron Tx, proton h=+1  
 $h_e=-1, h_p=+1, h_\gamma=-1$   
 electron Tx, proton h=-1  
 $h_e=+1, h_p=+1, h_\gamma=+1$   
 electron Tx, proton h=+1  
 $h_e=-1, h_p=-1, h_\gamma=-1$   
 electron Tx, proton h=-1

Leading initial-ensemble/final-state components (N=8, retained=87.7%)

