



low_Egamma TxTx: max C_{py} regions



TxTx characteristic max $C_{p\gamma}$ configuration

3D momenta




c_p_gamma_low_egamma_txtx_region_0 C_{p\gamma}=0.999931
 region: low_Egamma_TxTx_region_0 final-pair delta_xy=0 rad
 kinematic point: high_s_high_theta_in_low_Egamma
 incoming state: $|T_{x,e}T_{x,p}\rangle$

$s=21.5158$, $\sqrt{s}=4.63852$, $|\vec{P}|=2.22442$, $|\vec{P}'|=1.69835$
 $\theta_{in}=1.57079$, $\phi_l=4.71239$, $\phi_p=1.5708$
 $E_\gamma=0.5$, $\phi_\gamma=1.5708$
 $Q^2=8.60837e-12$, $x_B=3.55992e-11$, $t=-0.0521317$, $W^2=1.12166$, $y=0.0117181$
 $\theta(\ell', \gamma)=180$ deg

four-momenta $[E, p_x, p_y, p_z]$ GeV:
 ℓ $E= 2.2244$ $p_x= -0.0000$ $p_y= -2.2244$ $p_z= -0.0000$
 P $E= 2.4141$ $p_x= 0.0000$ $p_y= 2.2244$ $p_z= 0.0000$
 ℓ' $E= 2.1984$ $p_x= -0.0000$ $p_y= -2.1984$ $p_z= 0.0000$
 P' $E= 1.9402$ $p_x= 0.0000$ $p_y= 1.6984$ $p_z= 0.0000$
 q_γ $E= 0.5000$ $p_x= 0.0000$ $p_y= 0.5000$ $p_z= 0.0000$

Transverse plane




$h_e=-1, h_p=+1, h_\gamma=-1$
 electron Tx, proton Tx

$h_e=+1, h_p=-1, h_\gamma=+1$
 electron Tx, proton Tx

$h_e=+1, h_p=+1, h_\gamma=-1$
 electron Tx, proton Tx


$h_e=-1, h_p=-1, h_\gamma=+1$
 electron Tx, proton Tx

Leading initial-ensemble/final-state components (N=4, retained=100.0%)



TxTx characteristic max $C_{p\gamma}$ configuration

3D momenta




c_p_gamma_low_egamma_txtx_region_1 C_{p\gamma}=0.981124
 region: low_Egamma TxTx_region_1 final-pair delta_xy=1.37445 rad
 kinematic point: high_s_high_theta_in_low_Egamma
 incoming state: $|T_{x,e}T_{x,p}\rangle$

$s=21.5158$, $\sqrt{s}=4.63852$, $|\vec{P}|=2.22442$, $|\vec{P}'|=1.88003$
 $\theta_{in}=1.57079$, $\phi_l=4.71239$, $\phi_P=1.5708$
 $E_\gamma=0.5$, $\phi_\gamma=2.94524$
 $Q^2=0.266466$, $x_B=0.133185$, $t=-0.0205939$, $W^2=2.61411$, $y=0.0969534$
 $\theta(l', \gamma)=115.177$ deg

four-momenta $[E, p_x, p_y, p_z]$ GeV:
 l $E= 2.2244$ $p_x= -0.0000$ $p_y= -2.2244$ $p_z= -0.0000$
 P $E= 2.4141$ $p_x= 0.0000$ $p_y= 2.2244$ $p_z= 0.0000$
 l' $E= 2.0375$ $p_x= 0.4904$ $p_y= -1.9776$ $p_z= 0.0000$
 P' $E= 2.1010$ $p_x= 0.0000$ $p_y= 1.8800$ $p_z= 0.0000$
 q_γ $E= 0.5000$ $p_x= -0.4904$ $p_y= 0.0975$ $p_z= 0.0000$

Transverse plane




$h_e=+1, h_p=+1, h_\gamma=-1$
 electron Tx, proton Tx

$h_e=+1, h_p=-1, h_\gamma=+1$
 electron Tx, proton Tx

$h_e=-1, h_p=+1, h_\gamma=-1$
 electron Tx, proton Tx


$h_e=-1, h_p=-1, h_\gamma=+1$
 electron Tx, proton Tx

Leading initial-ensemble/final-state components (N=4, retained=99.3%)



TxTx characteristic max $C_{p\gamma}$ configuration

3D momenta




c_p_gamma_low_egamma_txtx_region_2 C_{p\gamma}=0.981124
 region: low_Egamma TxTx_region_2 final-pair delta_xy=1.37445 rad
 kinematic point: high_s_high_theta_in_low_Egamma
 incoming state: $|T_{x,e}T_{x,p}\rangle$

$s=21.5158$, $\sqrt{s}=4.63852$, $|\vec{P}|=2.22442$, $|\vec{P}'|=1.88003$
 $\theta_{in}=1.57079$, $\phi_l=4.71239$, $\phi_P=1.5708$
 $E_\gamma=0.5$, $\phi_\gamma=0.19635$
 $Q^2=0.266466$, $x_B=0.133185$, $t=-0.0205939$, $W^2=2.61411$, $y=0.0969534$
 $\theta(l', \gamma)=115.177$ deg

four-momenta $[E, p_x, p_y, p_z]$ GeV:
 l $E= 2.2244$ $p_x= -0.0000$ $p_y= -2.2244$ $p_z= -0.0000$
 P $E= 2.4141$ $p_x= 0.0000$ $p_y= 2.2244$ $p_z= 0.0000$
 l' $E= 2.0375$ $p_x= -0.4904$ $p_y= -1.9776$ $p_z= 0.0000$
 P' $E= 2.1010$ $p_x= 0.0000$ $p_y= 1.8800$ $p_z= 0.0000$
 q_γ $E= 0.5000$ $p_x= 0.4904$ $p_y= 0.0975$ $p_z= 0.0000$

Transverse plane



$h_e=-1, h_p=-1, h_\gamma=+1$
 electron Tx, proton Tx

$h_e=-1, h_p=+1, h_\gamma=-1$
 electron Tx, proton Tx

$h_e=+1, h_p=-1, h_\gamma=+1$
 electron Tx, proton Tx

$h_e=+1, h_p=+1, h_\gamma=-1$
 electron Tx, proton Tx

Leading initial-ensemble/final-state components (N=4, retained=99.3%)

