


coherent mixing angles: maximum C_{ep} configuration (gradient_local_minimum)

z--x plane




max_c_ep_mixing_angles_polarization_02_local_minimum_7
 $C_{\{ep\}}=1$
 region: polarization_cluster_02_local_minimum_7
 outgoing-state purity: 1
 kinematic point: gradient_local_minimum
 lepton mixing angle: $\alpha_e=0.59023842$ rad
 incoming lepton: $0.830808|+\rangle +0.556559|-\rangle$
 proton mixing angle: $\alpha_p=2.2386433$ rad
 incoming proton: $-0.619297|+\rangle +0.785157|-\rangle$


$s=13.9759$, $\sqrt{s}=3.73844$, $|\vec{P}|=1.74985$, $|\vec{P}'|=1.65798$
 $\theta_{p'}=3.14159$, $\theta_\gamma=3.12202$, $\phi_{p'}=4.39523$, $\phi_\gamma=0.000121515$
 $E_\gamma=0.0861769$, $\theta_{\ell'}=0.000966173$, $\phi_{\ell'}=3.13898$
 $Q^2=12.2079$, $x_B=0.996521$, $t=-11.6068$, $W^2=0.922461$, $y=0.936234$

four-momenta $[E, p_x, p_y, p_z]$ GeV:
 ℓ $E= 1.7530$ $p_x= 0.0000$ $p_y= 0.0000$ $p_z= -1.7498$
 P $E= 1.9854$ $p_x= 0.0000$ $p_y= 0.0000$ $p_z= 1.7498$
 ℓ' $E= 1.7473$ $p_x= -0.0017$ $p_y= 0.0000$ $p_z= 1.7441$
 P' $E= 1.9049$ $p_x= -0.0000$ $p_y= -0.0000$ $p_z= -1.6580$
 q_γ $E= 0.0862$ $p_x= 0.0017$ $p_y= 0.0000$ $p_z= -0.0862$

x--y plane



Leading normalized final-state amplitudes
 (N=4, retained=100.0%)



muon: polarization cluster P2, local minimum 7; pairwise projections of the 8D $1 - C_{ep} = (1 - C_{ep})_{\text{local}} + 0.05$ contour

