


coherent mixing angles: maximum $C_{p\gamma}$ configuration (gradient_local_minimum)

z--x plane




max_c_p_gamma_mixing_angles_polarization_02_local_minimum_8
 $C_{\{p\backslash\gamma\}}=1$
 region: polarization_cluster_02_local_minimum_8
 outgoing-state purity: 1
 kinematic point: gradient_local_minimum
 lepton mixing angle: $\alpha_e=1.5707933$ rad
 incoming lepton: $2.98005e-06|+\rangle +1|-\rangle$
 proton mixing angle: $\alpha_p=2.4088937$ rad
 incoming proton: $-0.743372|+\rangle +0.668878|-\rangle$

$s=14.288$, $\sqrt{s}=3.77995$, $|\vec{P}|=1.77192$, $|\vec{P}'|=0.453461$
 $\theta_{P'}=0$, $\theta_\gamma=0$, $\phi_{P'}=1.26489$, $\phi_\gamma=1.99329$
 $E_\gamma=1.14056$, $\theta_{\ell'}=3.14159$, $\phi_{\ell'}=0$
 $Q^2=0.000124587$, $x_B=9.28138e-05$, $t=-0.810925$, $W^2=2.22206$, $y=0.100197$


four-momenta $[E, p_x, p_y, p_z]$ GeV:

ℓ	$E=$	1.7751	$p_x=$	0.0000	$p_y=$	0.0000	$p_z=$	-1.7719
P	$E=$	2.0049	$p_x=$	0.0000	$p_y=$	0.0000	$p_z=$	1.7719
ℓ'	$E=$	1.5975	$p_x=$	0.0000	$p_y=$	-0.0000	$p_z=$	-1.5940
P'	$E=$	1.0419	$p_x=$	0.0000	$p_y=$	0.0000	$p_z=$	0.4535
q_γ	$E=$	1.1406	$p_x=$	-0.0000	$p_y=$	0.0000	$p_z=$	1.1406

x--y plane



Leading normalized final-state amplitudes
 (N=2, retained=100.0%)



muon: polarization cluster P2, local minimum 8; pairwise projections of the 8D $1 - C_{p\gamma} = (1 - C_{p\gamma})_{\text{local}} + 0.05$ contour

