


# L electron + Ty proton: $C_{ep}$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 0.25 \text{ GeV}$ ,  $\max C_{ep} = 0.102$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 1 \text{ GeV}$ ,  $\max C_{ep} = 0.026$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 1.8 \text{ GeV}$ ,  $\max C_{ep} = 0.014$




# L electron + Ty proton: $C_{e\gamma}$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 0.25 \text{ GeV}$ ,  $\max C_{e\gamma} = 0.098$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1 \text{ GeV}$ ,  $\max C_{e\gamma} = 0.015$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1.8 \text{ GeV}$ ,  $\max C_{e\gamma} = 0.005$




# L electron + Ty proton: $C_{p\gamma}$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 0.25 \text{ GeV}$ ,  $\max C_{p\gamma} = 1.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1 \text{ GeV}$ ,  $\max C_{p\gamma} = 0.999$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1.8 \text{ GeV}$ ,  $\max C_{p\gamma} = 0.991$




# L electron + Ty proton: $C_{e|p\gamma}$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 0.25 \text{ GeV}$ ,  $\max C_{e|p\gamma} = 0.137$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1 \text{ GeV}$ ,  $\max C_{e|p\gamma} = 0.029$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1.8 \text{ GeV}$ ,  $\max C_{e|p\gamma} = 0.015$




# L electron + Ty proton: $C_{p|e\gamma}$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 0.25 \text{ GeV}$ ,  $\max C_{p|e\gamma} = 1.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1 \text{ GeV}$ ,  $\max C_{p|e\gamma} = 0.999$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1.8 \text{ GeV}$ ,  $\max C_{p|e\gamma} = 0.991$




# L electron + Ty proton: $C_{\gamma|ep}$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 0.25 \text{ GeV}$ ,  $\max C_{\gamma|ep} = 1.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1 \text{ GeV}$ ,  $\max C_{\gamma|ep} = 0.999$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1.8 \text{ GeV}$ ,  $\max C_{\gamma|ep} = 0.991$




# L electron + Ty proton: $M_e$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 0.25 \text{ GeV}$ ,  $\max M_e = 0.008$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1 \text{ GeV}$ ,  $\max M_e = 0.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1.8 \text{ GeV}$ ,  $\max M_e = 0.000$




# L electron + Ty proton: $M_p$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 0.25 \text{ GeV}$ ,  $\max M_p = 0.008$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 1 \text{ GeV}$ ,  $\max M_p = 0.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 1.8 \text{ GeV}$ ,  $\max M_p = 0.000$




# L electron + Ty proton: $M_\gamma$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 0.25 \text{ GeV}$ ,  $\max M_\gamma = 0.008$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1 \text{ GeV}$ ,  $\max M_\gamma = 0.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1.8 \text{ GeV}$ ,  $\max M_\gamma = 0.000$




# L electron + Ty proton: $F_3$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 0.25 \text{ GeV}$ ,  $\max F_3 = 0.014$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 1 \text{ GeV}$ ,  $\max F_3 = 0.004$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 1.8 \text{ GeV}$ ,  $\max F_3 = 0.005$




# L electron + Ty proton: $P_{\text{GHZ}+}$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 0.25 \text{ GeV}$ ,  $\max P_{\text{GHZ}+} = 0.280$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1 \text{ GeV}$ ,  $\max P_{\text{GHZ}+} = 0.384$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1.8 \text{ GeV}$ ,  $\max P_{\text{GHZ}+} = 0.483$



# L electron + Ty proton: $P_W$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 0.25 \text{ GeV}$ ,  $\max P_W = 0.333$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 1 \text{ GeV}$ ,  $\max P_W = 0.334$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 1.8 \text{ GeV}$ ,  $\max P_W = 0.333$

