


# Tx electron + Ty proton: $C_{ep}$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 0.25 \text{ GeV}$ ,  $\max C_{ep} = 0.993$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1 \text{ GeV}$ ,  $\max C_{ep} = 0.845$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1.8 \text{ GeV}$ ,  $\max C_{ep} = 0.368$




# Tx electron + Ty proton: $C_{e\gamma}$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 0.25 \text{ GeV}$ ,  $\max C_{e\gamma} = 1.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1 \text{ GeV}$ ,  $\max C_{e\gamma} = 1.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1.8 \text{ GeV}$ ,  $\max C_{e\gamma} = 1.000$




# Tx electron + Ty proton: $C_{p\gamma}$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 0.25 \text{ GeV}$ ,  $\max C_{p\gamma} = 1.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1 \text{ GeV}$ ,  $\max C_{p\gamma} = 0.999$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1.8 \text{ GeV}$ ,  $\max C_{p\gamma} = 0.987$




# Tx electron + Ty proton: $C_{e|p\gamma}$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 0.25 \text{ GeV}$ ,  $\max C_{e|p\gamma} = 1.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1 \text{ GeV}$ ,  $\max C_{e|p\gamma} = 1.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1.8 \text{ GeV}$ ,  $\max C_{e|p\gamma} = 1.000$




# Tx electron + Ty proton: $C_{p|ey}$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 0.25 \text{ GeV}$ ,  $\max C_{p|ey} = 1.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 1 \text{ GeV}$ ,  $\max C_{p|ey} = 1.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 1.8 \text{ GeV}$ ,  $\max C_{p|ey} = 1.000$




# Tx electron + Ty proton: $C_{\gamma|ep}$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 0.25 \text{ GeV}$ ,  $\max C_{\gamma|ep} = 1.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1 \text{ GeV}$ ,  $\max C_{\gamma|ep} = 1.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1.8 \text{ GeV}$ ,  $\max C_{\gamma|ep} = 1.000$




# Tx electron + Ty proton: $M_e$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 0.25 \text{ GeV}$ ,  $\max M_e = 1.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 1 \text{ GeV}$ ,  $\max M_e = 1.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 1.8 \text{ GeV}$ ,  $\max M_e = 1.000$




# Tx electron + Ty proton: $M_p$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 0.25 \text{ GeV}$ ,  $\max M_p = 1.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 1 \text{ GeV}$ ,  $\max M_p = 1.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 1.8 \text{ GeV}$ ,  $\max M_p = 1.000$






# Tx electron + Ty proton: $M_Y$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 0.25 \text{ GeV}$ ,  $\max M_Y = 1.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 1 \text{ GeV}$ ,  $\max M_Y = 1.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 1.8 \text{ GeV}$ ,  $\max M_Y = 1.000$




# Tx electron + Ty proton: $F_3$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 0.25 \text{ GeV}$ ,  $\max F_3 = 1.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 1 \text{ GeV}$ ,  $\max F_3 = 1.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 1.8 \text{ GeV}$ ,  $\max F_3 = 1.000$




# Tx electron + Ty proton: $P_{\text{GHZ}+}$ + two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 0.25 \text{ GeV}$ ,  $\max P_{\text{GHZ}+} = 0.546$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1 \text{ GeV}$ ,  $\max P_{\text{GHZ}+} = 0.753$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1.8 \text{ GeV}$ ,  $\max P_{\text{GHZ}+} = 0.958$



# Tx electron + Ty proton: $P_W$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 0.25 \text{ GeV}$ ,  $\max P_W = 0.469$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 1 \text{ GeV}$ ,  $\max P_W = 0.502$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 1.8 \text{ GeV}$ ,  $\max P_W = 0.461$

