


Tx electron: C_{ep} two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 0.25 \text{ GeV}$, $\max C_{ep} = 0.000$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1 \text{ GeV}$, $\max C_{ep} = 0.000$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1.8 \text{ GeV}$, $\max C_{ep} = 0.000$




Tx electron: $C_{e\gamma}$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 0.25 \text{ GeV}$, $\max C_{e\gamma} = 1.000$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1 \text{ GeV}$, $\max C_{e\gamma} = 1.000$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1.8 \text{ GeV}$, $\max C_{e\gamma} = 1.000$




Tx electron: $C_{p\gamma}$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 0.25 \text{ GeV}$, $\max C_{p\gamma} = 0.000$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1 \text{ GeV}$, $\max C_{p\gamma} = 0.000$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1.8 \text{ GeV}$, $\max C_{p\gamma} = 0.000$




Tx electron: $C_{e|p\gamma}$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 0.25 \text{ GeV}$, $\max C_{e|p\gamma} = 0.000$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1 \text{ GeV}$, $\max C_{e|p\gamma} = 0.000$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1.8 \text{ GeV}$, $\max C_{e|p\gamma} = 0.000$




Tx electron: $C_{p|e\gamma}$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 0.25 \text{ GeV}$, $\max C_{p|e\gamma} = 0.000$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1 \text{ GeV}$, $\max C_{p|e\gamma} = 0.000$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1.8 \text{ GeV}$, $\max C_{p|e\gamma} = 0.000$




Tx electron: $C_{\gamma|ep}$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 0.25 \text{ GeV}$, $\max C_{\gamma|ep} = 0.000$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1 \text{ GeV}$, $\max C_{\gamma|ep} = 0.000$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1.8 \text{ GeV}$, $\max C_{\gamma|ep} = 0.000$




Tx electron: M_e two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_Y = 0.25 \text{ GeV}$, $\max M_e = -0.000$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_Y = 1 \text{ GeV}$, $\max M_e = -0.000$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_Y = 1.8 \text{ GeV}$, $\max M_e = -0.000$




Tx electron: M_p two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 0.25 \text{ GeV}$, $\max M_p = -0.000$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1 \text{ GeV}$, $\max M_p = -0.000$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1.8 \text{ GeV}$, $\max M_p = -0.000$




Tx electron: M_Y two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_Y = 0.25 \text{ GeV}$, $\max M_Y = -0.000$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_Y = 1 \text{ GeV}$, $\max M_Y = -0.000$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_Y = 1.8 \text{ GeV}$, $\max M_Y = -0.000$




Tx electron: F_3 two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 0.25 \text{ GeV}$, $\max F_3 = 0.000$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1 \text{ GeV}$, $\max F_3 = 0.000$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1.8 \text{ GeV}$, $\max F_3 = 0.000$




Tx electron: $P_{\text{GHZ}+}$ + two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_Y = 0.25 \text{ GeV}$, $\max P_{\text{GHZ}+} = 0.280$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_Y = 1 \text{ GeV}$, $\max P_{\text{GHZ}+} = 0.384$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_Y = 1.8 \text{ GeV}$, $\max P_{\text{GHZ}+} = 0.483$



Tx electron: P_W two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 0.25 \text{ GeV}$, $\max P_W = 0.332$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1 \text{ GeV}$, $\max P_W = 0.333$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1.8 \text{ GeV}$, $\max P_W = 0.333$

