




L proton: $C_{\ell_{Hp}}$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 0.25 \text{ GeV}$, $\max C_{\ell_{Hp}} = 0.017$




$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1 \text{ GeV}$, $\max C_{\ell_{Hp}} = 0.037$



L proton: $C_{\ell_H\gamma}$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 0.25 \text{ GeV}$, $\max C_{\ell_H\gamma} = 0.725$




$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1 \text{ GeV}$, $\max C_{\ell_H\gamma} = 0.576$



L proton: $C_{p\gamma}$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 0.25 \text{ GeV}$, $\max C_{p\gamma} = 0.776$




$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1 \text{ GeV}$, $\max C_{p\gamma} = 0.815$



L proton: $C_{\ell_H|p\gamma}$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 0.25 \text{ GeV}$, $\max C_{\ell_H|p\gamma} = 0.000$




$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1 \text{ GeV}$, $\max C_{\ell_H|p\gamma} = 0.000$



L proton: $C_{p|\ell_H\gamma}$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 0.25 \text{ GeV}$, $\max C_{p|\ell_H\gamma} = 0.000$




$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1 \text{ GeV}$, $\max C_{p|\ell_H\gamma} = 0.000$



L proton: $C_{\gamma|\ell_{HP}}$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 0.25 \text{ GeV}$, $\max C_{\gamma|\ell_{HP}} = 0.000$




$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1 \text{ GeV}$, $\max C_{\gamma|\ell_{HP}} = 0.000$



L proton: M_{ℓ_H} two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 0.25 \text{ GeV}$, $\max M_{\ell_H} = -0.000$




$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1 \text{ GeV}$, $\max M_{\ell_H} = -0.000$



L proton: M_p two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 0.25 \text{ GeV}$, $\max M_p = -0.000$




$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1 \text{ GeV}$, $\max M_p = -0.000$



L proton: M_Y two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_Y = 0.25 \text{ GeV}$, $\max M_Y = -0.000$




$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_Y = 1 \text{ GeV}$, $\max M_Y = -0.000$



L proton: F_3 two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 0.25 \text{ GeV}$, $\max F_3 = 0.000$




$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1 \text{ GeV}$, $\max F_3 = 0.000$



L proton: $P_{\text{GHZ}+}$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 0.25 \text{ GeV}$, $\max P_{\text{GHZ}+} = 0.280$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1 \text{ GeV}$, $\max P_{\text{GHZ}+} = 0.394$



L proton: P_W two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 0.25 \text{ GeV}$, $\max P_W = 0.215$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1 \text{ GeV}$, $\max P_W = 0.222$

