


Ty massless lepton: $C_{\ell_0 p}$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 0.25 \text{ GeV}$, $\max C_{\ell_0 p} = 0.000$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1 \text{ GeV}$, $\max C_{\ell_0 p} = 0.000$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1.8 \text{ GeV}$, $\max C_{\ell_0 p} = 0.000$




Ty massless lepton: $C_{\ell_0\gamma}$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 0.25 \text{ GeV}$, $\max C_{\ell_0\gamma} = 0.404$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1 \text{ GeV}$, $\max C_{\ell_0\gamma} = 0.672$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1.8 \text{ GeV}$, $\max C_{\ell_0\gamma} = 0.931$




Ty massless lepton: $C_{p\gamma}$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 0.25 \text{ GeV}$, $\max C_{p\gamma} = 0.000$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1 \text{ GeV}$, $\max C_{p\gamma} = 0.000$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1.8 \text{ GeV}$, $\max C_{p\gamma} = 0.000$




Ty massless lepton: $C_{\ell_0|p\gamma}$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 0.25 \text{ GeV}$, $\max C_{\ell_0|p\gamma} = 0.000$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1 \text{ GeV}$, $\max C_{\ell_0|p\gamma} = 0.000$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1.8 \text{ GeV}$, $\max C_{\ell_0|p\gamma} = 0.000$




Ty massless lepton: $C_{p|\ell_0\gamma}$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 0.25 \text{ GeV}$, $\max C_{p|\ell_0\gamma} = 0.000$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1 \text{ GeV}$, $\max C_{p|\ell_0\gamma} = 0.000$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1.8 \text{ GeV}$, $\max C_{p|\ell_0\gamma} = 0.000$




Ty massless lepton: $C_{\gamma|\ell_0 p}$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 0.25 \text{ GeV}$, $\max C_{\gamma|\ell_0 p} = 0.000$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1 \text{ GeV}$, $\max C_{\gamma|\ell_0 p} = 0.000$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1.8 \text{ GeV}$, $\max C_{\gamma|\ell_0 p} = 0.000$




Ty massless lepton: M_{ℓ_0} two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_Y = 0.25 \text{ GeV}$, $\max M_{\ell_0} = -0.000$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_Y = 1 \text{ GeV}$, $\max M_{\ell_0} = -0.000$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_Y = 1.8 \text{ GeV}$, $\max M_{\ell_0} = -0.000$




Ty massless lepton: M_p two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 0.25 \text{ GeV}$, $\max M_p = -0.000$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1 \text{ GeV}$, $\max M_p = -0.000$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1.8 \text{ GeV}$, $\max M_p = -0.000$




Ty massless lepton: M_Y two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_Y = 0.25 \text{ GeV}$, $\max M_Y = -0.000$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_Y = 1 \text{ GeV}$, $\max M_Y = -0.000$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_Y = 1.8 \text{ GeV}$, $\max M_Y = -0.000$




Ty massless lepton: F_3 two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 0.25 \text{ GeV}$, $\max F_3 = 0.000$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1 \text{ GeV}$, $\max F_3 = 0.000$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1.8 \text{ GeV}$, $\max F_3 = 0.000$




Ty massless lepton: $P_{\text{GHZ}+}$ + two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 0.25 \text{ GeV}$, $\max P_{\text{GHZ}+} = 0.280$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1 \text{ GeV}$, $\max P_{\text{GHZ}+} = 0.384$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1.8 \text{ GeV}$, $\max P_{\text{GHZ}+} = 0.483$



Ty massless lepton: P_W two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 0.25 \text{ GeV}$, $\max P_W = 0.229$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1 \text{ GeV}$, $\max P_W = 0.236$

$s = 21.5 \text{ GeV}^2$, $\theta_{\text{in}} = 1.57 \text{ rad}$
 $E_\gamma = 1.8 \text{ GeV}$, $\max P_W = 0.208$

