STAT 3011 Fall 2022 Exam 1 (A) Time Limit: 90 Minutes	Name (Print): Student ID:	
Instructions:		
• Do not begin or turn this page	e until you are instructed.	
• Enter all requested information the top of every page, in case	on on the top and bottom of this page, and put your initite pages become separated.	itials or
	including this cover page and the multiple choice answer missing. There are 13 multiple-choice problems and	,
• The exam is closed book. Do	not use your books, or any electronic devices on this e	exam.
	one sheet of paper (size A4 or 8.5" by 11") with form not share calculators or notes!	nulas oi
• Show all your work on each following rules apply:	problem for full credit except multiple choice problem	ıs. The

- Organize your work, in a reasonably neat and coherent way, in the space provided. Work scattered all over the page without a clear ordering will receive very little credit.
- Mysterious or unsupported answers will not receive full credit for short answer problems.
 A correct answer, unsupported by calculations, explanation, or algebraic work will not receive full credit; an incorrect answer supported by substantially correct calculations and explanations may still receive partial credit.
- If you need more space, use the back of the pages; clearly indicate when you have done this.

Honesty Statement and Pledge:

I have not given or received any aid or assistance to or from any other student in this course during the exam period. Everything I have written on this exam represents my own work and knowledge. I sign this knowing that infringements on the University's Academic Honest policy may result in failure or expulsion.

Signed By:	Date:	
§ • • —————————————————————————————————	•	

Problem 1. (40 points) Multiple Choice


Choose ONLY ONE answer for each question. Circle your answers to all questions in the answer sheet provided on page 15. (NO explanation is needed).

1. (3 points) A random sample of patients who received a hip transplant operation at Stanford University Hospital in 2022 will be followed for 10 years after their operation to determine the success (or failure) of the transplant.

Which of the following is correct?

- (A) Sample of this study is those who had a successful transplant operation in 2022.
- (B) This study is a randomized experiment.
- (C) This study uses a volunteer sample.
- (D) None of the above are correct.
- 2. (3 points) Which of the following graphs is *not* useful for judging the shape of distribution for a quantitative variable?
 - (A) Bar graph
 - (B) Box plot
 - (C) Histogram
 - (D) Stem and leaf plot
- 3. (3 points) Which of the following is true?
 - (A) Sample standard deviation can never be zero.
 - (B) Mean and median can never be the same.
 - (C) The mean is always one of the data points.
 - (D) All of the above are false.
- 4. (3 points) Let A and B be independent events. It is known that P(A) = 0.50 and $P(A \cap B) = 0.15$. Find the probability $P(A \cup B)$.
 - (A) 0.45
 - (B) 0.65
 - (C) 0.75
 - (D) 0.85
- 5. (3 points) Let C and D be events such that P(C) = 0.35, P(D) = 0.25, and $P(C \mid D) = 0.20$. What is the probability that EXACTLY ONE of C or D occurs (but NOT BOTH)?
 - (A) 0.40
 - (B) 0.50
 - (C) 0.55
 - (D) 0.60

6. (3 points) We have a random variable X with mean 0.5 and standard deviation 0.224. Its density curve is as follows. We know that the density at 0.9 is 0.54 and the probability that X is smaller than 0.1 equals to 0.028. What is the probability that X is smaller than or equal to 0.9?



- (A) 0.486
- (B) 0.54
- (C) pnorm(0.9, mean = 0.5, sd = 0.224)
- (D) 0.972
- 7. (3 points) The probability distribution of a random variable X is provided in the following table. Select the claim that is incorrect. (Hint: refer to the R command results.)

x	0	1	2	3
P(X=x)	0.343	0.441	0.189	0.027

```
> dbinom(x = 0, size = 3, prob = 0.3)
[1] 0.343
> dbinom(x = 1, size = 3, prob = 0.3)
[1] 0.441
> dbinom(x = 2, size = 3, prob = 0.3)
[1] 0.189
> dbinom(x = 3, size = 3, prob = 0.3)
[1] 0.027
```

- (A) $X \sim Bin(3, 0.3)$
- (B) The mean of X equals to 0.9.
- (C) This is a valid probability distribution.
- (D) The standard deviation of X equals to $\sqrt{0.07}$.

- 8. (3 points) Suppose we have a random variable $X \sim N(3,2)$, based on the 68-95-99.7 rule, what is the probability that X is greater than or equal to -3 and smaller than 7?
 - (A) 0.815
 - (B) 0.95
 - (C) 0.9735
 - (D) 0.975
- 9. (3 points) A college senior scored 95 on the psychology exam and scored 85 on the history exam. The mean score for the psychology exam is 89 with a standard deviation 3. The mean score for the history exam is 80 with a standard deviation 2.5. Suppose both distributions are normal. Relative to others, which exam did this student do better on?
 - (A) Psychology
 - (B) Equally well
 - (C) History
 - (D) Cannot decide based on the current information
- 10. (3 points) Suppose the height of female undergraduate students follows a normal distribution N(68, 2). What is the correct R command that gives us the probability that a randomly selected female undergraduate student is shorter than or equal to 69.5 inches?
 - (A) 1 pnorm(0.75)
 - (B) pnorm(0.75)
 - (C) pnorm(0.75, mean = 68, sd = 2)
 - (D) pnorm(69.5, mean = 68, sd = 2) + dnorm(69.5, mean = 68, sd = 2)
- 11. (3 points) Suppose the probability is p=0.2 that a person who purchases an instant lottery ticket wins money, and this probability holds for every ticket purchased. Consider different random samples of n=30 purchased tickets. Let \hat{p} be sample proportion of winning tickets in a sample of 30 tickets.

Which of the following is true?

- (A) Standard deviation of sampling distribution of sample proportion is $\sqrt{(0.2)(0.8)(30)} = 2.19$.
- (B) According to the 68-95-99.7 rule, approximately 95% of sample proportion of tickets that will be money winners will be between 12.7% and 27%.
- (C) The sampling distribution of sample proportion is *not* approximately normal.
- (D) All of the above are true.

- 12. (3 points) The Central Limit Theorem states:
 - (A) the shape of the sampling distribution of sample mean is always normal.
 - (B) For a large n, the shape of the sampling distribution of sample mean is approximately normal.
 - (C) the mean of sampling distribution of sample mean is equal to the population mean.
 - (D) the standard deviation of sampling distribution of sample mean is always less than the standard deviation of the population for n > 1.
- 13. (4 points) Did you circle multiple choice answers on page 15?
 - (A) Yes, I did.
 - (B) I will now.
 - (C) I will now.
 - (D) I will now.

Problem 2. (13 points) Be sure to show all work for full credit.


1. (4 points) In elections, television networks often declare the winner well before all the votes have been counted. They do this using exit polling, surveying voters after they leave the voting booth. To predict how (i) <u>9.5 million voters</u> in California voted in 2010, a TV exit poll used surveyed (ii) <u>3889 voters</u> and found (iii) <u>53.1%</u> of them voted for Jerry Brown (Democratic candidate in 2010 California gubernatorial race). After all the votes were counted, Jerry Brown won (iv) <u>53.8%</u> of votes.

Based on the information above, match each of the definitions a) - d) to one of (i)-(iv). If it is unknown, state "unknown".

- a) Population of interest
- b) Sample
- c) Parameter
- d) Statistic

Be sure to show all work for full credit.

Political scientists have noticed that presidential approval ratings tend to fluctuate with the seasons. To examine this claim, we use data on presidential quarterly approval ratings from 1945 to 1976. The data are displayed in the boxplots below.



Answer the following questions regarding the presidential approval rating data.

2. (3 points) During which quarter does the presidential approval rating seem to be highest? Clearly state your reasoning; state explicitly the statistic you are basing your answer on.

3. (3 points) Give an estimate for the IQR for the first quarter approval ratings.

4.	(3 points) Below i	is R comma	ands used to co	nstruct the side-by-	-side box-plot.	
	> dat <- read.c	sv(file.c	hoose())			
	> names(dat)					
	[1] "ratings" "	quarters"				
	> boxplot((a)	,	xlab="Quarter"	, ylab="Approval	Rating")
	Fill in the blank	(a)	that makes a s	side-by-side boxplot	j.	

Problem 3. (15 points) Be sure to show all work for full credit.

Below is a table that summarizes 302 students' college affiliations (College of Biology Science (CBS) or College of Liberal Arts (CLA)) and their favorite seasons.

	Favorite season				Total
	Spring Summer Fall Winter				
CBS	27	35	52	19	133
CLA	6	59	87	17	169
Total	33	94	139	36	302

Let:

- A be the event that a randomly selected student is in CLA
- B be the event a randomly selected student's favorite season is fall
- C be the event a randomly selected student's favorite season is summer.
- 1. (4 points) Find the probability that a randomly selected student is in CLA and this student likes fall the most.

Use A, B, and/or C to define this event and calculate the probability. Round your answer to the nearest 3 decimal places.

- 2. (4 points) Given that a randomly selected student is in CLA, what is the probability this student's favorite season is Fall?
 - Use A, B, and/or C to define this event and calculate the probability. Round your answer to the nearest 3 decimal places.

Copy of the table from the previous page

	Favorite season				Total
	Spring Summer Fall Winter				
CBS	27	35	52	19	133
CLA	6	59	87	17	169
Total	33	94	139	36	302

Let:

- A be the event that a randomly selected student is in CLA
- B be the event a randomly selected student's favorite season is fall
- C be the event a randomly selected student's favorite season is summer.
- 3. (3 points) Which pairs of events (A, B), (A, C), or (B, C), if any, are disjoint?

4. (4 points) Are A and B independent? Show your work to support your answer.

Problem 4. (20 points) Be sure to show all work for full credit.

Suppose the weights of eighteen-year-olds follow a normal distribution with mean 127 and standard deviation 12 (in pound). We randomly sample 4 eighteen-year-olds and ask if their weight is larger than 127 pounds.

You may find the following R command results helpful.

$$pnorm(0.5) = 0.6915$$
; $qnorm(0.242) = -0.7$; $qnorm(1 - 0.242) = 0.7$

- 1. (2 points) Let W represent the weight of a randomly selected eighteen-year-old. What is the distribution of W? Use statistical notation.
- 2. (4 points) Tom's weight is 133 pounds. What is his z-score? Interpret this z-score.

3. (4 points) What is the probability that a randomly selected teen is heavier than Tom? Round your answer to three decimal places.

Copy of Problem 4 descriptions from the previous page

Suppose the weights of eighteen-year-olds follow a normal distribution with mean 127 and standard deviation 12 (in pound). We randomly sample 4 eighteen-year-olds and ask if their weight is larger than 127 pounds.

You may find the following R command results helpful. pnorm(0.5) = 0.6915; qnorm(0.242) = -0.7; qnorm(1 - 0.242) = 0.7

4. (4 points) In order to be lighter than at least 24.2% of eighteen-year-olds, what is the maximum weight that Tom should have? Does he need to control his weight? Round your answer to one decimal place.

- 5. (2 points) We randomly sample 4 eighteen-year-olds and ask if their weight is larger than 127 pounds. Let V represent the number of teens who say they are heavier than 127 pounds. What is the distribution of V? Write down the statistical notation.
- 6. (4 points) What is the probability that at least 2 teens in our sample say they are heavier than 127 pounds? Show your work including formulas with numbers plugged in.

Problem 5. (12 points) Be sure to show all work for full credit.

Sheila's doctor is concerned that she may suffer from gestational diabetes (high blood glucose levels during pregnancy). There is variation both in the actual glucose level and in the blood test that measures the level. A patient is classified as having gestational diabetes if the glucose level is above 135 milligrams per deciliter (mg/dl) one hour after a sugary drink is digested. Sheila's measured glucose level one hour after ingesting the sugary drink varies according to the normal distribution with $\mu = 120 \text{mg/dl}$ and $\sigma = 10 \text{mg/dl}$.

You may find the following R code helpful.

```
pnorm(1.5) = 0.9332, pnorm(2.0) = 0.97725, pnorm(3.0) = 0.99865, qnorm(0.01) = -2.33, qnorm(0.025) = -1.96, qnorm(0.05) = -1.64.
```

1. (4 points) If measurements are made on randomly selected four separate days and sample mean is computed, describe the shape, center, and spread of the sampling distribution of sample mean. Write down the statistical notation for this sampling distribution.

2. (4 points) If the sample mean of Sheila's four readings is compared with the criterion 135mg/dl, what is the probability that Sheila is diagnosed as having gestational diabetes?

Copy of problem 5 descriptions from the previous page.

Sheila's doctor is concerned that she may suffer from gestational diabetes (high blood glucose levels during pregnancy). There is variation both in the actual glucose level and in the blood test that measures the level. A patient is classified as having gestational diabetes if the glucose level is above 135 milligrams per deciliter (mg/dl) one hour after a sugary drink is digested. Sheila's measured glucose level one hour after ingesting the sugary drink varies according to the normal distribution with $\mu = 120 \text{mg/dl}$ and $\sigma = 10 \text{mg/dl}$.

```
You may find the following R code helpful.

pnorm(1.5) = 0.9332, pnorm(2.0) = 0.97725, pnorm(3.0) = 0.99865,

qnorm(0.01) = -2.33, qnorm(0.025) = -1.96, qnorm(0.05) = -1.64.
```

3. (4 points) What is the level L such that there is a probability of 0.05 that sample mean glucose level of four test results falls above L for Sheila's glucose level distribution?

Name: _____

Lecture Section: 0011001006 016021Lecture time: $9:05~\mathrm{am}$ $8:00~\mathrm{am}$ $10{:}10~\mathrm{am}$ 12:20 pm 11:15 am(Circle One) Zhang Yang Park Park Park

Question	Answer					
1	A	В	С	D		
2	A	В	С	D		
3	A	В	С	D		
4	A	В	С	D		
5	A	В	С	D		
6	A	В	С	D		
7	A	В	С	D		
8	A	В	С	D		
9	A	В	С	D		
10	A	В	С	D		
11	A	В	С	D		
12	A	В	С	D		
13	A	В	С	D		

Please do NOT write in the following table. This is for grading purpose only!

Question	I	II	III	IV	V	100
Score						
Total	40	13	15	20	12	100