

A Docking score matrix for 138 receptor–ligand pairs

Gold borders indicate the six representative predicted poses; positive scores are shown in grey

sesamin	-8.7	-7.5	-11.5	-9.7	-7.7	-8.9
Linarin	-8.8	-3.7	-11.0	+2.6	-8.4	-9.0
isofucosterol	-7.6	-5.9	-10.6	-5.5	-8.5	-7.3
Lucidusculine	-7.3	-6.1	-10.5	-6.1	-7.2	-7.7
Lucidumoside D	-6.8	-3.8	-10.3	-4.4	-7.3	-7.2
cholesterol	-7.5	-5.9	-10.2	-3.1	-7.4	-7.9
demethylwedelolactone	-8.6	-7.5	-10.2	-6.2	-8.1	-8.4
NSC63551	-8.5	-3.9	-10.2	-6.2	-7.3	-8.1
3'-O-Methylorobol	-8.8	-7.0	-9.4	-10.0	-7.6	-7.7
beta-sitosterol	-7.8	-3.9	-10.0	-2.2	-8.2	-7.6
butin	-8.6	-7.4	-10.0	-8.7	-7.8	-7.9
eriodictyol	-8.3	-7.4	-9.9	-8.9	-8.0	-8.3
luteolin	-8.4	-7.2	-9.9	-9.0	-8.1	-8.4
wedelolactone	-8.3	-7.4	-9.8	-6.4	-8.0	-8.3
Olitoriside	-8.1	+11.4	-9.7	+24.1	-8.0	-9.0
quercetin	-8.6	-7.5	-9.7	-8.5	-8.1	-7.6
taxifolin	-8.4	-7.7	-9.7	-8.2	-7.7	-8.2
isorhamnetin	-8.1	-6.9	-9.6	-8.3	-7.6	-7.7
acacetin	-8.1	-6.8	-9.4	-9.2	-7.5	-7.9
Pratensein	-8.5	-6.8	-9.1	-9.4	-7.5	-7.6
campest-5-en-3beta-ol	-7.9	-6.4	-9.2	-6.6	-8.2	-7.6
kaempferol	-8.2	-7.2	-9.2	-8.1	-7.7	-7.7
matrine	-7.4	-7.5	-8.4	-6.4	-6.6	-6.9
	CDK6 (3NUP)	CA2 (1BCD)	PARP1 (1UK1)	KIT (4HVS)	SYK (1XBB)	GSK3B (1Q3D)

