

**1.**

(a)

$$\begin{aligned}x'_i x'_i &= a_{ij} a_{ik} x_j x_k \\x_i x_i &= \delta_{jk} x_j x_k\end{aligned}$$

Since  $x'_i x'_i = x_i x_i$  for all  $x_i$

$$\delta_{jk} = a_{ij} a_{ik}$$

(b)

$$\begin{aligned}x_i &= \delta_{ik} x_k \\&= a_{ji} a_{jk} x_k \\&= a_{ji} x'_j\end{aligned}$$

(c)

$$\begin{aligned}\frac{\partial f}{\partial x'_i} &= \frac{\partial f}{\partial x_j} \frac{\partial x_j}{\partial x'_i} \\&= \frac{\partial f}{\partial x_j} \frac{\partial a_{kj} x'_k}{\partial x'_i} \\&= a_{ij} \frac{\partial f}{\partial x_j}\end{aligned}$$

**2.**

(a)

$$\begin{aligned}\delta'_{ij} &= a_{im} a_{jn} \delta_{mn} \\&= a_{ik} a_{jk} \\&= \delta_{ij}\end{aligned}$$

(b)

$$\begin{aligned}C'_i T'_{ij} &= a_{ik} C_k a_{im} a_{jn} T_{mn} \\&= a_{ik} a_{im} a_{jn} C_k T_{mn} \\&= \delta_{km} a_{jn} C_k T_{mn} \\&= a_{jk} C_i T_{ik} \\&= (C_i T_{ik})'\end{aligned}$$

(c)

Since both  $A_i A_j$  and  $\delta_{ij}$  are second rank tensors and  $A^2$  is a scalar,  $T_{ij}$  is also a second rank tensor.

(d)

$$\begin{aligned}
 & \partial_i \left( A_i A_j - \frac{1}{2} \delta_{ij} A_k A_k \right) \\
 &= \partial_i (A_i A_j) - \frac{1}{2} \partial_j (A_k A_k) \\
 &= A_j \partial_i (A_i) + A_i \partial_i (A_j) - A_k \partial_j (A_k) \\
 &= A_j \partial_i (A_i) + A_i \partial_i (A_j) - A_i \partial_j (A_i) \\
 &= A_j \partial_i (A_i) + A_i (\delta_{ki} \delta_{jl} - \delta_{jk} \delta_{il}) \partial_k (A_l) \\
 &= A_j \partial_i (A_i) + \varepsilon_{mij} A_i \varepsilon_{mkl} \partial_k (A_l) \\
 &= A_j \partial_i (A_i) + \varepsilon_{jmi} (\varepsilon_{mkl} \partial_k (A_l)) A_i
 \end{aligned}$$

(e)

i.

$$\begin{aligned}
 T_{ij} n_j &= \left( A_i A_j - \frac{1}{2} \delta_{ij} A^2 \right) n_j \\
 &= A_i A_j n_j - \frac{1}{2} A^2 n_i
 \end{aligned}$$

Therefore  $T \cdot \vec{n}$  is a linear combination of  $\vec{A}$  and  $\vec{n}$

ii.

Let  $B_i = T_{ij} n_j$

$$\begin{aligned}
 B_i B_i &= \left( A_i A_j n_j - \frac{1}{2} A^2 n_i \right) \left( A_i A_k n_k - \frac{1}{2} A^2 n_i \right) \\
 &= (A_j n_j)^2 A^2 + \frac{1}{4} A^4 - (A_j n_j)^2 A^2 = \frac{1}{4} A^4 \\
 B_i n_i &= A_i A_j n_j n_i - \frac{1}{2} A^2 \\
 &= A^2 \left( \cos^2 \theta - \frac{1}{2} \right) \\
 &= \frac{A^2}{2} \cos 2\theta \\
 \cos \theta_{Bn} &= \cos 2\theta \\
 \theta_{Bn} &= 2\theta
 \end{aligned}$$

iii.

See above.