

1.

(a)

In  $x, y, z$  basis

$$\begin{aligned} Q_{ij} &= \begin{pmatrix} -2d^2q & & \\ & -2d^2q & \\ & & 4d^2q \end{pmatrix} \\ &= 2d^2q \begin{pmatrix} -1 & & \\ & -1 & \\ & & 2 \end{pmatrix} \end{aligned}$$

Potential

$$\begin{aligned} \Phi &= \frac{1}{8\pi\epsilon_0} \frac{Q_{ij}x_ix_j}{r^5} \\ &= \frac{d^2q}{4\pi\epsilon_0} \frac{-x^2 - y^2 + 2z^2}{r^5} \\ &= \frac{d^2q}{4\pi\epsilon_0} \frac{3\cos^2\theta - 1}{r^3} \end{aligned}$$

Electric field

$$\begin{aligned} E_r &= \frac{3d^2q}{4\pi\epsilon_0} \frac{3\cos^2\theta - 1}{r^4} \\ E_\theta &= \frac{d^2q}{4\pi\epsilon_0} \frac{3\sin 2\theta}{r^4} \end{aligned}$$

Field line

$$\begin{aligned} \frac{dr}{d\theta} &= r \frac{3\cos^2\theta - 1}{\sin 2\theta} \\ \frac{dr}{r} &= \frac{3\cos^2\theta - 1}{\sin 2\theta} d\theta \\ r &= R_0 \sin \theta \sqrt{\cos \theta} \end{aligned}$$

The field line is the furthest when  $\frac{dr}{d\theta} = 0$

$$\theta = \arccos \frac{1}{\sqrt{3}}$$

(b)

Magnetic field

$$\begin{aligned}
 \vec{B} &= -\frac{\mu_0}{24\pi c^2 r} \hat{r} \times (\hat{r} \cdot \ddot{\vec{Q}}) \\
 &= -\frac{\mu_0}{24\pi c^2 r} \hat{r} \times \left( \hat{r} \cdot \frac{d^3 2d^2 q (3\hat{z}\hat{z} - 1)}{dt^3} \right) \\
 &= -\frac{q\mu_0 d_0^2}{12\pi c^2 r} \hat{r} \times (3(\hat{r} \cdot \hat{z})\hat{z} - \hat{r}) \frac{d^3 \cos^2 \omega t}{dt^3} \\
 &= \frac{q\mu_0 d_0^2 \omega^3}{\pi c^2 r} (\hat{r} \cdot \hat{z})(\hat{r} \times \hat{z}) \sin 2\omega t \\
 &= -\frac{q\mu_0 d_0^2 \omega^3}{2\pi c^2 r} \sin 2\theta \sin 2\omega t \hat{e}_\phi
 \end{aligned}$$

Electric field

$$\begin{aligned}
 \vec{E} &= c\vec{B} \times \vec{r} \\
 &= -\frac{q\mu_0 d_0^2 \omega^3}{2\pi c r} \sin 2\theta \sin 2\omega t \hat{e}_\theta
 \end{aligned}$$

Power

$$\frac{dP}{d\Omega} = \frac{q^2 d_0^4 \omega^6}{8\pi^2 \varepsilon_0 c^5} \sin^2 2\theta$$

The radiation is the strongest for  $\theta = \frac{\pi}{4}$  and  $\theta = \frac{3\pi}{4}$

$$\begin{aligned}
 P &= \int_0^\pi d\theta \sin \theta \int_0^{2\pi} d\phi \frac{q^2 d_0^4 \omega^6}{8\pi^2 \varepsilon_0 c^5} \sin^2 2\theta \\
 &= \frac{4q^2 d_0^4 \omega^6}{15\pi \varepsilon_0 c^5}
 \end{aligned}$$

The radiation appears at  $2\omega$

(c)

The radiation will be a dipole radiation appears at  $\omega$  with total power

$$P = \frac{q^2 d_0^2 \omega^4}{12\pi \varepsilon_0 c^3}$$

Ratio of power

$$\begin{aligned}
 \frac{P_{quad}}{P} &= \frac{16d_0^2 \omega^2}{5c^2} \\
 &\ll 1
 \end{aligned}$$

**2.**

(a)

$$\begin{aligned}
 \vec{B} &= \nabla \times A \\
 &= \frac{\mu_0}{4\pi} \int d^3\vec{r}' \nabla \times \left( \frac{1}{|\vec{r} - \vec{r}'|} \vec{j} \left( \vec{r}', t - \frac{|\vec{r} - \vec{r}'|}{c} \right) \right) \\
 &= \frac{\mu_0}{4\pi} \int d^3\vec{r}' \frac{1}{|\vec{r} - \vec{r}'|} \nabla \times \vec{j} \left( \vec{r}', t - \frac{|\vec{r} - \vec{r}'|}{c} \right) + \left( \nabla \frac{1}{|\vec{r} - \vec{r}'|} \right) \times \vec{j} \left( \vec{r}', t - \frac{|\vec{r} - \vec{r}'|}{c} \right) \\
 &= \frac{\mu_0}{4\pi} \int d^3\vec{r}' \vec{j} \left( \vec{r}', t - \frac{|\vec{r} - \vec{r}'|}{c} \right) \times \frac{\vec{r} - \vec{r}'}{|\vec{r} - \vec{r}'|^3} + \frac{\partial}{\partial t} \vec{j} \left( \vec{r}', t - \frac{|\vec{r} - \vec{r}'|}{c} \right) \times \frac{\vec{r} - \vec{r}'}{c|\vec{r} - \vec{r}'|^2}
 \end{aligned}$$

(b)

The ratio of the two term is on the order of (replacing time derivative with  $T$  and replace  $|\vec{r} - \vec{r}'|$  with  $d$ )

$$\frac{d}{cT} \ll 1$$

**3.**

(a)

$$\begin{aligned}
 P &= \frac{e^4 E_0^2}{12\pi\epsilon_0 m_e^2 c^3} \\
 &= \langle S \rangle \frac{e^4}{6\pi\epsilon_0^2 m_e^2 c^4} \\
 &= \langle S \rangle \frac{8\pi}{3} r_e^2
 \end{aligned}$$

(b)

The radiation (and therefore the stress tensor) has mirror symmetry along each axis.

(c)

$$\begin{aligned}
 F &= \frac{P}{c} \\
 &= \langle S \rangle \frac{8\pi}{3c} r_e^2
 \end{aligned}$$

**4.**

(a)

**5.**

**6.**

(a)

(b)

(c)

(d)

(e)