

1.

(a)

$$\begin{aligned}
 H &= \hbar\omega b^\dagger b + i\hbar\Lambda(b^{\dagger 2} + b^2) \\
 \frac{db}{dt} &= \frac{i}{\hbar}[H, b] + \frac{\partial b}{\partial t} \\
 &= [i\omega b^\dagger b - \Lambda b^{\dagger 2}, b] + i\omega b \\
 &= 2\Lambda b^\dagger \\
 \frac{db^\dagger}{dt} &= 2\Lambda b
 \end{aligned}$$

(b)

$$\begin{aligned}
 \vec{E} &= i\mathcal{E}_\omega \vec{\varepsilon} \left(b e^{i(\vec{k} \cdot \vec{r} - \omega t)} - b^\dagger e^{-i(\vec{k} \cdot \vec{r} - \omega t)} \right) \\
 &= i\mathcal{E}_\omega \vec{\varepsilon} \left(b \cos(\vec{k} \cdot \vec{r} - \omega t) + ib \sin(\vec{k} \cdot \vec{r} - \omega t) - b^\dagger \cos(\vec{k} \cdot \vec{r} - \omega t) + ib^\dagger \sin(\vec{k} \cdot \vec{r} - \omega t) \right) \\
 &= -2\mathcal{E}_\omega \vec{\varepsilon} \left(\frac{b - b^\dagger}{2i} \cos(\vec{k} \cdot \vec{r} - \omega t) + \frac{b + b^\dagger}{2} \sin(\vec{k} \cdot \vec{r} - \omega t) \right) \\
 &= -2\mathcal{E}_\omega \vec{\varepsilon} \left(b_Q \cos(\vec{k} \cdot \vec{r} - \omega t) + b_P \sin(\vec{k} \cdot \vec{r} - \omega t) \right) \\
 \frac{db_\pm}{dt} &\equiv \frac{db \pm b^\dagger}{dt} \\
 &= 2\Lambda b^\dagger \pm 2\Lambda b \\
 &= \pm 2\Lambda b_\pm \\
 \frac{db_P}{dt} &= 2\Lambda b_P \\
 \frac{db_Q}{dt} &= -2\Lambda b_Q
 \end{aligned}$$

Therefore

$$\begin{aligned}
 b_P &= e^{2\Lambda t} b_{P0} \\
 b_Q &= e^{-2\Lambda t} b_{Q0} \\
 b &= b_P + ib_Q \\
 &= e^{2\Lambda t} b_{P0} + ie^{-2\Lambda t} b_{Q0} \\
 &= b_0 \cosh 2\Lambda t + b_0^\dagger \sinh 2\Lambda t \\
 b^\dagger &= b_0^\dagger \cosh 2\Lambda t + b_0 \sinh 2\Lambda t
 \end{aligned}$$

(c)


$$\begin{aligned}
 \langle N \rangle &= \langle 0 | b^\dagger b | 0 \rangle \\
 &= \langle 0 | \left(b_0^\dagger \cosh 2\Lambda t + b_0 \sinh 2\Lambda t \right) \left(b_0 \cosh 2\Lambda t + b_0^\dagger \sinh 2\Lambda t \right) | 0 \rangle \\
 &= \sinh^2 2\Lambda t \\
 \Delta b_P &= e^{2\Lambda t} \Delta b_{P0} \\
 &= \frac{1}{2} e^{2\Lambda t} \\
 \Delta b_Q &= e^{-2\Lambda t} \Delta b_{Q0} \\
 &= \frac{1}{2} e^{-2\Lambda t}
 \end{aligned}$$

The state is squeezed in Q direction while the product of the uncertainty in P and Q remains the same.

(d)


(e)

i.




ii.


$$\beta = i$$



iii.

$$\beta = i$$





Whether a phase or number squeezed state is created depends on the initial state.

2.

(a)

(b)

(c)

i.

ii.

iii.

iv.

3.

(a)

(b)

(c)

(d)