

# Building Single Molecules from Single Atoms

A DISSERTATION PRESENTED  
BY  
YICHAO YU  
TO  
THE DEPARTMENT OF PHYSICS

IN PARTIAL FULFILLMENT OF THE REQUIREMENTS  
FOR THE DEGREE OF  
DOCTOR OF PHILOSOPHY  
IN THE SUBJECT OF  
PHYSICS

HARVARD UNIVERSITY  
CAMBRIDGE, MASSACHUSETTS  
MARCH 2021

©2021 – YICHAO YU  
ALL RIGHTS RESERVED.

Thesis advisor: Professor Kang-Kuen Ni

Yichao Yu

# Building Single Molecules from Single Atoms

ABSTRACT

# Contents

o	INTRODUCTION	<b>I</b>
1	APPARATUS	<b>2</b>
1.1	Cooling and optical pumping beams . . . . .	2
1.2	Tweezer and imaging . . . . .	3
1.3	Molecular Raman frequency generation . . . . .	3
2	COMPUTER CONTROL OF THE EXPERIMENT	<b>4</b>
2.1	Overall structure . . . . .	4
2.2	Frontend . . . . .	4
2.3	Backends . . . . .	4
2.4	Automation of scan . . . . .	5
3	RAMAN SIDEBAND COOLING	<b>6</b>
3.1	Theory . . . . .	6
3.2	Setup . . . . .	6
3.3	Challenge with large Lamb-Dicky parameter . . . . .	7
3.4	Solution: High order sidebands . . . . .	7
3.5	Solution: Simulation based optimization . . . . .	7
3.6	Cooling performance . . . . .	7
4	INTERACTION OF SINGLE ATOMS	<b>9</b>
4.1	Scattering length . . . . .	9
4.2	Energy levels of two interacting atoms in an anisotropic trap . . . . .	10
4.3	Interaction shift spectroscopy . . . . .	10
4.4	Summary and Outlook . . . . .	10
5	PHOTOASSOCIATION OF SINGLE ATOMS	<b>11</b>
5.1	Energy levels . . . . .	11
5.2	Effect of the trap . . . . .	11
5.3	Photoassociation spectroscopy . . . . .	12
6	TWO-PHOTON SPECTROSCOPY OF NACs GROUND STATE	<b>13</b>
7	COHERENT OPTICAL CREATION OF NACs MOLECULE	<b>14</b>





# Acknowledgments

,

# 0

## Introduction



# 1

## Apparatus

### 1.1 COOLING AND OPTICAL PUMPING BEAMS

(MOT, OP, fiber back reflection)

(Mention Na Raman beam to be covered in later chapter?)

## 1.2 TWEEZER AND IMAGING

## 1.3 MOLECULAR RAMAN FREQUENCY GENERATION

(beam path, calibration)

# 2

## Computer control of the experiment

### 2.1 OVERALL STRUCTURE

### 2.2 FRONTEND

### 2.3 BACKENDS

(communication protocol)

2.3.1 FPGA BACKEND

2.3.2 NIDAQ BACKEND

2.3.3 USRP BACKEND

2.4 AUTOMATION OF SCAN

# 3

## Raman sideband cooling

### 3.1 THEORY

#### 3.1

### 3.2 SETUP

#### 3.2




**Figure 3.1:** Single Na atom Raman sideband cooling scheme. The Raman transitions between  $|2, 2; n\rangle$  and  $|1, 1; n + \Delta n\rangle$  have a one-photon detuning  $\Delta = 75$  GHz below the  $3^2S_{1/2}$  to  $3^2P_{3/2}$  transition. Two-photon detuning,  $\delta$ , is defined relative to the  $\Delta n = 0$  carrier transition. For optical pumping, we use two  $\sigma^+$  polarized transitions, one to pump the atom state out of  $|1, 1\rangle$  via  $3^2P_{3/2}$  and one to pump atoms out of  $|2, 1\rangle$  via  $3^2P_{1/2}$  to minimize heating of the  $|2, 2\rangle$  state.

### 3.3 CHALLENGE WITH LARGE LAMB-DICKY PARAMETER

### 3.4 SOLUTION: HIGH ORDER SIDEBANDS

### 3.5 SOLUTION: SIMULATION BASED OPTIMIZATION

### 3.6 COOLING PERFORMANCE



**Figure 3.2:** Geometry and polarizations of the Raman and optical pumping beams relative to the optical tweezer and bias magnetic field. Raman beams R1 and R4 address the radial  $x$ -mode. R1 and R2 address the radial  $y$ -mode. R3 and R4 address the axial  $z$ -mode, where the beams also couple to radial motion, but this coupling can be neglected when the atoms are cooled to the ground state of motion.

# 4

## Interaction of single atoms

### 4.1 SCATTERING LENGTH

(Importance/relation with binding energy etc.)



## 4.2 ENERGY LEVELS OF TWO INTERACTING ATOMS IN AN ANISOTROPIC TRAP

## 4.3 INTERACTION SHIFT SPECTROSCOPY

(motional sideband, scattering length result)

## 4.4 SUMMARY AND OUTLOOK

(Motional state selection)

# 5

## Photoassociation of single atoms

### 5.1 ENERGY LEVELS

### 5.2 EFFECT OF THE TRAP

(light shift, broadening)

### 5.3 PHOTOASSOCIATION SPECTROSCOPY

( $v=0, 12, 14$ , etc)

# 6

## Two-photon spectroscopy of NaCs ground state

(N=2, different HF states)

# 7

Coherent optical creation of NaCs

molecule

# 8

## Conclusion