

Doppler-free spectroscopy using saturated absorption

Yichao Yu

MIT

April 28, 2013

Introduction

- Saturated absorption.
- Precise spectrum measurement.
- Frequency stabilization and locking.

Introduction

- Saturated absorption.
- Precise spectrum measurement.
- Frequency stabilization and locking.

Introduction

- Saturated absorption.
- Precise spectrum measurement.
- Frequency stabilization and locking.

1 Doppler broadening and saturated absorption.

2 Apparatus and measurement.

3 Data and result.

4 Conclusion.

Doppler broadening and spectral hole burning.

- Natural line width.
- Doppler broadening.

$$\Delta f = \frac{v}{c} f_0 \approx 100 \text{ MHz}$$

- Cooling.
- He-Ne spectral hole burning discovered by Bennett in 1962.
- Saturated absorption in vapor cell.

Doppler broadening and spectral hole burning.

- Natural line width.
- Doppler broadening.

$$\Delta f = \frac{v}{c} f_0 \approx 100 \text{ MHz}$$

- Cooling.
- He-Ne spectral hole burning discovered by Bennett in 1962.
- Saturated absorption in vapor cell.

Doppler broadening and spectral hole burning.

- Natural line width.
- Doppler broadening.

$$\Delta f = \frac{v}{c} f_0 \approx 100 \text{ MHz}$$


- Cooling.
- He-Ne spectral hole burning discovered by Bennett in 1962.
- Saturated absorption in vapor cell.

Doppler broadening and spectral hole burning.

- Natural line width.
- Doppler broadening.

$$\Delta f = \frac{v}{c} f_0 \approx 100 \text{ MHz}$$

- Cooling.
- He-Ne spectral hole burning discovered by Bennett in 1962.
- Saturated absorption in vapor cell.




Doppler broadening and spectral hole burning.

- Natural line width.
- Doppler broadening.

$$\Delta f = \frac{v}{c} f_0 \approx 100 \text{ MHz}$$

- Cooling.
- He-Ne spectral whole burning discovered by Bennett in 1962.
- Saturated absorption in vapor cell.




Doppler broadening and spectral hole burning.

- Natural line width.
- Doppler broadening.


$$\Delta f = \frac{v}{c} f_0 \approx 100 \text{ MHz}$$

- Cooling.
- He-Ne spectral hole burning discovered by Bennett in 1962.
- Saturated absorption in vapor cell.




Saturated absorption.

- Saturation intensity.
- Saturate absorption with two beams and Lamb dip.
- Crossover peaks.




Saturated absorption.

- Saturation intensity.
- Saturate absorption with two beams and Lamb dip.
- Crossover peaks.




Saturated absorption.

- Saturation intensity.
- Saturate absorption with two beams and Lamb dip.
- Crossover peaks.




Saturated absorption.

- Saturation intensity.
- Saturate absorption with two beams and Lamb dip.
- Crossover peaks.




_____ G

Apparatus.




- Pump beam
- Probe and reference beam.
- Fabry-Pérot cavity

Apparatus.




- Pump beam
- Probe and reference beam.
- Fabry-Pérot cavity

Apparatus.



- Pump beam
- Probe and reference beam.
- Fabry-Pérot cavity

Apparatus.



- Pump beam
- Probe and reference beam.
- Fabry-Pérot cavity

Hyperfine structure of Rubidium D_2 line.

- D_2 line

$$5^2S_{1/2} \rightarrow 5^2P_{3/2}$$

- 2 “doppler distinguishable” “ground states”, each couples with 3 non-“doppler distinguishable” excited states.
-

$$\Delta E_{hfs} = \frac{1}{2}A_{hfs}K + B_{hfs}\frac{3K(K+1) - 2I(I+1)J(J+1)}{8I(2I-1)J(2J-1)}$$

$$K = F(F+1) - I(I+1) - J(J+1)$$

- Quantities to measure

$$A_{5^2S_{1/2}}, A_{5^2P_{3/2}} \text{ and } B_{5^2P_{3/2}} \text{ for } ^{85}\text{Rb} \text{ and } ^{87}\text{Rb}.$$

Hyperfine structure of Rubidium D_2 line.

- D_2 line
 $5^2S_{1/2} \rightarrow 5^2P_{3/2}$
- 2 “doppler distinguishable” “ground states”, each couples with 3 non-“doppler distinguishable” excited states.

$$\Delta E_{hfs} = \frac{1}{2}A_{hfs}K + B_{hfs}\frac{3K(K+1) - 2I(I+1)J(J+1)}{8I(2I-1)J(2J-1)}$$
$$K = F(F+1) - I(I+1) - J(J+1)$$

- Quantities to measure

$A_{5^2S_{1/2}}$, $A_{5^2P_{3/2}}$ and $B_{5^2P_{3/2}}$ for ^{85}Rb and ^{87}Rb .

Hyperfine structure of Rubidium D_2 line.

- D_2 line
 $5^2S_{1/2} \rightarrow 5^2P_{3/2}$
- 2 “doppler distinguishable” “ground states”, each couples with 3 non-“doppler distinguishable” excited states.

-

$$\Delta E_{hfs} = \frac{1}{2}A_{hfs}K + B_{hfs} \frac{3K(K+1) - 2I(I+1)J(J+1)}{8I(2I-1)J(2J-1)}$$

$$K = F(F+1) - I(I+1) - J(J+1)$$

- Quantities to measure

$A_{5^2S_{1/2}}$, $A_{5^2P_{3/2}}$ and $B_{5^2P_{3/2}}$ for ^{85}Rb and ^{87}Rb .

Hyperfine structure of Rubidium D_2 line.

- D_2 line
 $5^2S_{1/2} \rightarrow 5^2P_{3/2}$
- 2 “doppler distinguishable” “ground states”, each couples with 3 non-“doppler distinguishable” excited states.


-

$$\Delta E_{hfs} = \frac{1}{2}A_{hfs}K + B_{hfs} \frac{3K(K+1) - 2I(I+1)J(J+1)}{8I(2I-1)J(2J-1)}$$

$$K = F(F+1) - I(I+1) - J(J+1)$$


- Quantities to measure
 $A_{5^2S_{1/2}}$, $A_{5^2P_{3/2}}$ and $B_{5^2P_{3/2}}$ for ^{85}Rb and ^{87}Rb .

Spectrum measurement.




Probe beam intensity of the whole scan.

Spectrum measurment.



Probe beam intensity of the whole scan.



Difference between probe and reference beam intensity of the ^{87}Rb $F = 2$ lines.

Hyperfine Structure Constants.

Hyperfine Structure Constants.

Isotope	Constant	Measured	Expected	Deviation
^{85}Rb	$A_{5^2S_{1/2}}$	$0.986(40)\text{GHz}$	1.0119GHz	0.6σ
	$A_{5^2P_{3/2}}$	$24.44(81)\text{MHz}$	$25.0020(99)\text{MHz}$	0.7σ
	$B_{5^2P_{3/2}}$	$32.2(4.8)\text{MHz}$	$25.790(93)\text{MHz}$	1.3σ
^{87}Rb	$A_{5^2S_{1/2}}$	$3.285(65)\text{GHz}$	3.4173GHz	2σ
	$A_{5^2P_{3/2}}$	$84.58(52)\text{MHz}$	$84.7185(20)\text{MHz}$	0.2σ
	$B_{5^2P_{3/2}}$	$16.03(80)\text{MHz}$	$12.4965(37)\text{MHz}$	4σ

Conclusion.

- Observed Lamb dip.
- Measured hyperfine splitting and the hyperfine structure constants.

Conclusion.

- Observed Lamb dip.
- Measured hyperfine splitting and the hyperfine structure constants.

