

# Mössbauer Spectroscopy


Yichao Yu

MIT

March 6, 2013

# Mössbauer effect and Mössbauer spectroscopy.


- Nuclear  $\gamma$  spectrum.
- Simple setup.
- Super high resolution. ( $10^{12}$ )



Mössbauer spectrum of  $\text{FeC}_2\text{O}_4$ .

## Mössbauer effect and Mössbauer spectroscopy.


- Nuclear  $\gamma$  spectrum.
- Simple setup.
- Super high resolution. ( $10^{12}$ )



Mössbauer spectrum of  $\text{FeC}_2\text{O}_4$ .

## Mössbauer effect and Mössbauer spectroscopy.

- Nuclear  $\gamma$  spectrum.
- Simple setup.
- Super high resolution. ( $10^{12}$ )



Mössbauer spectrum of  $FeC_2O_4$ .


- 1 **Mössbauer effect.**
- 2 **Apparatus and samples.**
- 3 **Data and result.**
- 4 **Conclusion.**

## Nuclear spectrum and recoil.

- Radio active element radiate  $\gamma$ -ray at characteristic frequencies.
- Radiation  $\rightarrow$  Absorption.
- Recoil momentum and doppler shift.


## Nuclear spectrum and recoil.

- Radio active element radiate  $\gamma$ -ray at characteristic frequencies.
- Radiation  $\rightarrow$  Absorption.
- Recoil momentum and doppler shift.



## Nuclear spectrum and recoil.

- Radio active element radiate  $\gamma$ -ray at characteristic frequencies.
- Radiation  $\rightarrow$  Absorption.
- Recoil momentum and doppler shift.





# Mössbauer effect.


- Discovered in 1957 by Mössbauer.
- Momentum transfer with the whole crystal.  $\rightarrow$  Recoilless.
- Scanning frequency using doppler effect.

# Mössbauer effect.


- Discovered in 1957 by Mössbauer.
- Momentum transfer with the whole crystal.  $\rightarrow$  Recoilless.
- Scanning frequency using doppler effect.

# Mössbauer effect.


- Discovered in 1957 by Mössbauer.
- Momentum transfer with the whole crystal.  $\rightarrow$  Recoilless.
- Scanning frequency using doppler effect.




## Experiment Apparatus.




## Experiment Apparatus.



## Experiment Apparatus.



## Experiment Apparatus.




- Velocity of the source,

$$v = 2f\lambda$$

- Source:  $^{57}\text{Co}$ .
- First decay into excited state of  $^{57}\text{Fe}$ .
- Energy  $14.4\text{keV}$ .

## Experiment Apparatus.




- Velocity of the source,

$$v = 2f\lambda$$

- Source:  $^{57}\text{Co}$ .
- First decay into excited state of  $^{57}\text{Fe}$ .
- Energy  $14.4\text{keV}$ .



## Experiment Apparatus.




- Velocity of the source,

$$v = 2f\lambda$$

- Source:  $^{57}\text{Co}$ .
- First decay into excited state of  $^{57}\text{Fe}$ .
- Energy  $14.4\text{keV}$ .

## Experiment Apparatus.



- Velocity of the source,

$$v = 2f\lambda$$

- Source:  $^{57}\text{Co}$ .
- First decay into excited state of  $^{57}\text{Fe}$ .
- Energy  $14.4\text{keV}$ .

# Effects and samples.


## Effects

- Isomer Shift.
- Zeeman effect.

$$E = -g_N \mu_N B m_I$$

- Quadrupole Splitting.
- Temperature Shift.

$$\frac{\delta}{E} = \frac{v^2}{2c^2} = \frac{E_k}{mc^2}$$



# Effects and samples.


## Effects

- Isomer Shift.
- Zeeman effect.

$$E = -g_N \mu_N B m_I$$

- Quadrupole Splitting.
- Temperature Shift.

$$\frac{\delta}{E} = \frac{v^2}{2c^2} = \frac{E_k}{mc^2}$$



# Effects and samples.


## Effects

- Isomer Shift.
- Zeeman effect.

$$E = -g_N \mu_N B m_I$$

- Quadrapole Splitting.
- Temperature Shift.

$$\frac{\delta}{E} = \frac{v^2}{2c^2} = \frac{E_k}{mc^2}$$



# Effects and samples.


## Effects

- Isomer Shift.
- Zeeman effect.

$$E = -g_N \mu_N B m_I$$

- Quadrapole Splitting.
- Temperature Shift.

$$\frac{\delta}{E} = \frac{v^2}{2c^2} = \frac{E_k}{mc^2}$$



# Effects and samples.

## Effects

- Isomer Shift.
- Zeeman effect.

$$E = -g_N \mu_N B m_I$$

- Quadrapole Splitting.
- Temperature Shift.

$$\frac{\delta}{E} = \frac{v^2}{2c^2} = \frac{E_k}{mc^2}$$

## Samples

- $^{57}\text{Fe}$
- $\text{FeSO}_4$
- $\text{Fe}_2(\text{SO}_4)_3$
- $\text{Fe}_2\text{O}_3$
- Stainless steel (Varying temperature).
- $\text{Na}_4\text{Fe}(\text{CN})_6$  For line width.

# Effects and samples.

## Effects

- Isomer Shift.
- Zeeman effect.

$$E = -g_N \mu_N B m_I$$

- Quadrapole Splitting.
- Temperature Shift.

$$\frac{\delta}{E} = \frac{v^2}{2c^2} = \frac{E_k}{mc^2}$$

## Samples

- $^{57}\text{Fe}$
- $\text{FeSO}_4$
- $\text{Fe}_2(\text{SO}_4)_3$
- $\text{Fe}_2\text{O}_3$
- Stainless steel (Varying temperature).
- $\text{Na}_4\text{Fe}(\text{CN})_6$  For line width.



# Effects and samples.

## Effects

- Isomer Shift.
- Zeeman effect.

$$E = -g_N \mu_N B m_I$$

- Quadrapole Splitting.
- Temperature Shift.

$$\frac{\delta}{E} = \frac{v^2}{2c^2} = \frac{E_k}{mc^2}$$

## Samples

- $^{57}\text{Fe}$
- $\text{FeSO}_4$
- $\text{Fe}_2(\text{SO}_4)_3$
- $\text{Fe}_2\text{O}_3$
- Stainless steel (Varying temperature).
- $\text{Na}_4\text{Fe}(\text{CN})_6$  For line width.

# Effects and samples.

## Effects

- Isomer Shift.
- Zeeman effect.

$$E = -g_N \mu_N B m_I$$

- Quadrapole Splitting.
- Temperature Shift.

$$\frac{\delta}{E} = \frac{v^2}{2c^2} = \frac{E_k}{mc^2}$$

## Samples

- $^{57}\text{Fe}$
- $\text{FeSO}_4$
- $\text{Fe}_2(\text{SO}_4)_3$
- $\text{Fe}_2\text{O}_3$
- Stainless steel (Varying temperature).
- $\text{Na}_4\text{Fe}(\text{CN})_6$  For line width.

# Effects and samples.

## Effects

- Isomer Shift.
- Zeeman effect.

$$E = -g_N \mu_N B m_I$$

- Quadrapole Splitting.
- **Temperature Shift.**

$$\frac{\delta}{E} = \frac{v^2}{2c^2} = \frac{E_k}{mc^2}$$

## Samples

- $^{57}\text{Fe}$
- $\text{FeSO}_4$
- $\text{Fe}_2(\text{SO}_4)_3$
- $\text{Fe}_2\text{O}_3$
- Stainless steel (Varying temperature).
- $\text{Na}_4\text{Fe}(\text{CN})_6$  For line width.

# Effects and samples.

## Effects

- Isomer Shift.
- Zeeman effect.

$$E = -g_N \mu_N B m_I$$

- Quadrapole Splitting.
- Temperature Shift.

$$\frac{\delta}{E} = \frac{v^2}{2c^2} = \frac{E_k}{mc^2}$$

## Samples

- $^{57}\text{Fe}$
- $\text{FeSO}_4$
- $\text{Fe}_2(\text{SO}_4)_3$
- $\text{Fe}_2\text{O}_3$
- Stainless steel (Varying temperature).
- $\text{Na}_4\text{Fe}(\text{CN})_6$  For line width.

# Effects and samples.

## Effects

- Isomer Shift.
- Zeeman effect.

$$E = -g_N \mu_N B m_I$$


- Quadrapole Splitting.
- Temperature Shift.

$$\frac{\delta}{E} = \frac{v^2}{2c^2} = \frac{E_k}{mc^2}$$



## Samples

- $^{57}\text{Fe}$
- $\text{FeSO}_4$
- $\text{Fe}_2(\text{SO}_4)_3$
- $\text{Fe}_2\text{O}_3$
- Stainless steel (Varying temperature).
- $\text{Na}_4\text{Fe}(\text{CN})_6$  For line width.

# Calibration.




# Calibration.



# $^{57}\text{Fe}$ and $\text{FeSO}_4$ .

$^{57}\text{Fe}$




**Zeeman splitting for ground state ( $g_0$ ) and excited state ( $g_1$ ).**

$g_0$	$1.882(13) \cdot 10^{-7} eV$
$g_1$	$1.074(13) \cdot 10^{-7} eV$




# $^{57}\text{Fe}$ and $\text{FeSO}_4$ .

$^{57}\text{Fe}$



$\text{FeSO}_4$



**Zeeman splitting for ground state ( $g_0$ ) and excited state ( $g_1$ ).**

$g_0$	$1.882(13) \cdot 10^{-7} \text{eV}$
$g_1$	$1.074(13) \cdot 10^{-7} \text{eV}$

**Isomer shift ( $\epsilon$ ) and quadrupole splitting ( $\delta$ ).**

$\epsilon$	$5.9(1.4) \cdot 10^{-8} \text{eV}$
$\delta$	$1.562(2) \cdot 10^{-7} \text{eV}$

# Temperature effect.

## Samples

- Classic:

$$E_k = \frac{3}{2}k_B T$$

- Debye  $T^3$  approximation:

$$E_k = \frac{3\pi^4 * k_B * T^4}{10\Theta_D^3}$$

- Exact Debye Model:

$$E_k = \frac{9k_B T^4}{10\Theta_D^3} D_3 \left( \frac{T}{\Theta_D} \right)$$

# Temperature effect.

## Samples

- Classic:

$$E_k = \frac{3}{2}k_B T$$

- Debye  $T^3$  approximation:

$$E_k = \frac{3\pi^4 * k_B * T^4}{10\Theta_D^3}$$

- Exact Debye Model:

$$E_k = \frac{9k_B T^4}{10\Theta_D^3} D_3 \left( \frac{T}{\Theta_D} \right)$$

# Temperature effect.

## Samples

- Classic:

$$E_k = \frac{3}{2}k_B T$$

- Debye  $T^3$  approximation:

$$E_k = \frac{3\pi^4 * k_B * T^4}{10\Theta_D^3}$$

- Exact Debye Model:

$$E_k = \frac{9k_B T^4}{10\Theta_D^3} D_3 \left( \frac{T}{\Theta_D} \right)$$

# Temperature effect.

## Samples

- Classic:

$$E_k = \frac{3}{2}k_B T$$

- Debye  $T^3$  approximation:

$$E_k = \frac{3\pi^4 * k_B * T^4}{10\Theta_D^3}$$

- Exact Debye Model:

$$E_k = \frac{9k_B T^4}{10\Theta_D^3} D_3 \left( \frac{T}{\Theta_D} \right)$$

Model	$E_k$
Classic	$1.409(12) \cdot 10^{-2} eV$
Debye $T^3$	$4.59(22) \cdot 10^{-1} eV$
Exact Debye	$1.304(84) \cdot 10^{-2} eV$
(Measured)	$0.99(36) \cdot 10^{-2} eV$

## Conclusion.

- Calibrated the velocity using laser.
- Measured Mössbauer spectrum of a variety of materials.
- Calculated different kinds of splitting and shifting.

## Conclusion.

- Calibrated the velocity using laser.
- Measured Mössbauer spectrum of a variety of materials.
- Calculated different kinds of splitting and shifting.

## Conclusion.

- Calibrated the velocity using laser.
- Measured Mössbauer spectrum of a variety of materials.
- Calculated different kinds of splitting and shifting.





## Effects

- Isomer Shift:

$\varepsilon$

- Zeeman effect:

$g_0, g_1$

- Quadrupole Splitting:

$\delta$

$$\Delta E_1 = \varepsilon - \frac{g_0}{2} - \frac{3g_1}{2} - \frac{\delta}{2} \quad (1)$$

$$\Delta E_2 = \varepsilon - \frac{g_0}{2} - \frac{g_1}{2} + \frac{\delta}{2} \quad (2)$$

$$\Delta E_3 = \varepsilon - \frac{g_0}{2} + \frac{g_1}{2} + \frac{\delta}{2} \quad (3)$$

$$\Delta E_4 = \varepsilon + \frac{g_0}{2} - \frac{g_1}{2} + \frac{\delta}{2} \quad (4)$$

$$\Delta E_5 = \varepsilon + \frac{g_0}{2} + \frac{g_1}{2} + \frac{\delta}{2} \quad (5)$$

$$\Delta E_6 = \varepsilon + \frac{g_0}{2} + \frac{3g_1}{2} - \frac{\delta}{2} \quad (6)$$

