

# Coherent optical association of a single molecule – Supplemental material

Yichao Yu,<sup>1,2,3,\*</sup> Kenneth Wang,<sup>1,2,3</sup> Jonathan D. Hood,<sup>4</sup> Lewis R. B. Picard,<sup>1,2,3</sup> Jessie T. Zhang,<sup>1,2,3</sup>  
William B. Cairncross,<sup>2,1,3</sup> Jeremy M. Hutson,<sup>5</sup> Till Rosenband,<sup>1</sup> and Kang-Kuen Ni<sup>2,1,3,†</sup>

<sup>1</sup>*Department of Physics, Harvard University, Cambridge, Massachusetts 02138, USA*

<sup>2</sup>*Department of Chemistry and Chemical Biology, Harvard University, Cambridge, Massachusetts 02138, USA*

<sup>3</sup>*Harvard-MIT Center for Ultracold Atoms, Cambridge, Massachusetts 02138, USA*

<sup>4</sup>*Department of Chemistry, Purdue University, West Lafayette, Indiana, 47906, USA*

<sup>5</sup>*Joint Quantum Centre Durham-Newcastle, Department of Chemistry,  
Durham University, Durham, DH1 3LE, United Kingdom*

(Dated: November 24, 2020)

## S1. 3 LEVEL RAMAN TRANSFER WITH CROSS COUPLING

In our system, the separation in energy between the initial and target state is much smaller than the single photon detuning  $\Delta$  from the intermediate state. Thus, there is significant cross coupling for the scattering rate and light shift, where each state is coupled to the excited state by the power in both beams. Under these conditions, the scattering rate and light shift is proportional to the total power  $P_{\text{tot}} = P_m + P_a$ , where  $P_m(P_a)$  is the power in the beam that addresses the molecular (atomic) state. The Raman Rabi frequency is proportional to  $\sqrt{P_m P_a}$ . With a fixed total power and thus fixed scattering rate, the Raman Rabi frequency is maximized when  $P_m = P_a = P_{\text{tot}}/2$ .

For the purposes of this work, we denote  $\Omega_a(\Omega_m)$  as the Rabi frequencies to drive a transition from the atom (target molecule) state to the intermediate state used in the Raman transition corresponding to the power in a single beam,  $P_a$  ( $P_m$ ). Thus, the Raman Rabi frequency is  $\Omega_a \Omega_m / 2\Delta$ , where  $\Delta$  is the single photon detuning. Cross coupling affects the scattering rate and light shift, and assuming equal powers in both beams, these quantities become  $\Gamma_e \Omega_m^2 / 2\Delta^2 + \Gamma_e \Omega_a^2 / 2\Delta^2$  and  $\Omega_m^2 / 2\Delta - \Omega_a^2 / 2\Delta$ , where  $\Gamma_e$  is the excited state linewidth. Since  $\Omega_m \gg \Omega_a$ , the scattering rate and light shift can be approximated as  $\Gamma_e \Omega_m^2 / 2\Delta^2$  and  $\Omega_m^2 / 2\Delta$ .

## S2. RAMAN TRANSFER WITH MANY INTERMEDIATE STATES

With multiple intermediate states, the total Raman Rabi frequency, scattering rate, and light shift become a sum over all these possible states. In our model, we consider all vibrational states of  $\text{c}^3\Sigma^+(\Omega = 1)$  and states in the atomic continuum as intermediate states. The Rabi frequency between two states is proportional to the electronic dipole moment, arising from the electronic wavefunctions, and the overlap of the vibrational wavefunctions. Under the assumption that the electronic dipole moment  $d$  is the same for all transitions between the ground state and the excited electronic state, the total Raman Rabi frequency becomes:

$$\Omega_R = \frac{d^2 E^2}{2\hbar^2} \sum_{v'} \frac{\langle a|v'\rangle \langle v'|m\rangle}{\Delta_{v'}} \quad (\text{S.1})$$

where  $|a\rangle$  ( $|m\rangle$ ) are the relative coordinate wavefunctions for the trapped atomic (target molecular) state, and  $|v'\rangle$  is the relative-coordinate wavefunction for the excited state which includes both bound molecular states and the atomic continuum.  $\Delta_{v'}$  is a single photon detuning with respect to the  $|v'\rangle$  state,  $E$  is the electric field for the power in the single beam given a waist size. We can divide this sum into a sum over all the bound molecular states, and a sum over atomic continuum states.

$$\Omega_R = \frac{d^2 E^2}{2\hbar^2} \sum_{v' \leq v'_{\text{max}}} \frac{\langle a|v'\rangle \langle v'|m\rangle}{\Delta_{v'}} + \frac{d^2 E^2}{2\hbar^2} \sum_{v' > v'_{\text{max}}} \frac{\langle a|v'\rangle \langle v'|m\rangle}{\Delta_{v'}} \quad (\text{S.2})$$

where  $v'_{\text{max}}$  corresponds to the excited molecular state, which is most weakly bound. The first term can be calculated numerically given an excited state potential. To calculate the second term, we first make an assumption that we are

---

\*Electronic address: [yichaoyu@g.harvard.edu](mailto:yichaoyu@g.harvard.edu)

†Electronic address: [ni@chemistry.harvard.edu](mailto:ni@chemistry.harvard.edu)

far enough detuned from the contributing continuum states that they are all at approximately the same detuning,  $\delta_{\text{thresh}}$ . This allows us to pull it out of the sum, and we obtain:

$$\Omega_R \approx \frac{d^2 E^2}{2\hbar^2} \sum_{v' \leq v'_{\text{max}}} \frac{\langle a|v'\rangle \langle v'|m\rangle}{\Delta_{v'}} + \frac{d^2 E^2}{2\hbar^2 \delta_{\text{thresh}}} \sum_{v' > v'_{\text{max}}} \langle a|v'\rangle \langle v'|m\rangle \quad (\text{S.3})$$

From requiring orthogonality,

$$\sum_{v'} \langle a|v'\rangle \langle v'|m\rangle = \langle a|m\rangle = 0 \quad (\text{S.4})$$

we derive an expression for the second sum in equation S.3

$$\sum_{v' > v'_{\text{max}}} \langle a|v'\rangle \langle v'|m\rangle = - \sum_{v' \leq v'_{\text{max}}} \langle a|v'\rangle \langle v'|m\rangle \quad (\text{S.5})$$

which can be calculated from the vibrational states of the excited state potential. Thus, the Raman Rabi frequency can be approximated as:

$$\Omega_R \approx \frac{d^2 E^2}{2\hbar^2} \sum_{v' \leq v'_{\text{max}}} \frac{\langle a|v'\rangle \langle v'|m\rangle}{\Delta_{v'}} - \frac{d^2 E^2}{2\hbar^2 \delta_{\text{thresh}}} \sum_{v' \leq v'_{\text{max}}} \langle a|v'\rangle \langle v'|m\rangle \quad (\text{S.6})$$

We can account for the atomic continuum in the scattering rate in a similar way.

$$\Gamma_s = \frac{d^2 E^2}{2\hbar^2} \sum_{v' \leq v'_{\text{max}}} \Gamma_{v'} \frac{\langle m|v'\rangle \langle v'|m\rangle}{\Delta_{v'}^2} + \frac{d^2 E^2}{2\hbar^2} \sum_{v' > v'_{\text{max}}} \Gamma_{v'} \frac{\langle m|v'\rangle \langle v'|m\rangle}{\Delta_{v'}^2} \quad (\text{S.7})$$

$$\approx \frac{d^2 E^2}{2\hbar^2} \sum_{v' \leq v'_{\text{max}}} \Gamma_{v'} \frac{\langle m|v'\rangle \langle v'|m\rangle}{\Delta_{v'}^2} + \frac{d^2 E^2 \Gamma}{2\hbar^2 \delta_{\text{thresh}}^2} \sum_{v' > v'_{\text{max}}} \langle m|v'\rangle \langle v'|m\rangle \quad (\text{S.8})$$

where  $\Gamma_{v'}$  is the excited state linewidth for the  $|v'\rangle$  state, and we have assumed the atomic continuum states all have the same linewidth  $\Gamma$ . Using normalization, we derive an expression for the sum in the second term

$$\sum_{v' > v'_{\text{max}}} \langle m|v'\rangle \langle v'|m\rangle = 1 - \sum_{v' \leq v'_{\text{max}}} \langle m|v'\rangle \langle v'|m\rangle \quad (\text{S.9})$$

We then arrive at an expression for the scattering rate, accounting for the atomic continuum,

$$\Gamma_s \approx \frac{d^2 E^2}{2\hbar^2} \sum_{v' \leq v'_{\text{max}}} \Gamma_{v'} \frac{\langle m|v'\rangle \langle v'|m\rangle}{\Delta_{v'}^2} + \frac{d^2 E^2 \Gamma}{2\hbar^2 \delta_{\text{thresh}}^2} \left( 1 - \sum_{v' \leq v'_{\text{max}}} \langle m|v'\rangle \langle v'|m\rangle \right) \quad (\text{S.10})$$

We have replaced the full continuum with an effective state at the atomic threshold with strength proportional to the amount of wavefunction overlap between the target molecular state and all atomic continuum states. This model leads to a component in the scattering rate that is proportional to  $1/\delta_{\text{thresh}}^2$  and is smaller as one detunes farther from the threshold. We note that the first term can also scale nearly as  $1/\delta_{\text{thresh}}^2$ , if, as in our case, the largest terms in the summation are from states near the threshold.

A very similar result holds for the light shift and is written here for completeness

$$\frac{d^2 E^2}{2\hbar^2} \sum_{v' \leq v'_{\text{max}}} \frac{\langle m|v'\rangle \langle v'|m\rangle}{\Delta_{v'}} + \frac{d^2 E^2}{2\hbar^2 \delta_{\text{thresh}}} \left( 1 - \sum_{v' \leq v'_{\text{max}}} \langle m|v'\rangle \langle v'|m\rangle \right) \quad (\text{S.11})$$

### S3. COMPUTATIONAL METHODS FOR BOUND STATES AND SCATTERING

For scattering and near-threshold bound states, we solve the Schrödinger equation by coupled-channel methods [S1], using a basis set in which the electron spins  $s_{\text{Na}}$  and  $s_{\text{Cs}}$  and the nuclear spins  $i_{\text{Na}}$  and  $i_{\text{Cs}}$  are separately coupled,

$$|(s_{\text{Na}} s_{\text{Cs}} S)(i_{\text{Na}} i_{\text{Cs}} I) F M_F\rangle. \quad (\text{S.12})$$

Here  $S$  and  $I$  are the total electron and nuclear spins of the molecule or atomic pair. Their resultant  $F$  is the total spin, with projection  $M_F$  onto the axis of the magnetic field. In principle  $F$  can be further coupled to the rotational angular momentum  $L$ , but in the present work only functions with  $L = 0$  are included. In the absence of a magnetic field,  $F$  and  $M_F$  are both good quantum numbers, but in a magnetic field only  $M_F$  is conserved. The calculations in this paper use basis sets with all possible values of  $S$ ,  $I$  and  $F$  (for the required value of  $M_F$ ) that can be constructed with  $s_{\text{Na}} = s_{\text{Cs}} = 1/2$ ,  $i_{\text{Na}} = 3/2$  and  $i_{\text{Cs}} = 7/2$ .

Scattering calculations are carried out using the MOLSCAT package [S2, S3] and bound-state calculations using the related BOUND package [S3, S4]. The calculations use singlet and triplet interaction potentials for Na+Cs obtained by adjusting the potentials of Docenko *et al.* [S5] at short range, using the same procedure as for KCs [S6], to reproduce the experimental binding energy of the least-bound triplet state [S7] and the position of the s-wave Feshbach resonance near 864 G [S8]. The atomic nuclear hyperfine coupling and Zeeman effects are fully included, using the Hamiltonian described in [S9].

The bound-state calculations were initially used to predict the binding energies of states that lie within about 1 GHz of each atomic threshold of interest. At zero magnetic field, the states fall into groups from each hyperfine state of the atomic pair, labeled by  $(f_{\text{Na}}, f_{\text{Cs}})$ . For example, for the  $(f_{\text{Na}}, f_{\text{Cs}}) = (2, 3)$  states that are of principal interest here, each group comprises 5 zero-field levels with  $F = 1, 2, 3, 4$  and 5; one such group is spread out over a range of energies between 290 MHz and 770 MHz below the  $(f_{\text{Na}}, f_{\text{Cs}}) = (2, 3)$  threshold. In a magnetic field, each zero-field level splits into  $2F + 1$  Zeeman sublevels labeled by  $M_F$ . For each sublevel we can calculate the dependence of the energy on magnetic field, in order to choose a target state for which the Raman transition frequency depends only weakly on field.

The state that we ultimately chose as the target state for Raman transfer is the least-bound  $(F, M_F) = (5, 5)$  sublevel with  $(f_{\text{Na}}, f_{\text{Cs}}) = (2, 3)$ , which was predicted at a binding energy 763 MHz below the  $(f_{\text{Na}}, f_{\text{Cs}}) = (2, 3)$  threshold. As described in the text, it was subsequently located experimentally at a binding energy of 770.57150(9) MHz.

Multichannel bound-state wavefunctions, with components labeled by  $S$ ,  $I$  and  $F$ , are calculated using the method of Thornley and Hutson [S10]. For molecular bound states, they are obtained on a grid of internuclear distances from 2.4 Å to 50 Å. For states of confined atomic pairs, a harmonic confining potential is added to the molecular potentials and the grid is extended to 3000 Å.

#### S4. CALCULATING MATRIX ELEMENTS WITH COUPLED CHANNEL GROUND STATE

To calculate Rabi frequencies for transitions to intermediate vibrational levels of the  $c^3\Sigma^+(\Omega = 1)$  electronic state, we obtain their wavefunctions from a single-channel calculation using the potential curve of [S11]. We then calculate overlap integrals between this function and the triplet ( $S = 1$ ) components of the coupled-channel wavefunctions for either the target molecular state or the confined atomic state. These calculations assume that the intermediate state has pure triplet character and neglect its nuclear spin character, which is unknown.

The full calculation of the Raman Rabi frequency and scattering rate using the methods in section S2 is shown in figure S1. The strong contribution from the atomic threshold arises from the large overlap near-threshold bound states. This leads to a  $1/\delta_{\text{thresh}}^2$  dependence of the scattering rate and favors using deeply bound states as the intermediate state for the Raman transfer. The Raman Rabi frequency is larger than the scattering rate only for the deeply bound states.

Since our coupled-channel calculations include the harmonic confinement, we can calculate the enhancement in  $\Omega_a$  with increasing confinement. Using two calculations with  $\omega_{\text{trap}} = 36.5$  kHz and  $\omega_{\text{trap}} = 80$  kHz trapping frequencies, and assuming a power law scaling, we find  $\Omega_a \propto \omega_{\text{trap}}^{0.58}$ . Thus,  $\Omega_a \propto P^{0.29}$ , which was also experimentally verified. However, we note that in the actual experiment, our system does not have spherically symmetric confinement. The radial trapping frequency is about 5.7 times larger than the axial trapping frequency.

- 
- [S1] J. M. Hutson, *Comp. Phys. Commun.* **84**, 1 (1994).  
[S2] J. M. Hutson and C. R. Le Sueur, *Comp. Phys. Commun.* **241**, 9 (2019).  
[S3] J. M. Hutson and C. R. Le Sueur, MOLSCAT, BOUND and FIELD, version 2020.0, <https://github.com/molscat/molscat> (2020).  
[S4] J. M. Hutson and C. R. Le Sueur, *Comp. Phys. Commun.* **241**, 1 (2019).  
[S5] O. Docenko, M. Tamanis, J. Zaharova, R. Ferber, A. Pashov, H. Knöckel, and E. Tiemann, *J. Phys. B* **39**, S929 (2006).  
[S6] M. Gröbner, P. Weinmann, E. Kirilov, H.-C. Nägerl, P. S. Julienne, C. R. Le Sueur, and J. M. Hutson, *Phys. Rev. A* **95**, 022715 (2017).




FIG. S1: Full calculation of the Raman Rabi frequency and scattering rate as a function of the single-photon frequency including all vibrational states of  $c^3\Sigma^+(\Omega = 1)$  and the atomic continuum. The second frequency in the Raman transition is offset by about 770 MHz and resonant with the two photon transition. The assumed excited-state linewidth for all molecular lines is 20 MHz.

- [S7] J. D. Hood, Y. Yu, Y.-W. Lin, J. T. Zhang, K. Wang, L. R. Liu, B. Gao, and K.-K. Ni, *Phys. Rev. Research* **2**, 023108 (2020).
- [S8] J. T. Zhang, Y. Yu, W. B. Cairncross, K. Wang, L. R. B. Picard, J. D. Hood, Y.-W. Lin, J. M. Hutson, and K.-K. Ni, *Phys. Rev. Lett.* **124**, 253401 (2020).
- [S9] J. M. Hutson, E. Tiesinga, and P. S. Julienne, *Phys. Rev. A* **78**, 052703 (2008).
- [S10] A. E. Thornley and J. M. Hutson, *J. Chem. Phys.* **101**, 5578 (1994), <https://doi.org/10.1063/1.467345>.
- [S11] A. Grochola, P. Kowalczyk, J. Szczepkowski, W. Jastrzebski, A. Wakim, P. Zabawa, and N. P. Bigelow, *Phys. Rev. A* **84**, 012507 (2011).