

## 1.

$$\begin{aligned}
 |1, 1; 2, 2\rangle &= |1, 1\rangle_1 |1, 1\rangle_2 \\
 |1, 1; 2, -2\rangle &= |1, -1\rangle_1 |1, -1\rangle_2 \\
 |1, 1; 2, 1\rangle &= \frac{J_-}{2\hbar} |1, 1; 2, 2\rangle \\
 &= \frac{L_{1-} + L_{2-}}{2\hbar} |1, 1\rangle_1 |1, 1\rangle_2 \\
 &= \frac{|1, 0\rangle_1 |1, 1\rangle_2 + |1, 1\rangle_1 |1, 0\rangle_2}{\sqrt{2}} \\
 |1, 1; 2, -1\rangle &= \frac{J_+}{2\hbar} |1, 1; 2, 2\rangle \\
 &= \frac{L_{1+} + L_{2+}}{2\hbar} |1, -1\rangle_1 |1, -1\rangle_2 \\
 &= \frac{|1, 0\rangle_1 |1, -1\rangle_2 + |1, -1\rangle_1 |1, 0\rangle_2}{\sqrt{2}} \\
 |1, 1; 2, 0\rangle &= \frac{J_-}{\sqrt{6}\hbar} |1, 1; 2, 1\rangle \\
 &= \frac{L_{1-} + L_{2-}}{\sqrt{6}\hbar} \frac{|1, 0\rangle_1 |1, 1\rangle_2 + |1, 1\rangle_1 |1, 0\rangle_2}{\sqrt{2}} \\
 &= \frac{|1, -1\rangle_1 |1, 1\rangle_2 + 2|1, 0\rangle_1 |1, 0\rangle_2 + |1, 1\rangle_1 |1, -1\rangle_2}{\sqrt{6}}
 \end{aligned}$$

From orthogonality

$$\begin{aligned}
 |1, 1; 1, 1\rangle &= \frac{|1, 0\rangle_1 |1, 1\rangle_2 - |1, 1\rangle_1 |1, 0\rangle_2}{\sqrt{2}} \\
 |1, 1; 1, -1\rangle &= \frac{|1, 0\rangle_1 |1, -1\rangle_2 - |1, -1\rangle_1 |1, 0\rangle_2}{\sqrt{2}} \\
 |1, 1; 1, 0\rangle &= \frac{J_-}{\sqrt{2}\hbar} |1, 1; 1, 1\rangle \\
 &= \frac{L_{1-} + L_{2-}}{\sqrt{2}\hbar} \frac{|1, 0\rangle_1 |1, 1\rangle_2 - |1, 1\rangle_1 |1, 0\rangle_2}{\sqrt{2}} \\
 &= \frac{|1, -1\rangle_1 |1, 1\rangle_2 - |1, 1\rangle_1 |1, -1\rangle_2}{\sqrt{2}}
 \end{aligned}$$

From orthogonality

$$|1, 1; 0, 0\rangle = \frac{|1, -1\rangle_1 |1, 1\rangle_2 - |1, 0\rangle_1 |1, 0\rangle_2 + |1, 1\rangle_1 |1, -1\rangle_2}{\sqrt{3}}$$

## 2.

### (a)

Since  $J_z = L_z + S_z$  and  $[L_z, S_z] = 0$ , the  $J_z$  eigenstates are sum of  $L_z$  and  $S_z$  eigenstates that has the same sum of  $m_l$  and  $m_s$ . (The only variable on the RHS are  $m_l$  and  $m_s$  among which  $m_s$  can only be  $\pm \frac{1}{2}$  so the constraint above limit the decomposition to the given form.)

(b)

Hamiltonian

$$\begin{aligned} H &= \frac{L^2}{2ma^2} + V_0 + \frac{e^2 \vec{L} \cdot \vec{S}}{2mc^2 a^3} \\ &= \frac{L^2}{2ma^2} + V_0 + \frac{e^2}{4mc^2 a^3} (J^2 - L^2 - S^2) \\ &= \frac{l(l+1)}{2ma^2} + V_0 + \frac{e^2}{4mc^2 a^3} \left( j(j+1) - l(l+1) - \frac{3}{4} \right) \end{aligned}$$

When  $l = 0$  there's only one manifold instead of two.

**3.**

$$\begin{aligned} e^{\lambda A} B e^{-\lambda A} &= \left( \sum_{m=0}^{\infty} \frac{\lambda^m A^m}{m!} \right) B \left( \sum_{n=0}^{\infty} \frac{\lambda^n (-A)^n}{n!} \right) \\ &= \sum_{m,n=0}^{\infty} \frac{\lambda^m A^m}{m!} B \frac{\lambda^n (-A)^n}{n!} \\ &= \sum_{m=0}^{\infty} \sum_{n=0}^m \frac{\lambda^m A^{m-n} B (-A)^n}{(m-n)! n!} \\ &= \sum_{m=0}^{\infty} \frac{\lambda^m}{m!} \sum_{n=0}^m \frac{m! A^{m-n} B (-A)^n}{(m-n)! n!} \end{aligned}$$

So now we just need to show that

$$[A^{(m)}, B] = \sum_{n=0}^m \frac{m! A^{m-n} B (-A)^n}{(m-n)! n!}$$

where  $[A^{(0)}, B] \equiv B$ ,  $[A^{(n)}, B] \equiv [A, [A^{(n-1)}, B]]$ . For  $m = 0$ , this is trivially true. If it is true for  $m - 1$ ,

$$\begin{aligned} [A^{(m)}, B] &= \left[ A, \sum_{n=0}^{m-1} \frac{(m-1)! A^{m-n-1} B (-A)^n}{(m-n-1)! n!} \right] \\ &= \sum_{n=0}^{m-1} \frac{(m-1)! A^{m-n} B (-A)^n}{(m-n-1)! n!} + \sum_{n=1}^m \frac{(m-1)! A^{m-n} B (-A)^n}{(m-n)!(n-1)!} \end{aligned}$$

Extend the summation taking advantage of  $(-1)! = \infty$

$$\begin{aligned} [A^{(m)}, B] &= \sum_{n=0}^m \frac{(m-1)! A^{m-n} B (-A)^n}{(m-n-1)! n!} + \frac{(m-1)! A^{m-n} B (-A)^n}{(m-n)!(n-1)!} \\ &= \sum_{n=0}^m \frac{(m)! A^{m-n} B (-A)^n}{(m-n)! n!} \left( \frac{m-n}{m} + \frac{n}{m} \right) \\ &= \sum_{n=0}^m \frac{(m)! A^{m-n} B (-A)^n}{(m-n)! n!} \end{aligned}$$

Therefore, the equation is true for all  $m \geq 0$ .

If  $[A, B] = \gamma B$ ,  $[A^{(n)}, B] = \gamma^n B$

$$\begin{aligned} e^{\lambda A} B e^{-\lambda A} &= \sum_{m=0}^{\infty} \frac{\lambda^m}{m!} [A^{(m)}, B] \\ &= \sum_{m=0}^{\infty} \frac{\lambda^m \gamma^m}{m!} B \\ &= e^{\lambda \gamma} B \end{aligned}$$

#### 4.

Since  $[\lambda, A] = 0$  and  $[A, A] = 0$  (i.e. everything commutes),

$$\frac{d e^{\lambda A}}{d\lambda} = A e^{\lambda A}$$

We have

$$\begin{aligned} \frac{dG}{d\lambda} &= \frac{d e^{\lambda A} e^{\lambda B}}{d\lambda} \\ &= e^{\lambda A} \frac{d e^{\lambda B}}{d\lambda} + \frac{d e^{\lambda A}}{d\lambda} e^{\lambda B} \\ &= e^{\lambda A} B e^{\lambda B} + A e^{\lambda A} e^{\lambda B} \\ &= e^{\lambda A} B e^{-\lambda A} e^{\lambda A} e^{\lambda B} + A e^{\lambda A} e^{\lambda B} \\ &= (A + e^{\lambda A} B e^{-\lambda A}) G \\ &= \left( A + \sum_{m=0}^{\infty} \frac{\lambda^m}{m!} [A^{(m)}, B] \right) G \end{aligned}$$

Using this,

$$\begin{aligned} \frac{dG}{d\lambda} &= \left( \frac{d e^{\lambda B^\dagger} e^{\lambda A^\dagger}}{d\lambda} \right)^\dagger \\ &= \left( \left( B^\dagger + \sum_{m=0}^{\infty} \frac{\lambda^m}{m!} [B^{\dagger(m)}, A^\dagger] \right) G^\dagger \right)^\dagger \\ &= G \left( B + \sum_{m=0}^{\infty} \frac{\lambda^m}{m!} [A, B^{(m)}] \right) \end{aligned}$$

If  $C \equiv [A, B]$  commutes with both  $A$  and  $B$  (and therefore  $A + B$ ),

$$\begin{aligned} \frac{dG}{d\lambda} &= (A + B + \lambda C) G \\ G &= e^{A+B+\lambda^2 C/2} \\ e^A e^B &= e^{A+B+C/2} \end{aligned}$$

## 5.

### (a)

Eigenvalue  $\lambda$

$$\begin{aligned} 0 &= (h - \lambda)^2 - |g|^2 \\ h - \lambda &= \pm |g| \\ \lambda &= h \pm |g| \end{aligned}$$

Corresponding eigen vectors are  $\frac{1}{\sqrt{2}} \begin{pmatrix} 1, \pm \frac{g}{|g|} \end{pmatrix}$

### (b)

Initial state

$$|\psi_0\rangle = \frac{1}{\sqrt{2}}(|+\rangle + |-\rangle)$$

At time  $t$ ,

$$\begin{aligned} |\psi_t\rangle &= \frac{1}{\sqrt{2}} \left( \exp\left(-i\frac{h+|g|}{\hbar}t\right)|+\rangle + \exp\left(-i\frac{h-|g|}{\hbar}t\right)|-\rangle \right) \\ &= \frac{e^{-iht/\hbar}}{\sqrt{2}} \left( e^{i|g|t/\hbar}|+\rangle + e^{-i|g|t/\hbar}|-\rangle \right) \\ &= \frac{e^{-iht/\hbar}}{2} \left( e^{-i|g|t/\hbar} \left( |1\rangle + \frac{g}{|g|}|2\rangle \right) + e^{i|g|t/\hbar} \left( |1\rangle - \frac{g}{|g|}|2\rangle \right) \right) \\ &= e^{-iht/\hbar} \left( \cos\left(\frac{|g|t}{\hbar}\right)|1\rangle - i\frac{g}{|g|}\sin\left(\frac{|g|t}{\hbar}\right)|2\rangle \right) \end{aligned}$$

## 6.

### (a)

In Heisenberg picture,

$$\begin{aligned} H &= \hbar\omega_0 a^\dagger a - Fz_0(a^\dagger + a) \\ i\hbar \frac{\partial}{\partial t} a &= -[H, a] \\ &= \hbar\omega_0 a - Fz_0 \\ \frac{\partial}{\partial t} a &= -i\omega_0 a + i\frac{F_0 z_0}{\hbar} e^{-t/\tau} \\ a &= e^{-i\omega_0 t} a_0 + i\tau \frac{F_0 z_0}{\hbar} \left( e^{-i\omega_0 t} - e^{-t/\tau} \right) \end{aligned}$$

$$\begin{aligned}
 \langle n|\psi\rangle &= \langle n_t|0\rangle \\
 &= \frac{1}{\sqrt{n!}} \langle 0_t|a^n|0\rangle \\
 &= \frac{1}{\sqrt{n!}} \langle 0_t|\left(e^{-i\omega_0 t}a_0 + i\tau\frac{F_0 z_0}{\hbar}\left(e^{-i\omega_0 t} - e^{-t/\tau}\right)\right)^n|0\rangle \\
 &= \frac{1}{\sqrt{n!}} \left(i\tau\frac{F_0 z_0}{\hbar}\left(e^{-i\omega_0 t} - e^{-t/\tau}\right)\right)^n \langle 0_t|0\rangle
 \end{aligned}$$

From normalization,

$$\begin{aligned}
 1 &= \sum_n \frac{1}{n!} \left(\tau\frac{F_0 z_0}{\hbar}\left|e^{-i\omega_0 t} - e^{-t/\tau}\right|\right)^{2n} |\langle 0_t|0\rangle|^2 \\
 |\langle 0_t|0\rangle|^2 &= \exp\left(-\frac{F_0^2 z_0^2 \tau^2}{\hbar^2}\left|e^{-i\omega_0 t} - e^{-t/\tau}\right|^2\right) \\
 |\langle n|\psi\rangle|^2 &= \frac{1}{n!} \left|\frac{F_0^2 z_0^2 \tau^2}{\hbar^2}\left|e^{-i\omega_0 t} - e^{-t/\tau}\right|^2\right|^n \exp\left(-\frac{F_0^2 z_0^2 \tau^2}{\hbar^2}\left|e^{-i\omega_0 t} - e^{-t/\tau}\right|^2\right)
 \end{aligned}$$

(b)