

1.

2.

(a)

Since $J_z = L_z + S_z$ and $[L_z, S_z] = 0$, the J_z eigenstates are sum of L_z and S_z eigenstates that has the same sum of m_l and m_s . (The only variable on the RHS are m_l and m_s among which m_s can only be $\pm \frac{1}{2}$ so the constraint above limit the decomposition to the given form.)

(b)

3.

4.

5.

(a)

Eigenvalue λ

$$\begin{aligned} 0 &= (h - \lambda)^2 - |g|^2 \\ h - \lambda &= \pm |g| \\ \lambda &= h \pm |g| \end{aligned}$$

Corresponding eigen vectors are $\frac{1}{\sqrt{2}} \left(1, \pm \frac{g}{|g|} \right)$

(b)

Initial state

$$|\psi_0\rangle = \frac{1}{\sqrt{2}}(|+\rangle + |-\rangle)$$

At time t ,

$$\begin{aligned} |\psi_t\rangle &= \frac{1}{\sqrt{2}} \left(\exp\left(-i\frac{h+|g|}{\hbar}t\right)|+\rangle + \exp\left(-i\frac{h-|g|}{\hbar}t\right)|-\rangle \right) \\ &= \frac{e^{-iht/\hbar}}{\sqrt{2}} \left(e^{-i|g|t/\hbar}|+\rangle + e^{i|g|t/\hbar}|-\rangle \right) \\ &= \frac{e^{-iht/\hbar}}{2} \left(e^{-i|g|t/\hbar} \left(|1\rangle + \frac{g}{|g|}|2\rangle \right) + e^{i|g|t/\hbar} \left(|1\rangle - \frac{g}{|g|}|2\rangle \right) \right) \\ &= e^{-iht/\hbar} \left(\cos\left(\frac{|g|t}{\hbar}\right)|1\rangle - i\frac{g}{|g|}\sin\left(\frac{|g|t}{\hbar}\right)|2\rangle \right) \end{aligned}$$

6.

(a)

(b)