

# HOMOTOPY PRO-NILPOTENT STRUCTURED RING SPECTRA AND TOPOLOGICAL QUILLEN LOCALIZATION

YU ZHANG

**ABSTRACT.** The aim of this short paper is to show that the homotopy limit of any diagram of nilpotent structured ring spectra is **TQ**-local, where structured ring spectra are described as algebras over a spectral operad  $\mathcal{O}$ ; in particular, every homotopy pro-nilpotent structured ring spectrum is **TQ**-local. Here, **TQ** is short for topological Quillen homology, which is weakly equivalent to  $\mathcal{O}$ -algebra stabilization. As an application, we simultaneously extend the previously known connected and nilpotent **TQ**-Whitehead theorems to a homotopy pro-nilpotent **TQ**-Whitehead theorem. We also compare **TQ**-localization with **TQ**-completion and show that **TQ**-local  $\mathcal{O}$ -algebras that are **TQ**-good are **TQ**-complete. Finally, we show that every  $(-1)$ -connected  $\mathcal{O}$ -algebra with a principally refined Postnikov tower is **TQ**-local, provided that  $\mathcal{O}$  is  $(-1)$ -connected.

## 1. INTRODUCTION

Structured ring spectra are spectra with extra algebraic structure encoded by the action of an operad  $\mathcal{O}$  in the closed symmetric monoidal category of modules  $(\mathbf{Mod}_{\mathcal{R}}, \wedge, \mathcal{R})$  over a commutative ring spectrum  $\mathcal{R}$ . For a fixed operad  $\mathcal{O}$ , denote by  $\mathbf{Alg}_{\mathcal{O}}$  the category of  $\mathcal{O}$ -algebras. In this paper we are working with reduced operads  $\mathcal{O}$  (i.e., such that  $\mathcal{O}[0] = *$ , where  $*$  denotes the trivial  $\mathcal{R}$ -module); such  $\mathcal{O}$ -algebras are non-unital. Similar to the ordinary homology of spaces, a precisely analogous notion of homology for  $\mathcal{O}$ -algebras is topological Quillen (**TQ**) homology [4, 6, 11, 21, 27, 37, 35]; useful introductions to notions of homology for algebraic structures appear, for instance, in [20, 24, 41, 43, 44]. It turns out that **TQ**-homology is weakly equivalent to stabilization  $\Omega^{\infty}\Sigma^{\infty}$  in  $\mathcal{O}$ -algebras [5, 28, 42, 45, 46] under appropriate connectivity conditions—this is because **TQ** is 1-excisive and agrees to order 1 with the identity functor on  $\mathcal{O}$ -algebras (via the **TQ**-Hurewicz map  $\mathrm{id} \rightarrow \mathbf{TQ}$ ) in the sense of Goodwillie [25, 1.2]; see, for instance, [11, 28, 36].

The **TQ**-local homotopy theory for  $\mathcal{O}$ -algebras is established in [29], where the upshot is that: if  $X$  is a cofibrant  $\mathcal{O}$ -algebra, then its weak **TQ**-fibrant replacement  $X \rightarrow L_{\mathbf{TQ}}(X)$  is the **TQ**-localization of  $X$ . By construction, the comparison map  $X \rightarrow L_{\mathbf{TQ}}(X)$  is a cofibration that is also a **TQ**-equivalence such that  $L_{\mathbf{TQ}}(X)$  is **TQ**-local (Definition 2.2). Intuitively, the **TQ**-localization  $L_{\mathbf{TQ}}(X)$  can be thought of as “the part of  $X$  that **TQ**-homology sees”.

In this short paper we attack the following question: When is the comparison map  $X \rightarrow L_{\mathbf{TQ}}(X)$  a weak equivalence? In other words, when is an  $\mathcal{O}$ -algebra  $X$  already **TQ**-local? For instance, we know from [13] that every connected  $\mathcal{O}$ -algebra is **TQ**-complete and hence  $X \simeq L_{\mathbf{TQ}}(X)$ , but we also know from [28] that every connected  $\mathcal{O}$ -algebra is the homotopy limit of a tower of nilpotent  $\mathcal{O}$ -algebras and hence  $X$  is homotopy pro-nilpotent (Definition 1.2); here,  $\mathcal{R}, \mathcal{O}$  were assumed to be  $(-1)$ -connected.

This leads us to one of the motivations of our work: what amounts to the “first half” of a conjecture of Francis-Gaitsgory [18, 3.4.5] that (i) the natural map comparing  $X$  with its  $\mathbf{TQ}$ -completion should be a weak equivalence for every homotopy pro-nilpotent  $\mathcal{O}$ -algebra  $X$ . Our main result, Theorem 1.8, is that (i) is true in general, provided that in the comparison map we replace “ $\mathbf{TQ}$ -completion” with “ $\mathbf{TQ}$ -localization”. Our strategy of attack is to leverage the  $\mathbf{TQ}$ -local homotopy theory of  $\mathcal{O}$ -algebras in [29] with the fact, proved in [12], that  $M$ -nilpotent  $\mathcal{O}$ -algebras are  $\mathbf{TQ}|_{\mathrm{Nil}_M}$ -complete.

**Definition 1.1.** Let  $X$  be an  $\mathcal{O}$ -algebra and  $M \geq 2$ . We say that  $X$  is  *$M$ -nilpotent* if all the  $M$ -ary and higher operations  $\mathcal{O}[t] \wedge X^{\wedge t} \rightarrow X$  of  $X$  are trivial (i.e., if these maps factor through the trivial  $\mathcal{R}$ -module  $*$  for each  $t \geq M$ ).

**Definition 1.2.** An  $\mathcal{O}$ -algebra is *nilpotent* (resp. *homotopy pro-nilpotent*) if it is  $M$ -nilpotent for some  $M \geq 2$  (resp. if it is weakly equivalent to the homotopy limit of a tower of nilpotent  $\mathcal{O}$ -algebras).

It is worth pointing out that homotopy pro-nilpotent  $\mathcal{O}$ -algebras need not be nilpotent; the following describes a large class of such  $\mathcal{O}$ -algebras.

**Proposition 1.3.** *If  $X$  is a connected  $\mathcal{O}$ -algebra and  $\mathcal{O}, \mathcal{R}$  are  $(-1)$ -connected, then  $X$  is homotopy pro-nilpotent.*

*Proof.* This is proved in Harper-Hess [28, 1.12] by showing that the homotopy completion tower of  $X$  converges strongly to  $X$ .  $\square$

**Proposition 1.4** ( $\mathbf{TQ}$ -local Whitehead theorem). *A map  $X \rightarrow Y$  between  $\mathbf{TQ}$ -local  $\mathcal{O}$ -algebras is a weak equivalence if and only if it is a  $\mathbf{TQ}$ -homology equivalence.*

*Proof.* This follows from the definition of  $\mathbf{TQ}$ -local  $\mathcal{O}$ -algebras (Definition 2.2); see, for instance, Hirschhorn [32, 3.2.13].  $\square$

**Proposition 1.5** (Preservation of the  $\mathbf{TQ}$ -local property: Homotopy limits). *The homotopy limit of a small diagram of  $\mathbf{TQ}$ -local  $\mathcal{O}$ -algebras is  $\mathbf{TQ}$ -local.*

*Proof.* This follows from the definition of  $\mathbf{TQ}$ -local  $\mathcal{O}$ -algebras; see, for instance, Dror Farjoun [14, 1.A.8, 1.G] and Hirschhorn [32, 19.4.4]. Here is another, essentially equivalent, proof: It follows from Proposition 1.4 that the homotopy limit in  $\mathbf{Alg}_{\mathcal{O}}$  of a small diagram of  $\mathbf{TQ}$ -local  $\mathcal{O}$ -algebras is weakly equivalent to its homotopy limit calculated in the  $\mathbf{TQ}$ -local homotopy theory [29, 5.14]; hence, verifying that the homotopy limit in  $\mathbf{Alg}_{\mathcal{O}}$  is  $\mathbf{TQ}$ -local reduces to the usual fibrancy property of homotopy limits in a homotopy theory (in this case, in the  $\mathbf{TQ}$ -local homotopy theory); see, for instance, Hirschhorn [32, 18.5.2], together with Ching-Harper [13, 8.9] for a discussion of homotopy limits in the context of  $\mathcal{O}$ -algebras.  $\square$

For instance, consider any pullback diagram of the form

$$\begin{array}{ccc} A & \longrightarrow & B \\ \downarrow & & \downarrow p \\ C & \longrightarrow & D \end{array}$$

in  $\mathbf{Alg}_{\mathcal{O}}$ . It follows from Proposition 1.5 that if  $B, C, D$  are  $\mathbf{TQ}$ -local and  $p$  is a fibration, then  $A$  is  $\mathbf{TQ}$ -local. Taking  $C = *$ , for instance, shows that  $\mathbf{TQ}$ -local  $\mathcal{O}$ -algebras play nicely with fibration sequences; this is not expected to be true, in

general, if we replace “TQ-local” with “TQ-complete” and is one of the reasons why TQ-localization is often better behaved than TQ-completion (at the expense of a much larger construction).

**Proposition 1.6.** *If  $X$  is an  $M$ -nilpotent  $\mathcal{O}$ -algebra (resp.  $Z$  is an  $\mathcal{O}$ -algebra) for some  $M \geq 2$ , then its  $\mathrm{TQ}|_{\mathrm{Nil}_M}$ -completion  $X_{\mathrm{TQ}|_{\mathrm{Nil}_M}}^\wedge$  (resp. TQ-completion  $Z_{\mathrm{TQ}}^\wedge$ ) is TQ-local.*

*Proof.* By Proposition 1.5, it suffices to verify that  $X_{\mathrm{TQ}|_{\mathrm{Nil}_M}}^\wedge$  (resp.  $Z_{\mathrm{TQ}}^\wedge$ ) is the homotopy limit of a small diagram of TQ-local  $\mathcal{O}$ -algebras—we defer the proof of this to Section 2 (see Propositions 2.10 and 2.9, respectively).  $\square$

**Proposition 1.7.** *Let  $M \geq 2$ .*

- (a) *If  $X$  is an  $M$ -nilpotent  $\mathcal{O}$ -algebra, then the natural map  $X \simeq X_{\mathrm{TQ}|_{\mathrm{Nil}_M}}^\wedge$  is a weak equivalence.*
- (b) *If  $Z$  is a connected  $\mathcal{O}$ -algebra and  $\mathcal{O}, \mathcal{R}$  are  $(-1)$ -connected, then the natural map  $Z \simeq Z_{\mathrm{TQ}}^\wedge$  is a weak equivalence.*

*Proof.* Part (a) is proved in Ching-Harper [12, 2.12] and part (b) is proved in Ching-Harper [13, 1.2].  $\square$

The following theorem is the main result of this paper. Our proof is embarrassingly simple—one of the significant advantages of having available the TQ-local homotopy theory framework to work within and organize our arguments in—it follows immediately from the above propositions, together with verifying that  $\mathrm{TQ}|_{\mathrm{Nil}_M}$ -resolutions have TQ-local fibrant replacements in  $\mathrm{Alg}_{\mathcal{O}}$ ; we defer the proof of this to Section 2 (see Proposition 2.10).

**Theorem 1.8** (Homotopy pro-nilpotent TQ-localization theorem). *Let  $X$  be a fibrant  $\mathcal{O}$ -algebra.*

- (a) *If  $X$  is nilpotent, then  $X$  is TQ-local.*
- (b) *If  $X$  is homotopy pro-nilpotent, then  $X$  is TQ-local.*
- (c) *If  $X$  is the homotopy limit of any small diagram of nilpotent  $\mathcal{O}$ -algebras, then  $X$  is TQ-local.*
- (d) *If  $X$  is connected and  $\mathcal{O}, \mathcal{R}$  are  $(-1)$ -connected, then  $X$  is TQ-local.*

*Remark 1.9.* It is worth pointing out (Proposition 2.7) that if some fibrant replacement of an  $\mathcal{O}$ -algebra  $X$  is TQ-local, then every fibrant replacement of  $X$  is TQ-local.

*Proof of Theorem 1.8.* Part (a) follows from Propositions 1.6 and 1.7. Parts (b) and (c) follow from part (a), together with Proposition 1.5. Part (d) follows from part (c), together with Proposition 1.3; alternately, it follows from Propositions 1.6 and 1.7.  $\square$

As an application of the main result, we obtain the following homotopy pro-nilpotent TQ-Whitehead theorem that simultaneously extends the previously known connected and nilpotent TQ-Whitehead theorems.

**Theorem 1.10** (Homotopy pro-nilpotent TQ-Whitehead theorem). *A map  $X \rightarrow Y$  between homotopy pro-nilpotent  $\mathcal{O}$ -algebras is a weak equivalence if and only if it is a TQ-homology equivalence; more generally, this remains true if  $X, Y$  are homotopy limits of small diagrams of nilpotent  $\mathcal{O}$ -algebras.*

*Proof.* This follows from Theorem 1.8, together with Proposition 1.4.  $\square$

We also compare TQ-localization with TQ-completion and show that TQ-local  $\mathcal{O}$ -algebras that are TQ-good are TQ-complete (Theorem 3.1). Finally we show that  $\mathcal{O}$ -algebras with a principally refined Postnikov tower are TQ-local, provided that mild connectivity assumptions are satisfied; here is the theorem, whose proof we defer to Section 4.

**Theorem 1.11.** *Let  $X$  be a fibrant  $\mathcal{O}$ -algebra. If  $X, \mathcal{O}, \mathcal{R}$  are  $(-1)$ -connected and  $X$  has a principally refined Postnikov tower, then  $X$  is TQ-local.*

**1.12. Conventions and notations.** We work in the category  $\mathbf{Alg}_{\mathcal{O}}$  of algebras over an operad  $\mathcal{O}$  in  $\mathbf{Mod}_{\mathcal{R}}$ , the category of  $\mathcal{R}$ -modules, where  $\mathcal{R}$  is a commutative monoid in the category of symmetric spectra [33, 47]. Throughout this paper, we assume that: (i)  $\mathcal{O}$  is reduced, meaning that  $\mathcal{O}[0] = *$  (i.e.,  $\mathcal{O}$ -algebras are non-unital) and (ii) the natural maps  $\mathcal{R} \rightarrow \mathcal{O}[1]$  and  $* \rightarrow \mathcal{O}[n]$  are flat stable cofibrations [28] in  $\mathcal{R}$ -modules for each  $n \geq 0$ ; see, for instance, [13, 2.1, 6.12]. Unless otherwise specified, we work with the positive flat stable model structure on  $\mathbf{Alg}_{\mathcal{O}}$  [28, 47]. To keep this paper appropriately concise, we freely use notation from [13, 28, 29]. Finally, “connected” is short for “0-connected”.

**Acknowledgments.** The author would like to thank John E. Harper and Niko Schonsheck for enlightening discussions and helpful suggestions, and Michael Ching and Mark W. Johnson for helpful conversations. The author is grateful to Oscar Randal-Williams for detailed and helpful critical comments on an early draft of this paper. The author was supported in part by the Simons Foundation: Collaboration Grants for Mathematicians #638247.

## 2. TQ-LOCAL $\mathcal{O}$ -ALGEBRAS AND $\mathrm{TQ}|_{\mathrm{Nil}_M}$ -RESOLUTIONS

The purpose of this section is to recall the definition of TQ-local  $\mathcal{O}$ -algebras, discuss their basic properties, and finally to verify that TQ-resolutions of  $\mathcal{O}$ -algebras and  $\mathrm{TQ}|_{\mathrm{Nil}_M}$ -resolutions of  $M$ -nilpotent  $\mathcal{O}$ -algebras have TQ-local fibrant replacements in  $\mathbf{Alg}_{\mathcal{O}}$  (Propositions 2.9 and 2.10).

TQ-homology and its relative form, topological Andre-Quillen (TAQ) homology, first introduced in [4] for commutative ring spectra (see also [3, 5, 6, 21, 35, 38, 39, 45]), are defined as derived indecomposables of  $\mathcal{O}$ -algebras analogous to Quillen homology of commutative algebras [1, 44]; see also [19, 24, 20, 41]. More precisely, factoring the canonical truncation map  $\mathcal{O} \rightarrow \tau_1 \mathcal{O}$  (see [28]) in the category of operads as  $\mathcal{O} \rightarrow J \rightarrow \tau_1 \mathcal{O}$ , a cofibration followed by a weak equivalence, we get the corresponding change of operads adjunction

$$(1) \quad \mathbf{Alg}_{\mathcal{O}} \begin{matrix} \xrightarrow{Q} \\ \xleftarrow{U} \end{matrix} \mathbf{Alg}_J$$

with left adjoint on top, where  $Q(X) = J \circ_{\mathcal{O}}(X)$  and  $U$  denotes the forgetful functor (or less concisely, the “forget along the map  $\mathcal{O} \rightarrow J$  functor”).

**Definition 2.1.** Let  $X$  be an  $\mathcal{O}$ -algebra. The *topological Quillen homology* (or TQ-homology, for short) of  $X$  is

$$\mathrm{TQ}(X) := RU(\mathrm{LQ}(X))$$

the  $\mathcal{O}$ -algebra defined via the indicated composite of total right and left derived functors. If  $X$  is cofibrant, then  $\mathrm{TQ}(X) \simeq UQ(X)$  and the unit of the  $(Q, U)$  adjunction in (1) is the  $\mathrm{TQ}$ -Hurewicz map  $X \rightarrow UQX$  of the form  $X \rightarrow \mathrm{TQ}(X)$ .

$\mathrm{TQ}$ -homology has been shown to enjoy several properties analogous to the ordinary homology of spaces; see, for instance, [4, 5, 11, 13, 28, 36]. Furthermore, it turns out that  $\mathrm{TQ}$ -homology is weakly equivalent to stabilization in the category of  $\mathcal{O}$ -algebras, provided that  $\mathcal{O}, \mathcal{R}$  are  $(-1)$ -connected; see, for instance, [5, 36, 42, 45, 46]; a simple proof using Goodwillie's functor calculus [25] is given in [28, 1.14].

Localization has been proved to be a useful method in algebraic topology. The idea is to focus on the information one cares about and ignore everything else, hence simplifying problems accordingly. See [2, 9, 10, 16, 31, 48] for a discussion of localization methods. Also see, for useful developments and ideas, [8, 14, 16, 32, 34, 40].

In this paper we consider localization of  $\mathcal{O}$ -algebras with respect to  $\mathrm{TQ}$ -homology. Recall the following from [29].

**Definition 2.2.** Let  $f: A \rightarrow B$  be a map in  $\mathrm{Alg}_{\mathcal{O}}$ . We say that  $f$  is a  $\mathrm{TQ}$ -equivalence if  $f$  induces a weak equivalence  $\mathrm{TQ}(A) \simeq \mathrm{TQ}(B)$  on  $\mathrm{TQ}$ -homology. We say that  $f$  is a  $\mathrm{TQ}$ -acyclic strong cofibration if  $f$  is a cofibration between cofibrant objects which is also a  $\mathrm{TQ}$ -equivalence. An  $\mathcal{O}$ -algebra  $X$  is called  $\mathrm{TQ}$ -local if (i)  $X$  is fibrant in  $\mathrm{Alg}_{\mathcal{O}}$  and (ii) every  $\mathrm{TQ}$ -acyclic strong cofibration  $A \rightarrow B$  induces a weak equivalence

$$\mathrm{Hom}(A, X) \xleftarrow{\simeq} \mathrm{Hom}(B, X)$$

on mapping spaces in  $\mathbf{sSet}$ ; here we are using the simplicial model structure on  $\mathrm{Alg}_{\mathcal{O}}$  (see, for instance, [13, 17, 22, 23, 28]).

It is useful to note that the collection of  $\mathrm{TQ}$ -local  $\mathcal{O}$ -algebras is exactly the collection of weak  $\mathrm{TQ}$ -fibrant objects in the  $\mathrm{TQ}$ -local homotopy theory [29]. Intuitively, a  $\mathrm{TQ}$ -local object in  $\mathrm{Alg}_{\mathcal{O}}$  is an  $\mathcal{O}$ -algebra  $X$  which only sees  $\mathrm{TQ}$ -homology information. The following proposition will be useful for detecting  $\mathrm{TQ}$ -local  $\mathcal{O}$ -algebras.

**Proposition 2.3.** *Let  $X$  be a fibrant  $\mathcal{O}$ -algebra. Then  $X$  is  $\mathrm{TQ}$ -local if and only if  $X \rightarrow *$  satisfies the right lifting property with respect to every  $\mathrm{TQ}$ -acyclic strong cofibration  $A \rightarrow B$  in  $\mathrm{Alg}_{\mathcal{O}}$ .*

*Proof.* This is proved in [29, 3.12]. □

**Proposition 2.4.** *Let  $Y$  be a fibrant object in  $\mathrm{Alg}_J$ . Then  $UY \in \mathrm{Alg}_{\mathcal{O}}$  is  $\mathrm{TQ}$ -local.*

*Proof.* This follows from Proposition 2.3 by using the  $(Q, U)$  adjunction (1). □

Next we observe that the  $\mathrm{TQ}$ -local property is preserved by weak equivalences between fibrant  $\mathcal{O}$ -algebras.

**Proposition 2.5.** *Let  $X \rightarrow Y$  be a weak equivalence between fibrant objects in  $\mathrm{Alg}_{\mathcal{O}}$ . Then  $X$  is  $\mathrm{TQ}$ -local if and only if  $Y$  is  $\mathrm{TQ}$ -local.*

*Proof.* Let  $A \rightarrow B$  be a TQ-acyclic strong cofibration. Consider the commutative diagram of mapping spaces of the form

$$\begin{array}{ccc} \mathbf{Hom}(A, X) & \longleftarrow & \mathbf{Hom}(B, X) \\ \downarrow \sim & & \downarrow \sim \\ \mathbf{Hom}(A, Y) & \longleftarrow & \mathbf{Hom}(B, Y) \end{array}$$

in  $\mathbf{sSet}$ . Since the vertical maps are weak equivalences, it follows that the top map is a weak equivalence if and only if the bottom map is a weak equivalence.  $\square$

This observation generalizes as follows.

**Proposition 2.6.** *Consider any weak equivalence  $X \rightarrow Y$  in  $\mathbf{Alg}_{\mathcal{O}}$ . Let  $X', Y'$  be fibrant replacements of  $X, Y$ , respectively, in  $\mathbf{Alg}_{\mathcal{O}}$ . Then  $X'$  is TQ-local if and only if  $Y'$  is TQ-local.*

*Proof.* By assumption, the comparison map  $X \rightarrow X'$  is an acyclic cofibration and  $Y'$  is fibrant. Then it follows immediately (via lifting) that there exists a map  $\xi$  that makes the diagram

$$\begin{array}{ccc} X & \xrightarrow{\sim} & Y \\ \downarrow \sim & & \downarrow \sim \\ X' & \xrightarrow{\xi} & Y' \end{array}$$

in  $\mathbf{Alg}_{\mathcal{O}}$  commute. By Proposition 2.5,  $X'$  is TQ-local if and only if  $Y'$  is TQ-local since  $\xi$  is a weak equivalence between fibrant objects.  $\square$

**Proposition 2.7.** *Let  $X$  be an  $\mathcal{O}$ -algebra and suppose  $X', X''$  are fibrant replacements of  $X$  in  $\mathbf{Alg}_{\mathcal{O}}$ . Then  $X'$  is TQ-local if and only if  $X''$  is TQ-local.*

*Proof.* This follows from Proposition 2.6.  $\square$

The following generalization of Proposition 2.4 will be used in our proof of Propositions 2.9 and 2.10.

**Proposition 2.8.** *Let  $Y$  be a (not necessarily fibrant) object in  $\mathbf{Alg}_J$ , then every fibrant replacement of  $UY$  in  $\mathbf{Alg}_{\mathcal{O}}$  is TQ-local.*

*Proof.* Let  $UY \rightarrow \widetilde{UY}$  be a fibrant replacement of  $UY$  in  $\mathbf{Alg}_{\mathcal{O}}$ ; in particular,  $UY \rightarrow \widetilde{UY}$  is an acyclic cofibration. Let  $Y'$  be a fibrant replacement of  $Y$  in  $\mathbf{Alg}_J$ . Then it follows immediately (via lifting) that there exists a map  $\xi$  that makes the diagram

$$\begin{array}{ccc} UY & \xrightarrow{\sim} & UY' \\ \downarrow \sim & \nearrow \xi & \downarrow \\ \widetilde{UY} & \longrightarrow & * \end{array}$$

in  $\mathbf{Alg}_{\mathcal{O}}$  commute. Since  $UY'$  is TQ-local (Proposition 2.4) and  $\xi$  is a weak equivalence between fibrant objects,  $\widetilde{UY}$  is TQ-local by Proposition 2.5.  $\square$

Now we are in a good position to prove Propositions 2.9 and 2.10. First recall the  $\mathbf{TQ}$ -completion construction. Let  $Z$  be a cofibrant  $\mathcal{O}$ -algebra and consider the cosimplicial resolution of  $Z$  with respect to  $\mathbf{TQ}$ -homology of the form

$$Z \longrightarrow (UQ)Z \begin{array}{c} \xleftarrow{\quad} \\ \xrightarrow{\quad} \end{array} (UQ)^2 Z \begin{array}{c} \xleftarrow{\quad} \\ \xrightarrow{\quad} \end{array} (UQ)^3 Z \cdots$$

in  $\mathbf{Alg}_{\mathcal{O}}$ , denoted  $Z \rightarrow \mathbf{C}(Z)$ , with coface maps obtained by iterating the  $\mathbf{TQ}$ -Hurewicz map  $\mathrm{id} \rightarrow UQ$  (Definition 2.1) and codegeneracy maps built from the counit map of the adjunction  $(Q, U)$  in the usual way. Taking the homotopy limit (over  $\Delta$ ) gives the  $\mathbf{TQ}$ -completion map [13, 28] of the form

$$(2) \quad Z \rightarrow Z_{\mathbf{TQ}}^{\wedge} = \mathrm{holim}_{\Delta} \mathbf{C}(Z) \simeq \mathrm{holim}_{\Delta} \widetilde{\mathbf{C}(Z)}$$

in  $\mathbf{Alg}_{\mathcal{O}}$ , where  $\widetilde{\mathbf{C}(Z)}$  denotes any functorial fibrant replacement functor  $\widetilde{(-)}$  on  $\mathbf{Alg}_{\mathcal{O}}$  (obtained, for instance, by running the small object argument with respect to the generating acyclic cofibrations in  $\mathbf{Alg}_{\mathcal{O}}$ ) applied to the cosimplicial  $\mathcal{O}$ -algebra  $\mathbf{C}(Z)$ .

The following proposition verifies that  $\widetilde{\mathbf{C}(Z)}$  is an objectwise  $\mathbf{TQ}$ -local diagram—this amounts to checking the weak  $\mathbf{TQ}$ -fibrancy condition in the  $\mathbf{TQ}$ -local homotopy theory [29, 5.14] using the above technical propositions.

**Proposition 2.9.** *If  $Z$  is an  $\mathcal{O}$ -algebra, then the  $\mathbf{TQ}$ -completion  $Z_{\mathbf{TQ}}^{\wedge}$  of  $Z$  is the homotopy limit of a small diagram of  $\mathbf{TQ}$ -local  $\mathcal{O}$ -algebras.*

*Proof.* It suffices to consider the case when  $Z$  is a cofibrant  $\mathcal{O}$ -algebra; e.g., otherwise, work with its cofibrant replacement in  $\mathbf{Alg}_{\mathcal{O}}$ . We want to show that the  $\Delta$ -shaped diagram  $\widetilde{\mathbf{C}(Z)}$  in (2) is objectwise  $\mathbf{TQ}$ -local; i.e., that  $\widetilde{\mathbf{C}(Z)}^s$  is  $\mathbf{TQ}$ -local for each  $s \geq 0$ . This follows from Proposition 2.8. In more detail: Consider the case  $s = 0$ . Let  $Y := QZ \in \mathbf{Alg}_J$ . Then  $UY = (UQ)Z$ , hence it suffices to verify that  $\widetilde{UY}$  is  $\mathbf{TQ}$ -local which is true by Proposition 2.8. Similarly, consider the case  $s \geq 1$ . Let  $Y := Q(UQ)^s Z \in \mathbf{Alg}_J$ . Then  $UY = (UQ)^{s+1} Z$ , hence it suffices to verify that  $\widetilde{UY}$  is  $\mathbf{TQ}$ -local and Proposition 2.8 completes the proof.  $\square$

Recall the  $\mathbf{TQ}|_{\mathrm{Nil}_M}$ -completion construction from [12]. For each  $n \geq 1$ ,  $\tau_n \mathcal{O}$  is the operad associated to  $\mathcal{O}$  where

$$(\tau_n \mathcal{O})[t] := \begin{cases} \mathcal{O}[t] & \text{for } t \leq n \\ * & \text{otherwise} \end{cases}$$

and consider the associated commutative diagram of operad maps [12]

$$\begin{array}{ccccc} \mathcal{O} & \longrightarrow & J_n & \longrightarrow & J_1 = J \\ & \searrow & \downarrow \sim & & \downarrow \sim \\ & & \tau_n \mathcal{O} & \longrightarrow & \tau_1 \mathcal{O} \end{array}$$

where the upper horizontal maps are cofibrations of operads, the left-hand and bottom horizontal maps are the natural truncations, and the vertical maps are weak equivalences of operads; for notational simplicity, here we take  $J = J_1$ . The

corresponding change of operad adjunctions have the form

$$\mathrm{Alg}_{\mathcal{O}} \begin{array}{c} \xrightarrow{R_n} \\ \xleftarrow{V_n} \end{array} \mathrm{Alg}_{J_n} \begin{array}{c} \xrightarrow{Q_n} \\ \xleftarrow{U_n} \end{array} \mathrm{Alg}_J \quad \mathrm{Alg}_{\mathcal{O}} \begin{array}{c} \xrightarrow{Q} \\ \xleftarrow{U} \end{array} \mathrm{Alg}_J$$

with left adjoints on top, where  $R_n = J_n \circ_{\mathcal{O}} (-)$ ,  $Q_n = J \circ_{J_n} (-)$ ,  $Q = J \circ_{\mathcal{O}} (-)$ , and  $V_n, U_n, U$  denote the indicated forgetful functors; in particular, the adjunction on the right is the composite of the adjunctions on the left.

Next recall the  $\mathrm{TQ}|_{\mathrm{Nil}_M}$ -completion construction. Let  $n \geq 1$  and define  $M := n + 1$ . Let  $X$  be a cofibrant  $J_n$ -algebra and consider the cosimplicial resolution of  $X$  with respect to  $\mathrm{TQ}|_{\mathrm{Nil}_M}$ -homology of the form

$$X \longrightarrow (U_n Q_n) X \begin{array}{c} \xleftarrow{\quad} \\ \xrightarrow{\quad} \end{array} (U_n Q_n)^2 X \begin{array}{c} \xleftarrow{\quad} \\ \xrightarrow{\quad} \end{array} (U_n Q_n)^3 X \cdots$$

in  $\mathrm{Alg}_{J_n}$ , denoted  $X \rightarrow \mathbf{N}(X)$ , with coface maps obtained by iterating the  $\mathrm{TQ}|_{\mathrm{Nil}_M}$ -Hurewicz map  $\mathrm{id} \rightarrow U_n Q_n$  and codegeneracy maps built from the counit map of the adjunction  $(Q_n, U_n)$  in the usual way; see Ching-Harper [11]. Applying the forgetful functor  $V_n$  gives the diagram  $V_n X \rightarrow V_n \mathbf{N}(X)$  of the form

$$V_n X \longrightarrow V_n (U_n Q_n) X \begin{array}{c} \xleftarrow{\quad} \\ \xrightarrow{\quad} \end{array} V_n (U_n Q_n)^2 X \begin{array}{c} \xleftarrow{\quad} \\ \xrightarrow{\quad} \end{array} V_n (U_n Q_n)^3 X \cdots$$

in  $\mathrm{Alg}_{\mathcal{O}}$ . Taking the homotopy limit (over  $\Delta$ ) gives the  $\mathrm{TQ}|_{\mathrm{Nil}_M}$ -completion map of the form

$$(3) \quad X \rightarrow X_{\mathrm{TQ}|_{\mathrm{Nil}_M}}^{\wedge} = \mathrm{holim}_{\Delta} V_n \mathbf{N}(X) \simeq \mathrm{holim}_{\Delta} \widetilde{V_n \mathbf{N}(X)}$$

in  $\mathrm{Alg}_{\mathcal{O}}$ , where  $\widetilde{V_n \mathbf{N}(X)}$  denotes any functorial fibrant replacement functor  $\widetilde{(-)}$  on  $\mathrm{Alg}_{\mathcal{O}}$  applied to the cosimplicial  $\mathcal{O}$ -algebra  $V_n \mathbf{N}(X)$ .

**Proposition 2.10.** *If  $X$  is a  $J_n$ -algebra, then the  $\mathrm{TQ}|_{\mathrm{Nil}_M}$ -completion  $X_{\mathrm{TQ}|_{\mathrm{Nil}_M}}^{\wedge}$  of  $X$  is the homotopy limit of a small diagram of  $\mathrm{TQ}$ -local  $\mathcal{O}$ -algebras.*

*Proof.* It suffices to consider the case when  $X$  is a cofibrant  $J_n$ -algebra; e.g., otherwise, work with its cofibrant replacement in  $\mathrm{Alg}_{J_n}$ . We want to show that the  $\Delta$ -shaped diagram  $V_n \mathbf{N}(X)$  in (3) is objectwise  $\mathrm{TQ}$ -local; i.e., that  $\widetilde{V_n \mathbf{N}(X)}^s$  is  $\mathrm{TQ}$ -local for each  $s \geq 0$ . This follows from Proposition 2.8. In more detail: Consider the case  $s = 0$ . Let  $Y := Q_n X \in \mathrm{Alg}_J$ , then  $UY = V_n(U_n Q_n) X$  and hence it suffices to verify that  $\widetilde{UY}$  is  $\mathrm{TQ}$ -local; this is true by Proposition 2.8. Similarly, consider the case  $s \geq 1$ . Let  $Y := Q_n(U_n Q_n)^s X \in \mathrm{Alg}_J$ . Then  $UY = V_n(U_n Q_n)^{s+1} X$  and hence it suffices to verify that  $\widetilde{UY}$  is  $\mathrm{TQ}$ -local; this is true by Proposition 2.8 which completes the proof.  $\square$

### 3. COMPARING $\mathrm{TQ}$ -LOCALIZATION WITH $\mathrm{TQ}$ -COMPLETION

In this section we discuss the relation between  $\mathrm{TQ}$ -localization and  $\mathrm{TQ}$ -completion. Let  $X$  be a cofibrant  $\mathcal{O}$ -algebra. One can construct the  $\mathrm{TQ}$ -localization map  $X \rightarrow L_{\mathrm{TQ}}(X)$  (see [29]) by running the small object argument. By construction,  $L_{\mathrm{TQ}}(X)$  is  $\mathrm{TQ}$ -local and the  $\mathrm{TQ}$ -localization map  $X \rightarrow L_{\mathrm{TQ}}(X)$  is a  $\mathrm{TQ}$ -acyclic strong cofibration (Definition 2.2).

The  $\mathrm{TQ}$ -completion map  $X \rightarrow X_{\mathrm{TQ}}^{\wedge}$  can be thought of as an approximation of the  $\mathrm{TQ}$ -localization map. For instance, we know that the  $\mathrm{TQ}$ -completion  $X_{\mathrm{TQ}}^{\wedge}$  of  $X$  is always  $\mathrm{TQ}$ -local (Proposition 1.6).



**Theorem 3.1** (Recognizing when  $\mathbf{TQ}$ -local  $\mathcal{O}$ -algebras are  $\mathbf{TQ}$ -complete). *Let  $X$  be a cofibrant  $\mathcal{O}$ -algebra. Then the  $\mathbf{TQ}$ -completion map  $c: X \rightarrow X_{\mathbf{TQ}}^\wedge$  factors through the  $\mathbf{TQ}$ -localization map  $l: X \rightarrow L_{\mathbf{TQ}}(X)$  via a commutative diagram of the form*

$$\begin{array}{ccc} X & \xrightarrow{c} & X_{\mathbf{TQ}}^\wedge \\ \downarrow l & \nearrow \xi & \downarrow \\ L_{\mathbf{TQ}}(X) & \longrightarrow & * \end{array}$$

in  $\mathbf{Alg}_{\mathcal{O}}$ . Furthermore, if  $X$  is  $\mathbf{TQ}$ -local, then the following are equivalent:

- (i) The natural map  $X \rightarrow X_{\mathbf{TQ}}^\wedge$  is a  $\mathbf{TQ}$ -equivalence; i.e.,  $X$  is  $\mathbf{TQ}$ -good.
- (ii) The natural map  $X \simeq X_{\mathbf{TQ}}^\wedge$  is a weak equivalence; i.e.,  $X$  is  $\mathbf{TQ}$ -complete.
- (iii) The comparison map  $\xi$  is a weak equivalence.

*Proof.* This is analogous to the Bousfield-Kan completion of spaces [10]. Since  $X_{\mathbf{TQ}}^\wedge$  is  $\mathbf{TQ}$ -local and  $l: X \rightarrow L_{\mathbf{TQ}}(X)$  is a  $\mathbf{TQ}$ -acyclic strong cofibration, there exists a lift  $\xi$  that makes the diagram commute (Proposition 2.3) in  $\mathbf{Alg}_{\mathcal{O}}$ . Suppose  $X$  is  $\mathbf{TQ}$ -local, then  $l$  is a  $\mathbf{TQ}$ -equivalence between  $\mathbf{TQ}$ -local objects, hence a weak equivalence by the  $\mathbf{TQ}$ -local Whitehead theorem (Proposition 1.4). Therefore  $\xi$  is a weak equivalence if and only if  $c$  is a weak equivalence. This verifies (ii)  $\Leftrightarrow$  (iii). Since  $X, X_{\mathbf{TQ}}^\wedge$  are  $\mathbf{TQ}$ -local,  $c$  is a  $\mathbf{TQ}$ -equivalence if and only if  $c$  is a weak equivalence by the  $\mathbf{TQ}$ -local Whitehead theorem. This verifies (i)  $\Leftrightarrow$  (ii).  $\square$

It is worth pointing out the following two propositions.

**Proposition 3.2.** *A map  $f: X \rightarrow Y$  between  $\mathcal{O}$ -algebras is a  $\mathbf{TQ}$ -homology equivalence if and only if the induced map  $f_{\mathbf{TQ}}^\wedge: X_{\mathbf{TQ}}^\wedge \rightarrow Y_{\mathbf{TQ}}^\wedge$  is a weak equivalence.*

*Proof.* This is proved by arguing exactly as in [10, I.5], but here is the basic idea: The “if” direction is proved using retract argument and the “only if” direction is because  $\mathrm{holim}_\Delta$  preserves weak equivalences.  $\square$

**Proposition 3.3.** *Let  $X$  be an  $\mathcal{O}$ -algebra, then the following are equivalent:*

- (i)  $X$  is  $\mathbf{TQ}$ -good.
- (ii)  $X_{\mathbf{TQ}}^\wedge$  is  $\mathbf{TQ}$ -complete.
- (iii)  $X_{\mathbf{TQ}}^\wedge$  is  $\mathbf{TQ}$ -good.

*Proof.* This follows from exactly the same argument as in [10, I.5].  $\square$

#### 4. POSTNIKOV TOWERS AND $\mathbf{TQ}$ -LOCALIZATION

In this section we assume that  $\mathcal{O}, \mathcal{R}$  are  $(-1)$ -connected. We show that a  $(-1)$ -connected  $\mathcal{O}$ -algebra is  $\mathbf{TQ}$ -local if it has a principally refined Postnikov tower.

**Proposition 4.1.** *Let  $X$  be a  $(-1)$ -connected cofibrant  $\mathcal{O}$ -algebra. Then there exists a coaugmented tower  $\{X\} \rightarrow \{X_n\}$  (the Postnikov tower of  $X$ ) of the form*

$$\begin{array}{c} X \\ \downarrow \\ * = X_{-1} \longleftarrow X_0 \longleftarrow X_1 \longleftarrow X_2 \longleftarrow \cdots \end{array}$$

in  $\mathbf{Alg}_{\mathcal{O}}$  such that for each  $n \geq -1$ :

- (a)  $X_n$  is a cofibrant and fibrant  $\mathcal{O}$ -algebra.

- (b) the structure map  $X \rightarrow X_n$  is  $(n+1)$ -connected and  $\pi_k X_n = *$  for all  $k \geq n+1$ .
- (c) the structure map  $X_{n+1} \rightarrow X_n$  is a fibration.

*Proof.* The Postnikov tower can be constructed using small object arguments analogous to the arguments in [11, 15, 17, 47]. In more detail: Let  $I_n$  be the set of  $n$ -connected generating cofibrations in  $\mathbf{Alg}_{\mathcal{O}}$  and let  $J$  be the set of generating acyclic cofibrations in  $\mathbf{Alg}_{\mathcal{O}}$  (see, for instance, [26]). Start by setting  $X_{-1} = *$ . For each  $n \geq 0$ , we inductively run the small object argument with respect to  $I_{n+1} \cup J$  to factor the map  $X \rightarrow X_{n-1}$  in  $\mathbf{Alg}_{\mathcal{O}}$  as  $X \rightarrow X_n \rightarrow X_{n-1}$ . Then  $X \rightarrow X_n$  is a cofibration,  $X_n \rightarrow X_{n-1}$  is a fibration and  $\pi_k X_n = *$  for all  $k \geq n+1$ . By assumption,  $\mathcal{O}, \mathcal{R}$  are  $(-1)$ -connected, hence  $X \rightarrow X_n$  is  $(n+1)$ -connected by the small object argument construction.  $\square$

Analogous to the definition for spaces, principal Postnikov towers and principally refined Postnikov towers are defined as follows.

**Definition 4.2.** Let  $X$  be a  $(-1)$ -connected  $\mathcal{O}$ -algebra. We say that a Postnikov tower  $\{X_n\}$  of  $X$  is *principal* if for each  $n \geq 0$ , the structure map  $X_n \rightarrow X_{n-1}$  fits into a homotopy pullback diagram of the left-hand form

$$(4) \quad \begin{array}{ccc} X_n & \longrightarrow & *^{\text{fat}} \\ \downarrow & & \downarrow \\ X_{n-1} & \longrightarrow & K(\pi_n X, n+1) \end{array} \quad \begin{array}{ccc} M_i & \longrightarrow & *^{\text{fat}} \\ \downarrow & & \downarrow \\ M_{i-1} & \longrightarrow & K(G_i, n+1) \end{array}$$

in  $\mathbf{Alg}_{\mathcal{O}}$ , where  $*^{\text{fat}}$  is an  $\mathcal{O}$ -algebra that is weakly equivalent to  $*$  (i.e., a “fat point” in  $\mathbf{Alg}_{\mathcal{O}}$ ) and  $K(\pi_n X, n+1)$  is an object in  $\mathbf{Alg}_{\mathcal{O}}$  with  $\pi_n X$  as the only nontrivial homotopy group concentrated at level  $n+1$ .

We say that  $\{X_n\}$  is *principally refined* if, for each  $n \geq 0$ , the structure map  $X_n \rightarrow X_{n-1}$  can be factored as a finite composite  $X_n = M_{t_n} \rightarrow \cdots \rightarrow M_2 \rightarrow M_1 \rightarrow M_0 = X_{n-1}$  of maps such that, for each  $t_n \geq i \geq 1$ , the map  $M_i \rightarrow M_{i-1}$  fits into a homotopy pullback diagram of the right-hand form (4) in  $\mathbf{Alg}_{\mathcal{O}}$ , where the  $G_i$ ’s are abelian groups and  $K(G_i, n+1)$  is an object in  $\mathbf{Alg}_{\mathcal{O}}$  with  $G_i$  as the only nontrivial homotopy group concentrated at level  $n+1$ . In particular, every principal Postnikov tower is principally refined.

*Proof of Theorem 1.11.* We know that every 0-connected fibrant  $\mathcal{O}$ -algebra is  $\mathbf{TQ}$ -local (Theorem 1.8), hence, in particular, each Eilenberg-MacLane object  $K(G_i, n+1)$  appearing in the principally refined Postnikov tower of  $X$  has  $\mathbf{TQ}$ -local fibrant replacements in  $\mathbf{Alg}_{\mathcal{O}}$ . By inducting up the principally refined Postnikov tower, it follows that each  $X_n$  is  $\mathbf{TQ}$ -local (Proposition 1.5). Since  $X$  is the homotopy limit of its Postnikov tower  $\{X_n\}$ , which is objectwise  $\mathbf{TQ}$ -local, it follows that  $X$  is  $\mathbf{TQ}$ -local (Proposition 1.5) which completes the proof.  $\square$

We provide some examples of  $(-1)$ -connected algebras which admit principally refined Postnikov towers.

- (i) Let  $X$  be a cofibrant 0-connected  $\mathcal{O}$ -algebra. Analogous to results in [4, 7, 15, 30], one can show that the Postnikov tower of  $X$  is principal.
- (ii) Consider  $\Omega X$  for any 0-connected cofibrant  $\mathcal{O}$ -algebra  $X$ . Since the loop functor  $\Omega$  commutes with homotopy pullbacks in  $\mathbf{Alg}_{\mathcal{O}}$ ,  $\Omega X$  has a principal Postnikov tower by applying  $\Omega$  to the principal Postnikov tower of  $X$ .

- (iii) Consider  $UY$  for any  $(-1)$ -connected cofibrant  $J$ -algebra  $Y$ . The category  $\mathbf{Alg}_J$  is Quillen equivalent to  $\mathbf{Alg}_{\tau_1 \mathcal{O}} \cong \mathbf{Mod}_{\mathcal{O}[1]}$  [28, 7.21], hence the homotopy category of  $\mathbf{Alg}_J$  is stable. Therefore, the Postnikov tower of  $Y$  in  $\mathbf{Alg}_J$  is already principal. Applying forgetful functor  $U$  induces principal Postnikov tower for  $UY$  in  $\mathbf{Alg}_{\mathcal{O}}$ .
- (iv) One can construct additional examples by pulling back quotient towers along cocellular maps as described in [40, 3.3].

## REFERENCES

- [1] M. André. *Homologie des algèbres commutatives*. Springer-Verlag, Berlin, 1974. Die Grundlehren der mathematischen Wissenschaften, Band 206.
- [2] M. Artin and B. Mazur. *Etale homotopy*. Lecture Notes in Mathematics, No. 100. Springer-Verlag, Berlin-New York, 1969.
- [3] A. Baker and B. Richter, editors. *Structured ring spectra*, volume 315 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 2004.
- [4] M. Bastera. André-Quillen cohomology of commutative  $S$ -algebras. *J. Pure Appl. Algebra*, 144(2):111–143, 1999.
- [5] M. Bastera and M. A. Mandell. Homology and cohomology of  $E_{\infty}$  ring spectra. *Math. Z.*, 249(4):903–944, 2005.
- [6] M. Bastera and M. A. Mandell. Homology of  $E_n$  ring spectra and iterated  $THH$ . *Algebr. Geom. Topol.*, 11(2):939–981, 2011.
- [7] M. Bastera and M. A. Mandell. The multiplication on BP. *Journal of Topology*, 6(2):285–310, 2013.
- [8] H. J. Baues. *Combinatorial foundation of homology and homotopy*. Springer Monographs in Mathematics. Springer-Verlag, Berlin, 1999.
- [9] A. K. Bousfield. The localization of spaces with respect to homology. *Topology*, 14:133–150, 1975.
- [10] A. K. Bousfield and D. M. Kan. *Homotopy limits, completions and localizations*. Lecture Notes in Mathematics, Vol. 304. Springer-Verlag, Berlin, 1972.
- [11] M. Ching and J. E. Harper. Higher homotopy excision and Blakers-Massey theorems for structured ring spectra. *Adv. Math.*, 298:654–692, 2016.
- [12] M. Ching and J. E. Harper. A nilpotent Whitehead theorem for TQ-homology of structured ring spectra. *Tbilisi Math. J.*, 11:69–79, 2018.
- [13] M. Ching and J. E. Harper. Derived Koszul duality and TQ-homology completion of structured ring spectra. *Adv. Math.*, 341:118–187, 2019.
- [14] E. Dror Farjoun. *Cellular spaces, null spaces and homotopy localization*, volume 1622 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1996.
- [15] D. Dugger and B. Shipley. Postnikov extensions of ring spectra. *Algebr. Geom. Topol.*, 6:1785–1829 (electronic), 2006.
- [16] W. G. Dwyer. Localizations. In *Axiomatic, enriched and motivic homotopy theory*, volume 131 of *NATO Sci. Ser. II Math. Phys. Chem.*, pages 3–28. Kluwer Acad. Publ., Dordrecht, 2004.
- [17] A. D. Elmendorf, I. Kriz, M. A. Mandell, and J. P. May. *Rings, modules, and algebras in stable homotopy theory*, volume 47 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 1997. With an appendix by M. Cole.
- [18] J. Francis and D. Gaijsory. Chiral Koszul duality. *Selecta Math. (N.S.)*, 18(1):27–87, 2012.
- [19] S. Galatius, A. Kupers, and O. Randal-Williams. Cellular  $E_k$ -algebras. [arXiv:1805.07184 \[math.AT\]](https://arxiv.org/abs/1805.07184), 2018.
- [20] P. G. Goerss. On the André-Quillen cohomology of commutative  $\mathbf{F}_2$ -algebras. *Astérisque*, 186:169, 1990.
- [21] P. G. Goerss and M. J. Hopkins. André-Quillen (co)-homology for simplicial algebras over simplicial operads. In *Une dégustation topologique [Topological morsels]: homotopy theory in the Swiss Alps (Arolla, 1999)*, volume 265 of *Contemp. Math.*, pages 41–85. Amer. Math. Soc., Providence, RI, 2000.
- [22] P. G. Goerss and M. J. Hopkins. Moduli problems for structured ring spectra. *preprint*, 2005. Available at <http://hopf.math.purdue.edu>.

- [23] P. G. Goerss and J. F. Jardine. *Simplicial homotopy theory*, volume 174 of *Progress in Mathematics*. Birkhäuser Verlag, Basel, 1999.
- [24] P. G. Goerss and K. Schemmerhorn. Model categories and simplicial methods. In *Interactions between homotopy theory and algebra*, volume 436 of *Contemp. Math.*, pages 3–49. Amer. Math. Soc., Providence, RI, 2007.
- [25] T. G. Goodwillie. Calculus. III. Taylor series. *Geom. Topol.*, 7:645–711, 2003.
- [26] J. E. Harper. Homotopy theory of modules over operads in symmetric spectra. *Algebr. Geom. Topol.*, 9(3):1637–1680, 2009. Corrigendum: *Algebr. Geom. Topol.*, 15(2):1229–1237, 2015.
- [27] J. E. Harper. Bar constructions and Quillen homology of modules over operads. *Algebr. Geom. Topol.*, 10(1):87–136, 2010.
- [28] J. E. Harper and K. Hess. Homotopy completion and topological Quillen homology of structured ring spectra. *Geom. Topol.*, 17(3):1325–1416, 2013.
- [29] J. E. Harper and Y. Zhang. Topological Quillen localization of structured ring spectra. *Tbilisi Math. J.*, 12:67–89, 2019.
- [30] A. Hatcher. *Algebraic Topology*. Cambridge University Press, 2002.
- [31] P. Hilton, G. Mislin, and J. Roitberg. *Localization of nilpotent groups and spaces*. North-Holland Publishing Co., Amsterdam-Oxford; American Elsevier Publishing Co., Inc., New York, 1975. North-Holland Mathematics Studies, No. 15, Notas de Matemática, No. 55. [Notes on Mathematics, No. 55].
- [32] P. S. Hirschhorn. *Model categories and their localizations*, volume 99 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2003.
- [33] M. Hovey, B. Shipley, and J. H. Smith. Symmetric spectra. *J. Amer. Math. Soc.*, 13(1):149–208, 2000.
- [34] J. F. Jardine. *Local homotopy theory*. Springer Monographs in Mathematics. Springer, New York, 2015.
- [35] I. Kriz. Towers of  $E_\infty$  ring spectra with an application to BP. *preprint*, 1993.
- [36] N. J. Kuhn. Localization of André-Quillen-Goodwillie towers, and the periodic homology of infinite loopspaces. *Adv. Math.*, 201(2):318–378, 2006.
- [37] N. J. Kuhn and L. A. Pereira. Operad bimodules and composition products on André-Quillen filtrations of algebras. *Algebr. Geom. Topol.*, 17(2):1105–1130, 2017.
- [38] A. Lazarev. Cohomology theories for highly structured ring spectra. In *Structured ring spectra*, volume 315 of *London Math. Soc. Lecture Note Ser.*, pages 201–231. Cambridge Univ. Press, Cambridge, 2004.
- [39] M. A. Mandell. Topological André-Quillen cohomology and  $E_\infty$  André-Quillen cohomology. *Adv. Math.*, 177(2):227–279, 2003.
- [40] J. P. May and K. Ponto. *More concise algebraic topology*. Chicago Lectures in Mathematics. University of Chicago Press, Chicago, IL, 2012. Localization, completion, and model categories.
- [41] H. R. Miller. The Sullivan conjecture on maps from classifying spaces. *Ann. of Math. (2)*, 120(1):39–87, 1984. Correction: *Ann. of Math. (2)*, 121(3):605–609, 1985.
- [42] L. A. Pereira. Goodwillie calculus in the category of algebras over a spectral operad. 2013. Available at: <http://math.mit.edu/~luisalex/>.
- [43] D. Quillen. *Homotopical algebra*. Lecture Notes in Mathematics, No. 43. Springer-Verlag, Berlin, 1967.
- [44] D. Quillen. On the (co-) homology of commutative rings. In *Applications of Categorical Algebra (Proc. Sympos. Pure Math., Vol. XVII, New York, 1968)*, pages 65–87. Amer. Math. Soc., Providence, R.I., 1970.
- [45] S. Schwede. Spectra in model categories and applications to the algebraic cotangent complex. *J. Pure Appl. Algebra*, 120(1):77–104, 1997.
- [46] S. Schwede. Stable homotopy of algebraic theories. *Topology*, 40(1):1–41, 2001.
- [47] S. Schwede. Symmetric spectra. Available at: <http://www.math.uni-bonn.de/people/schwede/SymSpec.pdf>, 2012.
- [48] D. Sullivan. *Geometric topology. Part I*. Massachusetts Institute of Technology, Cambridge, Mass., 1971. Localization, periodicity, and Galois symmetry, Revised version.

DEPARTMENT OF MATHEMATICS, THE OHIO STATE UNIVERSITY, 231 WEST 18TH AVE, COLUMBUS, OH 43210, USA

Email address: [zhang.4841@osu.edu](mailto:zhang.4841@osu.edu)