

# THE SECONDARY PERIODIC ELEMENT $\beta_{p^2/p^2-1}$ AND ITS APPLICATIONS

JIANGUO HONG, XIANGJUN WANG, AND YU ZHANG\*

ABSTRACT. In this paper, we prove that  $\beta_{p^2/p^2-1}$  survives to  $E_\infty$  in the Adams-Novikov spectral sequence for all  $p \geq 5$ . From the Thom map  $\Phi : Ext_{BP_*BP}^{s,t}(BP_*, BP_*) \rightarrow Ext_A^{s,t}(\mathbb{Z}/p, \mathbb{Z}/p)$ , we also see that  $h_0h_3$  survives to  $E_\infty$  in the classical Adams spectral sequence. As an application, we prove that  $\beta_{p/p}^p$  is divisible by  $\beta_1$ .

## 1. INTRODUCTION

Let  $p \geq 5$  be an odd prime. The Adams-Novikov spectral sequence (ANSS) based on the Brown-Peterson spectrum  $BP$  is one of the most powerful tools to compute the  $p$ -component of the stable homotopy groups of spheres  $\pi_*(S^0)$  (cf. [1, 6, 10, 22]).

The  $E_2$ -term of the ANSS is  $Ext_{BP_*BP}^{s,t}(BP_*, BP_*)$ , which has been extensively studied in low dimensions. For  $s = 1$ ,  $Ext_{BP_*BP}^{1,*}(BP_*, BP_*)$  is generated by  $\alpha_{sp^n/n+1}$  for  $n \geq 0$ ,  $p \nmid s \geq 1$ , where  $\alpha_{sp^n/n+1}$  has order  $p^{n+1}$  (cf. [12, 10]). For  $s = 2$ ,  $Ext_{BP_*BP}^{2,*}(BP_*, BP_*)$  is the direct sum of cyclic groups generated by  $\beta_{sp^n/j, i+1}$  for suitable  $(n, s, j, i)$  (cf. [10, 22, 23]),  $\beta_{sp^n/j, i+1}$  has order  $p^{i+1}$ . For  $s \geq 3$ , only partial results of  $Ext_{BP_*BP}^{s,*}(BP_*, BP_*)$  are known (cf. [11]).

In order to compute the stable homotopy groups of the sphere, we still need to know which elements of the  $E_2$ -page could survive to the  $E_\infty$ -page of the ANSS. It is known that each element  $\alpha_{sp^n/n+1}$  is a permanent cycle in the ANSS which represents an element of  $\text{Im} J$  with the same order. But we are far from fully determining which elements of the  $\beta_{sp^n/j, i+1}$  family could survive to  $E_\infty$ .

Let  $\beta_{sp^n/j}$  denote  $\beta_{sp^n/j, 1}$ . H. Toda [26, 27] proved that  $\alpha_1\beta_1^p$  is zero in  $\pi_*(S^0)$ . This relation supports a non-trivial Adams-Novikov differential called the Toda differential

$$(1.1) \quad d_{2p-1}(\beta_{p/p}) = a \cdot \alpha_1\beta_1^p \neq 0$$

where  $a$  is a non-zero scalar *mod*  $p$ . Hence  $\beta_{p/p}$  could not survive the ANSS.

Based on the Toda differential (1.1), D. Ravenel [19] generalized the result and proved that there are nontrivial differentials

$$d_{2p-1}(\beta_{p^n/p^n}) \equiv a \cdot \alpha_1\beta_{p^{n-1}/p^{n-1}}^p, \quad \text{mod } \ker \beta_1^{p(p^{n-1}-1)/(p-1)}$$

for  $n \geq 1$ . Consequently,  $\beta_{p^n/p^n}$  also can not survive to  $E_\infty$  in the ANSS. From this one can see that only  $\beta_{sp^n/j} \in H^2(BP_*)$  for  $s \geq 2$ ,  $1 \leq j \leq p^n$  or  $s = 1$ ,  $1 \leq j \leq p^n - 1$  might survive to  $E_\infty$  in the ANSS. The following are some known results in this area:

Oka [13] proved that for  $s = 1$ ,  $1 \leq j \leq p - 1$  or  $s \geq 2$ ,  $1 \leq j \leq p$ ,  $\beta_{sp/j}$  is a permanent cycle in the ANSS.

---

2020 *Mathematics Subject Classification.* 18G40, 55Q45, 55T15.

*Key words and phrases.* stable homotopy groups of spheres, Adams-Novikov spectral sequence, infinite descent method.

The authors were supported by NSFC grant No. 11871284.

\* Corresponding author.

Oka [15] proved that for  $s = 1$ ,  $1 \leq j \leq 2p - 2$  or  $s \geq 2$ ,  $1 \leq j \leq 2p$ ,  $\beta_{sp^2/j}$  is a permanent cycle in the ANSS.

Later Oka [17, 18] generalized the result to  $n \geq 2$ , i.e. for  $n \geq 2$ ;  $s = 1$ ,  $1 \leq j \leq 2^{n-1}(p-1)$  or  $s \geq 2$ ,  $1 \leq j \leq 2^{n-1}p$ ,  $\beta_{sp^n/j}$  survives to  $E_\infty$  in the ANSS.

Shimomura [25] proved that for  $s \geq 1$ ,  $1 \leq j \leq p^2 - 2$ ,  $\beta_{sp^2/j}$  survives to  $E_\infty$  in the ANSS.

In this paper, we prove:

**Theorem A** *Let  $p \geq 5$  be an odd prime. Then  $\beta_{p^2/p^2-1}$  is a permanent cycle in the Adams-Novikov spectral sequence.*

Let  $M$  be the mod  $p$  Moore spectrum and  $M(1, p^n - 1)$  be the cofiber of the map  $v_1^{p^n-1}$

$$\Sigma^* M \xrightarrow{v_1^{p^n-1}} M \longrightarrow M(1, p^n - 1).$$

D. Ravenel ([24] Theorem 7.12) claimed that if  $M(1, p^n - 1)$  is a ring spectrum and  $\beta_{p^n/p^n-1}$  is a permanent cycle, then  $\beta_{sp^n/j}$  is a permanent cycle for all  $s \geq 1$ ,  $j \leq p^n - 1$ .

Between the ANSS and the classical Adams spectral sequence (ASS), there is the Thom reduction map

$$\Phi : Ext_{BP_*BP}^*(BP_*, BP_*) \longrightarrow Ext_A^*(\mathbb{Z}/p, \mathbb{Z}/p)$$

such that  $\Phi(\beta_{p^n/p^n-1}) = h_0 h_{n+1}$ . Thus

**Corollary B** *Let  $p \geq 5$  be an odd prime. Then  $h_0 h_3$  is a permanent cycle in the classical Adams spectral sequence.*

In [3] R. Cohen and P. Goerss claimed the existence of  $h_0 h_{n+1}$  in the classical ASS. One can see that the existence of  $h_0 h_{n+1}$  in ASS is equivalent to the existence of  $\beta_{p^n/p^n-1}$  in the Adams-Novikov spectral sequence. But N. Minami found a fatal error in their proof, so it is still an open problem in odd primary stable homotopy theory. Due to its extreme importance, M. Hovey [4] listed the convergence of  $h_0 h_{n+1}$  as one of the major open problems in algebraic topology.

Consider the ANSS for the Moore spectrum  $Ext_{BP_*BP}^{*,*}(BP_*, BP_*(M)) \implies \pi_*(M)$ . From the Toda differential, one can see that in the ANSS for the Moore spectrum

$$d_{2p-1}(h_{n+2}) = v_1 \beta_{p^n/p^n}^p, \quad d_{2p-1}(v_1 h_{n+2}) = v_1^2 \beta_{p^n/p^n}^p.$$

Applying the connecting homomorphism  $\delta : Ext_{BP_*BP}^{1,*}(BP_*, BP_*(M)) \longrightarrow Ext_{BP_*BP}^{2,*}(BP_*, BP_*)$  induced by the cofiber sequence

$$S^0 \xrightarrow{p} S^0 \longrightarrow M$$

one get an Adams differential in the ANSS for sphere

$$d_{2p-1}(\beta_{p^{n+1}/p^{n+1}-1}) = \alpha_2 \beta_{p^n/p^n}^p.$$

In Section 6, we prove that  $\beta_{p/p}^p$  is divisible by  $\beta_1$ , i.e.  $\beta_{p/p}^p = \beta_1 \mathbf{g}$ . Note  $\alpha_2 \beta_1 = 0$ , this provides another perspective for understanding why we could have

$$d_{2p-1}(\beta_{p^2/p^2-1}) = \alpha_2 \beta_{p/p}^p = 0 \text{ in } Ext_{BP_*BP}^{2p+1,*}(BP_*, BP_*).$$

in Theorem A.

Based on the analysis of  $\beta_{p/p}^p$ , we conjecture that:

**Conjecture C** For  $n < p - 1$ ,  $\beta_{p^n/p^n}^p$  is divisible by  $\beta_1$  and

$$\begin{aligned}\beta_{p/p}^p &= \beta_1 h_{11} b_{20}^{p-3} \gamma_2 \\ \beta_{p^2/p^2}^p &= \beta_1 h_{21} h_{11} b_{30}^{p-4} \delta_3 \\ &\dots \\ \beta_{p^i/p^i}^p &= \beta_1 h_{i,1} h_{i-1,1} \dots h_{11} b_{i+1,0}^{p-i-2} \alpha_{i+1}^{(i+2)} \\ &\dots \\ \beta_{p^{p-2}/p^{p-2}}^p &= \beta_1 h_{p-2,1} h_{p-3,1} \dots h_{11} \alpha_{p-1}^{(p)}\end{aligned}$$

where  $\alpha_{i+1}^{(i+2)}$  is the  $i + 2$ -th Greek letter elements. Thus  $\alpha_2 \beta_{p^n/p^n}^p = \alpha_2 \beta_1 \mathfrak{g} = 0$  for  $n < p - 1$ .

For  $n \geq p - 1$ ,  $\beta_{p^n/p^n}^p$  is not divisible by  $\beta_1$  and  $\alpha_2 \beta_{p^n/p^n}^p$  might be non-zero. This implies that  $\beta_{p^{n+1}/p^{n+1}-1}$  does not survives to  $E_\infty$  in the ANSS when  $n \geq p - 1$ .

This paper is arranged as follows. In section 2 we recall the construction of the topological small descent spectral sequence (TSDSS) and the small descent spectral sequence (SDSS), where the SDSS is a spectral sequence that converges to  $Ext_{BP_*BP}^{s,t}(BP_*, BP_*)$  started from the  $Ext$  groups of a complex with  $p$ -cells. Then we describe the  $E_1$ -terms of the SDSS in the form of Generator, total degree  $t - s$  and  $t - s \bmod pq - 2$  and range of index. This gives a method to compute the  $E_2$ -page of the ANSS with specialized  $t - s$ . In section 3 we compute the Adams-Novikov  $E_2$ -term  $Ext_{BP_*BP}^{s,t}(BP_*, BP_*)$  subject to  $t - s = q(p^3 + 1) - 3$  by the SDSS. In section 4, a non-trivial Adams-Novikov differential  $d_{2p-1}(h_{20} b_{11} \gamma_s) = \alpha_1 \beta_1^p h_{20} \gamma_s$  is proved. From which we prove our main theorem by showing that  $d_r(\beta_{p^2/p^2-1}) = 0$  in section 5. At last, in section 6, we prove that  $\beta_{p/p}^p$  is divisible by  $\beta_1$  and give our conjecture.

## 2. THE SMALL DESCENT SPECTRAL SEQUENCE AND THE ABC THEOREM

In 1985, D. Ravenel [20, 21, 22, 23] introduced the *method of infinite descent* and used it to compute the first thousand stems of the stable homotopy groups of spheres at the prime 5. This method is an approach to finding the  $E_2$ -term of the ANSS by the spectral sequence referred as the *small descent spectral sequence* (SDSS).

Hereafter we set that  $q = 2p - 2$ . Let  $T(n)$  be the Ranevel spectrum (cf. [22] Section 5, Chapter 6) characterized by

$$BP_*(T(n)) = BP_*[t_1, t_2, \dots, t_n].$$

Then we have the following diagram

$$S^0 = T(0) \longrightarrow T(1) \longrightarrow T(2) \longrightarrow \dots \longrightarrow T(n) \longrightarrow \dots \longrightarrow BP,$$

where  $S^0$  denote the sphere spectrum localized at an odd prime  $p \geq 5$ . Let  $T(0)_{p-1}$  and  $T(0)_{p-2}$  denote the  $q(p-1)$  and  $q(p-2)$  skeletons of  $T(1)$  respectively, they are denoted by  $X$  and  $\bar{X}$  for simple. Then

$$X = S^0 \cup_{\alpha_1} e^q \cup \dots \cup_{\alpha_1} e^{(p-2)q} \cup_{\alpha_1} e^{(p-1)q} \quad \text{and} \quad \bar{X} = S^0 \cup_{\alpha_1} e^q \cup \dots \cup_{\alpha_1} e^{(p-2)q}.$$

The  $BP$ -homologies of them are

$$BP_*(X) = BP_*[t_1]/\langle t_1^p \rangle \quad \text{and} \quad BP_*(\bar{X}) = BP_*[t_1]/\langle t_1^{p-1} \rangle.$$

From the definition above we get the following cofibre sequences

$$(2.1) \quad S^0 \xrightarrow{i'} X \xrightarrow{j'} \Sigma^q \bar{X} \xrightarrow{k'} S^1,$$

$$(2.2) \quad \overline{X} \xrightarrow{i''} X \xrightarrow{j''} S^{(p-1)q} \xrightarrow{k''} \Sigma \overline{X},$$

and the short exact sequences of  $BP$ -homologies

$$(2.3) \quad 0 \longrightarrow BP_*(S^0) \xrightarrow{i'_*} BP_*(X) \xrightarrow{j'_*} BP_*(\Sigma^q \overline{X}) \longrightarrow 0,$$

$$(2.4) \quad 0 \longrightarrow BP_*(\overline{X}) \xrightarrow{i''_*} BP_*(X) \xrightarrow{j''_*} BP_*(S^{(p-1)q}) \longrightarrow 0.$$

Put (2.3) and (2.4) together, one has the following long exact sequence

$$(2.5) \quad 0 \longrightarrow BP_*(S^0) \longrightarrow BP_*(X) \longrightarrow BP_*(\Sigma^q X) \longrightarrow BP_*(\Sigma^{pq} X) \longrightarrow \dots$$

Put (2.1) and (2.2) together, one has the following Adams diagram of cofibres

$$(2.6) \quad \begin{array}{ccccccc} S^0 & \longleftarrow & \Sigma^{q-1} \overline{X} & \longleftarrow & S^{pq-2} & \longleftarrow & \Sigma^{(p+1)q-3} \overline{X} \longleftarrow \dots \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ X & & \Sigma^{q-1} X & & \Sigma^{pq-2} X & & \Sigma^{(p+1)q-3} X. \end{array}$$

Thus one has:

**Proposition 2.1** (Ravenel [22] 7.4.2 Proposition) *Let  $X$  be as above. Then*

(a) *There is a spectral sequence converging to  $Ext_{BP_*BP}^{s+u,*}(BP_*, BP_*(S^0))$  with  $E_1$ -term*

$$E_1^{s,t,u} = Ext_{BP_*BP}^{s,t}(BP_*, BP_*(X)) \otimes E[\alpha_1] \otimes P[\beta_1], \quad \text{where} \\ E_1^{s,t,0} = Ext_{BP_*BP}^{s,t}(BP_*, BP_*(X)), \quad \alpha_1 \in E_1^{0,q,1}, \quad \beta_1 \in E_1^{0,q,p,2}$$

and  $d_r : E_r^{s,t,u} \longrightarrow E_r^{s-r+1,t,u+r}$ . Where  $E[-]$  denotes the exterior algebra and  $P[-]$  denotes the polynomial algebra on the indicated generators. This spectral sequence is referred as the small descent spectral sequence (SDSS).

(b) *There is a spectral sequence converging to  $\pi_*(S^0)$  with  $E_1$ -term*

$$E_1^{s,t} = \pi_*(X) \otimes E[\alpha_1] \otimes P[\beta_1], \quad \text{where} \\ E_1^{0,t} = \pi_t(X), \quad \alpha_1 \in E_1^{1,q}, \quad \beta_1 \in E_1^{2,p,q}$$

and  $d_r : E_r^{s,t} \longrightarrow E_r^{s+r,t+r-1}$ . This spectral sequence is referred as the topological small descent spectral sequence (TSDSS).

The above two spectral sequences produce the 0-line and the 1-line  $Ext_{BP_*BP}^{0,*}(BP_*, BP_*(S^0))$ ,  $Ext_{BP_*BP}^{1,*}(BP_*, BP_*(S^0))$  or the corresponding elements in  $\pi_*(S^0)$  by  $Ext_{BP_*BP}^{0,*}(BP_*, BP_*(X))$  and  $Ext_{BP_*BP}^{1,*}(BP_*, BP_*(X))$ .  $Ext_{BP_*BP}^{s,*}(BP_*, BP_*(S^0))$  ( $s \geq 2$ ) or the corresponding elements in  $\pi_*(S^0)$  is produced by  $Ext_{BP_*BP}^{s,*}(BP_*, BP_*(X))$  ( $s \geq 2$ ) as described as the following ABC Theorem.

**ABC Theorem** ([23] 7.5.1 ABC Theorem) *For  $p > 2$  and  $t - s < q(p^3 + p - 1) - 3$ ,  $s \geq 2$*

$$Ext_{BP_*BP}^{s,t}(BP_*, BP_*(X)) = Ext_{BP_*BP}^{s-2,*}(BP_*, BP_*(X) \otimes E_1^2) = A \oplus B \oplus C,$$

where  $A$  is the  $\mathbb{Z}/p$ -vector space spanned by

$$A = \{\beta_{ip}, \beta_{ip+1} | i \leq p-1\} \cup \{\beta_{p^2/p^2-j} | 0 \leq j \leq p-1\}, \\ B = R \otimes \{\gamma_i | i \geq 2\}$$

where

$$R = P[b_{20}^p] \otimes E[h_{20}] \otimes \mathbb{Z}/p \{ \{b_{11}^k | 0 \leq k \leq p-1\} \cup \{h_{11}b_{20}^k | 0 \leq k \leq p-2\} \},$$

and

$$C^{s,t} = \bigoplus_{i \geq 0} R^{s+2i, t+i(p^2-1)q}.$$

From the generators of  $R$  and [20] 4.11, 4.12 Theorem, we read the generators of  $C$  as follows:

Let  $i = jp + m$ . Then from  $R^{s+2i, t+i(p^2-1)q} \subset C^{s,t}$  we have:

- (1)  $b_{20}^{(j+1)p} \in R^{2(p-m)+2(jp+m), t+(jp+m)(p^2-1)q} \subset C^{2(p-m), t}$ , which is represented by

$$b_{20}^{p-m-1} u_{jp+m}$$

for  $p-1 \geq m \geq 1$ . From which we get

$$b_{20}^{p-m-1} u_{jp+m} \otimes E[h_{20}] \otimes \{b_{11}^k | 0 \leq k \leq p-1\} \cup \{h_{11} b_{20}^k | 0 \leq k \leq p-2\},$$

where

$$u_{jp+m} \in C^{2, q[(j+1)p^2 + (j+m+1)p+m]}.$$

- (2)  $b_{11}^k b_{20}^{jp} \in R^{2(k-m)+2(jp+m), t+(jp+m)(p^2-1)q} \subset C^{2(k-m), t}$ , which is represented by

$$b_{11}^{k-m-1} \beta_{(j+1)p/p-m}$$

for  $p-1 \geq k \geq m+1 \geq 1$ . From which we get

$$b_{11}^{k-m-1} \beta_{(j+1)p/p-m} \otimes E[h_{20}],$$

where

$$\beta_{(j+1)p/p-m} \in C^{2, q[(j+1)p^2 + jp+m]}.$$

- Especially  $h_{20} b_{11}^{p-1} b_{20}^{jp} \in R^{3+2(jp+p-2), t+(jp+p-2)(p^2-1)q} \subset C^{3, t}$  is represented by  $h_{11} \beta_{(j+1)p/1, 2}$ , which is an element of order  $p^2$ .

- (3)  $h_{11} b_{20}^k b_{20}^{jp} \in R^{2(k-m)+1+2(jp+m), t+(jp+m)(p^2-1)q} \subset C^{2(k-m)+1, t}$ , which is represented by

$$b_{20}^{k-m-1} \eta_{jp+m+1}$$

for  $p-2 \geq k \geq m+1 \geq 1$ , where

$$\eta_{jp+m+1} = h_{11} u_{jp+m} \in C^{3, q[(j+1)p^2 + (j+m+2)p+m]}.$$

- (4)  $h_{20} h_{11} b_{20}^k b_{20}^{jp} \in R^{2(k-m+1)+2(jp+m), t+(jp+m)(p^2-1)q} \subset C^{2(k-m+1), t}$ , which is represented by

$$b_{20}^{k-m} \beta_{jp+m+2}$$

for  $p-2 \geq k \geq m \geq 0$ , where

$$\beta_{jp+m+2} \in C^{2, q[jp^2 + (j+m+2)p+m+1]}.$$

- Especially  $h_{20} h_{11} b_{20}^{p-2} b_{20}^{jp} \in R^{2+2(jp+p-2), t+(jp+p-2)(p^2-1)q} \subset C^{2, t}$  is represented by  $\beta_{(j+1)p/1, 2}$ , which is an element of order  $p^2$ .

From the ABC Theorem above, we know that  $Ext_{BP_* BP}^{s, t}(BP_*, BP_*(X))$  for  $s \geq 2$ ,  $t-s < q(p^3 + p - 1) - 3$  is the  $\mathbb{Z}_{(p)}$ -module generated by the following generators, here the generators are listed as generators, total degree  $t-s$  and  $t-s \bmod pq-2$ , range of index; where  $pq-2 = 2p^2 - 2p - 2$  is the total degree of  $\beta_1 \in E_1^{0, qp, 2}$  in the SDSS.

Generators of A		$t - s$ and $t - s \bmod pq - 2$	Range of index
$\beta_{ip}$		$q[ip^2 + ip - 1] - 2$	
	$\equiv$	$2(i - 1)p + 2i$	if $i \leq p - 2$
	$\equiv$	0	if $i = p - 1$
$\beta_{ip+1}$		$q[ip^2 + (i + 1)p] - 2$	
	$\equiv$	$2ip + 2i$	if $i \leq p - 2$
	$\equiv$	$\underline{2p}_{2p}$	if $i = p - 1$
$\beta_{p^2/p^2-j}$		$q[p^3 + j] - 2$	
	$\equiv$	$2(j + 1)p - 2j_{2p}$	if $j \leq p - 2$
	$\equiv$	4	if $j = p - 1$
Generators of B		$t - s$ and $t - s \bmod pq - 2$	Range of index
$h_{11}b_{20}^k\gamma_i$		$q[(i + k)p^2 + (i + k)p + i - 2] - 2k - 4$	for $2 \leq i \leq p - 1, 0 \leq k \leq p - 2, 2 \leq i + k \leq p - 1$
	$\equiv$	$2(k + 2i - 2)p$	if $k + 2i \leq p$
	$\equiv$	$2(k + 2i - p - 1)p + 2$	if $k + 2i > p$
$h_{20}h_{11}b_{2,0}^k\gamma_i$		$q[(i + k)p^2 + (i + k + 1)p + i - 1] - 2k - 5$	for $2 \leq i \leq p - 1, 0 \leq k \leq p - 2, 2 \leq i + k \leq p - 1$
	$\equiv$	$2(k + 2i - 1)p - 1$	if $k + 2i < p$
	$\equiv$	$2(k + 2i - p)p + 1$	if $k + 2i \geq p$
$b_{11}^k\gamma_i$		$q[(i + k)p^2 + (i - 1)p + i - 2] - 2k - 3$	for $2 \leq i \leq p - 1, 0 \leq k \leq p - 1, 2 \leq i + k \leq p - 1$
	$\equiv$	$2(k + 2i - 2)p - 2k - 1$	if $k + 2i \leq p + 1$
	$\equiv$	1	if $k = 0, 2i = p + 1$
$h_{20}b_{11}^k\gamma_i$		$\equiv \underline{2(k + 2i - p - 1)p - 2k + 1}_{4p-3}$	if $k + 2i \geq p + 2$
		$q[(i + k)p^2 + ip + i - 1] - 2k - 4$	for $2 \leq i \leq p - 1, 0 \leq k \leq p - 1, 2 \leq i + k \leq p - 1$
	$\equiv$	$2(k + 2i - 1)p - 2(k + 1)$	if $k + 2i \leq p$
		$\equiv \underline{2(k + 2i - p)p - 2k}_{2p}$	if $k + 2i > p$

Generators of C	$t-s$ and $t-s \bmod pq-2$	Range of index
$b_{11}^k b_{20}^{p-m-1} u_{jp+m}$	$q[(p-m+j+k+1)p^2+jp+m]-2(p-m+k)$ $\equiv 2(j+k+1)p+2(j-k+1)$	for $1 \leq m < p, 0 \leq j \leq p-2, 0 \leq k < p$ and $j+k < m$
$h_{20} b_{11}^k b_{2,0}^{p-m-1} u_{jp+m}$	$q[(p-m+j+k+1)p^2+(j+1)p+m+1]-2(p-m+k)-1$ $\equiv 2(j+k+2)p+2(j-k+1)-1$ $\equiv 2(j-k+2)p-1$	for $1 \leq m < p, 0 \leq j \leq p-2, 0 \leq k < p$ and $j+k < m, j+k \leq p-3$ if $j+k \leq p-4$ or $j+k = p-3, 2j < p-5$ if $j+k = p-3, 2j \geq p-5$
$h_{11} b_{20}^{k+p-m-1} u_{jp+m}$	$q[(p-m+j+k+1)p^2+(j+k+1)p+m]-2(p-m+k)-1$ $\equiv 2(j+k+2)p+2(j-p)+3$	for $1 \leq m < p, 0 \leq j \leq p-2, 0 \leq k \leq p-2$ and $j+k < m, j+k \leq p-3$
$h_{20} h_{11} b_{2,0}^{k+p-m-1} u_{jp+m}$	$q[(p-m+j+k+1)p^2+(j+k+2)p+m]+2(m-k-2)$ $\equiv 2(j+k+2)p+2j+2$ $\equiv 2j+4$	for $1 \leq m < p, 0 \leq j \leq p-2, 0 \leq k \leq p-2$ and $j+k < m, j+k \leq p-3$ if $j+k \leq p-4$ if $j+k = p-3$
$b_{11}^{k-m-1} \beta_{(j+1)p/p-m}$	$q[(j+k-m)p^2+jp+m]-(2k-2m)$ $\equiv 2(j+k)p+2(j-k)$ $\equiv \frac{2(j+k-p+1)p+2(j-k+1)}{2p}$	for $1 \leq m+1 \leq k < p, 0 \leq j \leq p-2$ if $j+k \leq p-2$ or $j+k = p-1, 2j < p-1$ if $j+k \geq p$ or $j+k = p-1, 2j \geq p-1$
$h_{20} b_{11}^{k-m-1} \beta_{(j+1)p/p-m}$	$q[(j+k-m)p^2+(j+1)p+m+1]-(2k-2m+1)$ $\equiv \frac{2(j+k+1)p+2(j-k)-1}{4p-3}$ $\equiv 2(j+k-p+2)p+2(j-k)+1$	for $1 \leq m+1 \leq k < p, 0 \leq j \leq p-2$ if $j+k \leq p-3$ or $j+k = p-2, 2j \leq p-3$ if $j+k > p-2$ or $j+k = p-2, 2j > p-3$

continue to next page. . .

Generators of C	$t-s$ and $t-s \bmod pq-2$	Range of index
$h_{1,1}\beta_{(j+1)p/1,2}$	$q[(j+1)p^2 + (j+2)p - 1] - 3$ $\equiv 2jp + 2(j+1) + 1$ $\equiv 1$	for $0 \leq j \leq p-2$ if $j \leq p-3$ if $j = p-2$
$b_{2,0}^{k-m-1}\eta_{jp+m+1}$	$q[(j+k-m)p^2 + (j+k+1)p + m] - (2k-2m+1)$ $\equiv 2(j+k)p + 2j + 1$ $\equiv \frac{2(j+k-p+2)p + 2(j-p) + 3}{4p-3}$	for $1 \leq m+1 \leq k \leq p-2, 0 \leq j \leq p-2$ if $j+k \leq p-2$ if $j+k > p-2$
$b_{2,0}^{k-m}\beta_{jp+m+2}$	$q[(j+k-m)p^2 + (j+k+2)p + m+1] - 2(k-m+1)$ $\equiv \frac{2(j+k+1)p + 2j_{2p}}{4p-3}$ $\equiv 2(j+k-p+3)p + 2(j-p) + 2$ $\equiv 0$	for $0 \leq m \leq k \leq p-2, 0 \leq j \leq p-2$ if $j+k \leq p-3$ if $j+k > p-3$ if $j=k=p-2$
$\beta_{(j+1)p/1,2}$	$q[(j+1)p^2 + (j+1)p - 1] - 2$ $\equiv 2jp + 2(j+1)$ $\equiv 0$	for $0 \leq j \leq p-2$ if $j \leq p-3$ if $j = p-2$



Let us take  $h_{11}b_{20}^k\gamma_i$  from the  $B$ -family as an example to illustrate the calculation.  
The total degree of  $h_{11}b_{20}^k\gamma_i$  is

$$q[(i+k)p^2 + (i+k)p + i - 2] - 2k - 4 = 2(i+k)p^3 - 2(k+2)p - 2(i+k)$$

for  $2 \leq i \leq p-1, 0 \leq k \leq p-2$ . To ensure that the total degree of  $h_{11}b_{20}^k\gamma_i$  is less than  $q(p^3 + p - 1) - 3$ , we need  $i+k < p$ . Straightforward computation shows

$$2(i+k)p^3 - 2(k+2)p - 2(i+k) \equiv 2(k+2i-2)p \pmod{pq-2}$$

Notice that  $2(k+2i-2)p > pq-2$  if  $k+2i > p$ , the total degree of  $h_{11}b_{20}^k\gamma_i$  is

$$2(k+2i-2)p - (pq-2) = 2(k+2i-p-1)p + 2 \pmod{pq-2}$$

if  $k+2i > p$ .

**Remark.** The Adams-Novikov spectral sequence for the spectrum  $X$  collapses from  $E_2$ -term  $Ext_{BP_*BP}^{s,t}(BP_*, BP_*(X))$  in the range  $t-s < q(p^3 + p - 1) - 3$ , since there are no elements with filtration  $> 2p$ . Thus we actually get the homotopy groups  $\pi_{t-s}(X)$  in this range.

### 3. THE ANSS $E_2$ -TERM $Ext_{BP_*BP}^{s,t}(BP_*, BP_*)$ AT $t-s = q(p^3 + 1) - 3$

Consider the Adams-Novikov differential  $d_r : E_r^{s,t} \rightarrow E_r^{s+r, t+r-1}$  in the ANSS. From the total degree of  $\beta_{p^2/p^2-1}$ , we know that  $d_r(\beta_{p^2/p^2-1}) \in Ext_{BP_*BP}^{s,t}(BP_*, BP_*)$  such that  $t-s = q(p^3 + 1) - 3$ . The SDSS  $E_1^{s,t,u}$  converges to  $Ext_{BP_*BP}^{s+u,t}(BP_*, BP_*)$ . Fix  $t-s-u = q(p^3 + 1) - 3$ , we have:

**Lemma 3.1.** Fix  $t-s-u = q(p^3 + 1) - 3$ , the  $E_1$ -term  $E_1^{s,t,u}$  of the SDSS is the  $\mathbb{Z}/p$ -module generated by the following 8 generators:

$$\begin{aligned} \mathfrak{g}_1 &= \alpha_1 \beta_1^{p^2-1} \beta_2 \in E_1^{2,*,2p^2-1}; & \mathfrak{g}_2 &= \beta_1^{p^2-p} h_{20} \beta_{p/p} \in E_1^{3,*,2p^2-2p}; \\ \mathfrak{g}_3 &= \alpha_1 \beta_1^{\frac{p^2-2p-1}{2}} h_{2,0} \gamma_{\frac{p+1}{2}} \in E_1^{4,*,p^2-2p}; & \mathfrak{g}_4 &= \beta_1^{\frac{p^2-6p+1}{2}} b_{11}^2 \gamma_{\frac{p+1}{2}} \in E_1^{7,*,p^2-6p+1}; \\ \mathfrak{g}_5 &= \alpha_1 \beta_1^{mp-\frac{p-1}{2}} b_{11}^{\frac{p-1}{2}-m} \beta_{(\frac{p+1}{2})p/p-m} \in E_1^{p+1-2m,*,*}; & \mathfrak{g}_6 &= \beta_1^{p-1} \eta_{(p-3)p+3} \in E_1^{3,*,2p-2}; \\ \mathfrak{g}_7 &= \alpha_1 \beta_{(p-1)p+1} \in E_1^{2,q(p^3+1),1}; & \mathfrak{g}_8 &= \alpha_1 \beta_{p^2/p^2} \in E_1^{2,q(p^3+1),1}. \end{aligned}$$

*Proof.* Fix  $t-s-u = q(p^3 + 1) - 3$ . From the ABC Theorem, we know that the generators of the  $E_1$ -terms in the SDSS are of the form  $W = \beta_1^k w$  or  $W = \alpha_1 \beta_1^k w$ , where  $w$  is an element listed in the ABC Theorem.

1. If a generator of  $E_1^{s,t,u}$  is of the form  $W = \beta_1^k w$ , then the total degree of  $\beta_1^k w$  is  $q(p^3 + 1) - 3$  and the total degree of  $w$  is  $q(p^3 + 1) - 3$  modulo the total degree of  $\beta_1$  which is  $t-u = qp-2$ . Note that

$$q(p^3 + 1) - 3 \equiv 4p - 3 \pmod{qp-2},$$

we list all the generators whose total degree might be  $4p-3 \pmod{qp-2}$ , which are marked with underline and subscript  $4p-3$  in the table for ABC Theorem.

$$\begin{aligned} b_{11}^k \gamma_i & \text{ at } k=2 \text{ and } i=(p+1)/2; \\ h_{20} b_{11}^{k-m-1} \beta_{(j+1)p/p-m} & \text{ at } k=1 \text{ and } j=0; \\ b_{20}^{k-m-1} \eta_{jp+m+1} & \text{ at } k=3 \text{ and } j=p-3. \end{aligned}$$

From which we get the following generators in  $E_1^{s,t,u}$ :

$$\begin{aligned} b_{11}^2 \gamma_{\frac{p+1}{2}} &\implies \mathfrak{g}_4 = \beta_1^{\frac{p^2-6p+1}{2}} b_{11}^2 \gamma_{\frac{p+1}{2}} \in E_1^{7,*,p^2-6p+1}; \\ h_{20} \beta_{p/p} &\implies \mathfrak{g}_2 = \beta_1^{p^2-p} h_{20} \beta_{p/p} \in E_1^{3,*,2p^2-2p}; \\ \eta_{(p-3)p+3} &\implies \mathfrak{g}_6 = \beta_1^{p-1} \eta_{(p-3)p+3} \in E_1^{3,*,2p-2}. \end{aligned}$$

**2.** If a generator of  $E_1^{s,t,u}$  is of the form  $W = \alpha_1 \beta_1^k w_1$ , then from the total degree of  $\alpha_1$  being  $t - u = 2p - 3$  we see that the total degree of  $w_1$  is  $2p$  modulo  $qp - 2$ . Similarly we can find all such  $w_1$ 's, which are marked with underline and subscript  $2p$  in the ABC Theorem:

$$\beta_{(p-1)p+1}; \quad \beta_{p^2/p^2}; \quad h_{20} \gamma_{\frac{p+1}{2}}; \quad b_{11}^{\frac{p-1}{2}-m} \beta_{(\frac{p+1}{2})p/p-m}; \quad \beta_2.$$

From which we get the following generators in  $E_1^{s,t,u}$ :

$$\begin{aligned} \mathfrak{g}_7 &= \alpha_1 \beta_{(p-1)p+1}; & \mathfrak{g}_8 &= \alpha_1 \beta_{p^2/p^2}; \\ \mathfrak{g}_3 &= \alpha_1 \beta_1^{\frac{p^2-2p-1}{2}} h_{2,0} \gamma_{\frac{p+1}{2}}; & \mathfrak{g}_5 &= \alpha_1 \beta_1^{mp-\frac{p-1}{2}} b_{11}^{\frac{p-1}{2}-m} \beta_{(\frac{p+1}{2})p/p-m}; \\ \mathfrak{g}_1 &= \alpha_1 \beta_1^{p^2-1} \beta_2. \end{aligned}$$

Computing the filtration of the corresponding generators, we get the lemma.  $\square$

**Remark:** The method in proving Lemma 3.1 is a general method in computing the  $E_1$ -term  $E_1^{s,t,u}$  of the SDSS with specialized  $t - s - u$ .

**Theorem 3.2.** Fix  $t - s = q(p^3 + 1) - 3$ , the Adams-Novikov  $E_2$ -term  $Ext_{BP_*BP}^{s,t}(BP_*, BP_*)$  is the  $\mathbb{Z}/p$ -module generated by the following 6 elements

$$\begin{aligned} \mathfrak{g}_1 &= \alpha_1 \beta_1^{p^2-1} \beta_2 \in Ext_{BP_*BP}^{2p^2+1,*}; \\ \mathfrak{g}_3 &= \alpha_1 \beta_1^{\frac{p^2-2p-1}{2}} h_{2,0} \gamma_{\frac{p+1}{2}} \in Ext_{BP_*BP}^{p^2-2p+4,*}; & \mathfrak{g}_4 &= \beta_1^{\frac{p^2-6p+1}{2}} b_{11}^2 \gamma_{\frac{p+1}{2}} \in Ext_{BP_*BP}^{p^2-6p+8,*}; \\ & & \mathfrak{g}_6 &= \beta_1^{p-1} \eta_{(p-3)p+3} \in Ext_{BP_*BP}^{2p+1,*}; \\ \mathfrak{g}_7 &= \alpha_1 \beta_{(p-1)p+1} \in Ext_{BP_*BP}^{3,q(p^3+1)}; & \mathfrak{g}_8 &= \alpha_1 \beta_{p^2/p^2} \in Ext_{BP_*BP}^{3,q(p^3+1)}. \end{aligned}$$

*Proof.* Following D. Ravenel [22] page 287, we compute in the cobar complex of  $N_0^2 = BP_*/(p^\infty, v_1^\infty)$

$$\begin{aligned} d \left( \frac{v_2^{jp}}{pv_1^p} (t_2 - t_1^{p+1}) \right) &= \frac{v_2^{jp}}{pv_1^p} t_1^p \otimes t_1 + \frac{v_2^{jp}}{pv_1^{p-1}} b_{10}, \\ -d \left( \frac{v_2^{jp+1}}{pv_1^{p+1}} t_1 \right) &= -\frac{v_2^{jp}}{pv_1^p} t_1^p \otimes t_1 - j \frac{v_2^{(j-1)p+1}}{pv_1} t_1^{p^2} \otimes t_1 + \frac{v_2^{jp}}{pv_1} t_1 \otimes t_1, \\ d \left( j \frac{v_2^{(j-1)p} v_3}{pv_1} t_1 \right) &= j \frac{v_2^{(j-1)p+1}}{pv_1} t_1^{p^2} \otimes t_1 - j \frac{v_2^{jp}}{pv_1} t_1 \otimes t_1, \\ -(j-1)/2d \left( \frac{v_2^{jp}}{pv_1} t_1^2 \right) &= (j-1) \frac{v_2^{jp}}{pv_1} t_1 \otimes t_1. \end{aligned}$$

Straightforward calculation shows that the coboundary of

$$\frac{v_2^{jp}}{pv_1^p} t_2 - \frac{v_2^{jp}}{pv_1^p} t_1^{p+1} - \frac{v_2^{jp+1}}{pv_1^{p+1}} t_1 + j \frac{v_2^{(j-1)p} v_3}{pv_1} t_1 - (j-1)/2 \frac{v_2^{jp}}{pv_1} t_1^2$$

is  $\frac{v_2^{jp}}{pv_1^{p-1}}b_{10}$ . Then from  $\delta\delta\left(\frac{v_2^{jp}}{pv_1^p}\right) = \beta_{jp/p}$ , we get a differential in the SDSS


$$d_2(h_{20}\beta_{jp/p}) = \beta_1\beta_{jp/p-1}.$$

Similarly, we have

$$(3.1) \quad d_2(h_{20}\beta_{jp/i}) = \beta_1\beta_{jp/i-1} \quad \text{for } 2 \leq i \leq p.$$

Applying formula (3.1), we get the following differentials in the SDSS

$$\begin{aligned} d_2(\mathfrak{g}_2) &= d_2(\beta_1^{p^2-p}h_{20}\beta_{p/p}) = \beta_1^{p^2-p+1}\beta_{p/p-1}, \\ d_2(\alpha_1\beta_1^{mp-\frac{p-1}{2}-1}b_{11}^{\frac{p-1}{2}-m}h_{20}\beta_{(\frac{p+1}{2})p/p-m+1}) &= \alpha_1\beta_1^{mp-\frac{p-1}{2}}b_{11}^{\frac{p-1}{2}-m}\beta_{(\frac{p+1}{2})p/p-m} = \mathfrak{g}_5 \end{aligned}$$



The theorem follows.  $\square$

#### 4. A DIFFERENTIAL IN THE ANSS

This section is armed at showing that

$$(4.1) \quad d_{2p-1}(h_{20}b_{11}\gamma_s) = \alpha_1\beta_1^p h_{20}\gamma_s$$

in the ANSS, which will be used in proving Theorem A in section 5.

We begin from showing that  $\pi_{q(p^2+2p+2)-2}(V(2)) = 0$ . From which we show that the Toda bracket  $\langle \alpha_1\beta_1, p, \gamma_s \rangle = 0$  and the Toda bracket  $\langle \alpha_1\beta_1^{p-1}, \alpha_1\beta_1, p, \gamma_s \rangle$  is well defined. Then from the relation

$$\langle \alpha_1\beta_1^{p-1}, \alpha_1\beta_1, p, \gamma_s \rangle = \alpha_1\beta_1^{p-1}h_{20}\gamma_s = \beta_{p/p-1}\gamma_s$$

in  $\pi_*(S^0)$  and  $d(h_{20}b_{11}) = \beta_1\beta_{p/p-1}$ , we get the desired differential in the ANSS.

Let  $p \geq 5$  be an odd prime and  $V(2)$  be the Smith-Toda spectrum characterized by

$$BP_*(V(2)) = BP_*/I_3$$

where  $I_3$  is the invariant ideal of  $BP_* = \mathbb{Z}_{(p)}[v_1, v_2, \dots, v_i, \dots]$  generated by  $p, v_1$  and  $v_2$ . To compute the homotopy groups of  $V(2)$ , one has the ANSS  $\{E_r^{s,t}V(2), d_r\}$  that converges to  $\pi_*(V(2))$ . The  $E_2$ -page of this spectral sequence is

$$E_2^{s,t}V(2) = Ext_{BP_*BP}^{s,t}(BP_*, BP_*(V(2)))$$

Let

$$\Gamma = BP_*/I_3 \otimes_{BP_*} BP_*BP \otimes_{BP_*} BP_*/I_3 = BP_*/I_3[t_1, t_2, \dots].$$

Then  $(BP_*/I_3, \Gamma)$  is a Hopf algebroid, and its structure map is deduced from that of  $(BP_*, BP_*(BP))$ . By a change of ring theorem, one sees that

$$Ext_{BP_*BP}^{s,t}(BP_*, BP_*(V(2))) = Ext_{\Gamma}^{s,t}(BP_*, BP_*/I_3) \implies \pi_*(V(2))$$

**Lemma 4.1.** *The  $q(p^2 + 2p + 2) - 2$  dimensional stable homology group of  $V(2)$  is trivial, i.e.,*

$$\pi_{q(p^2+2p+2)-2}(V(2)) = 0.$$

*Proof.* Fix  $t - s = q(p^2 + 2p + 2) - 2$ , we know that the Adams-Novikov  $E_2$ -term

$$Ext_{BP_*BP}^{s, s+q(p^2+2p+2)-2}(BP_*, BP_*(V(2))) = Ext_{\Gamma}^{s, s+q(p^2+2p+2)-2}(BP_*, BP_*/I_3)$$

converges to  $\pi_{q(p^2+2p+2)-2}(V(2))$ . We will prove that  $\pi_{q(p^2+2p+2)-2}(V(2)) = 0$  by showing that  $Ext_{BP_*BP}^{s, s+q(p^2+2p+2)-2}(BP_*, BP_*(V(2))) = 0$ .

In the cobar complex  $C_{\Gamma}^s BP_*/I_3$ , the inner degree of  $v_i$ ,  $|v_i| = |t_i| \geq q(p^3 + p^2 + p + 1)$  for  $i \geq 4$ . It follows that in the range  $t - s \leq q(p^3 + p^2 + p + 1) - 1$ ,

$$Ext_{BP_*BP}^{s,t}(BP_*, BP_*/I_3) = Ext_{\Gamma}^{s,t}(BP_*, BP_*/I_3) = Ext_{\Gamma'}^{s,t}(BP_*, BP_*/I_3).$$

where  $\Gamma' = \mathbb{Z}/p[v_3][t_1, t_2, t_3]$ . From  $\eta_R(v_3) \equiv v_3 \pmod{I_3}$ , we see that

$$Ext_{\mathbb{Z}/p[v_3][t_1, t_2, t_3]}^{s,*}(BP_*, BP_*/I_3) \cong Ext_{\mathbb{Z}/p[t_1, t_2, t_3]}^{s,*}(\mathbb{Z}/p, \mathbb{Z}/p) \otimes \mathbb{Z}/p[v_3].$$

To compute the  $Ext$  groups  $Ext_{\mathbb{Z}/p[t_1, t_2, t_3]}^*(\mathbb{Z}/p, \mathbb{Z}/p)$ , we can use the modified May spectral sequence (MSS) introduced in [7, 8, 9, 23].

There is the May spectral sequence  $\{E_r^{s,t,*}, \delta_r\}$  that converges to  $Ext_{\mathbb{Z}/p[t_1, t_2, t_3]}^{s,t}(\mathbb{Z}/p, \mathbb{Z}/p)$ . The  $E_1$ -term of this spectral sequence is

$$(4.2) \quad E_1^{*,*,*} = E[h_{ij} | 0 \leq j, i = 1, 2, 3] \otimes P[b_{ij} | 0 \leq j, i = 1, 2, 3],$$

where

$$h_{ij} \in E_1^{1, q(1+p+\dots+p^{i-1})p^j, 2i-1} \quad \text{and} \quad b_{ij} \in E_1^{2, q(1+p+\dots+p^{i-1})p^{j+1}, p(2i-1)}.$$

The first May differential is given by

$$(4.3) \quad \delta_1(h_{i,j}) = \sum_{0 < k < i} h_{i-k, k+j} h_{k,j} \quad \text{and} \quad \delta_1(b_{i,j}) = 0.$$

For the reason of the total degree, to compute  $Ext_{BP_*BP}^{s, s+(q(p^2+2p+2)-2)}(BP_*, BP_*/I_3)$  we only need to consider the sub-module generated by  $h_{30}, h_{20}, h_{10}, h_{21}, h_{11}, h_{12}$  and  $b_{20}, b_{10}, b_{11}$ , i. e. the subcomplex

$$E[h_{ij} | 1 \leq i, i+j \leq 3] \otimes E[b_{20}, b_{11}] \otimes P[b_{10}].$$

From (4.3), we know that within  $t - s \leq q(p^2 + 2p + 2) - 2$  the May's  $E_2$ -term

$$E_2^{s,*,*} = H^{s,*,*}(E_1^{s,*,*}, \delta_1) = H^{s,*,*}(E[h_{ij} | 0 \leq j, i+j \leq 3], \delta_1) \otimes E[b_{20}, b_{11}] \otimes P[b_{10}].$$

H. Toda in [28] computed the cohomology of  $(E[h_{ij} | 0 \leq j, i+j \leq 3], \delta_1)$ . Here we only jot down the even dimensional elements within that range.

$$\begin{array}{ll} h_{20}h_{10}, & q(p+2)-2; & h_{20}h_{11}, & q(2p+1)-2; \\ h_{12}h_{10}, & q(p^2+1)-2; & h_{21}h_{11}, & q(p^2+2p)-2. \end{array}$$

Thus within  $t - s \leq q(p^2 + 2p + 2) - 2$ , the even dimensional May's  $E_2$ -term  $E_2^{s,t,*}$  is a submodule of

$$\mathbb{Z}/p\{1, h_{20}h_{10}, h_{20}h_{11}, h_{12}h_{10}, h_{21}h_{11}\} \otimes E[b_{20}, b_{11}] \otimes P[b_{10}].$$

Suppose we have a generator  $y$  in  $Ext_{\mathbb{Z}/p[v_3][t_1, t_2, t_3]}^{s, s+q(p^2+2p+2)-2}(BP_*, BP_*/I_3)$ . Then  $y$  is the form of  $x$  or  $v_3x$  where  $x$  is an even dimensional generator in  $H^*(E[h_{ij}|i+j \leq 3]) \otimes E[b_{20}, b_{11}] \otimes P[b_{10}]$ .

- (1) If  $y = v_3x$ , then  $x \in E_2^{s, t, *}$  subject to  $t - s = q(p+1) - 2$ . An easy computation shows that the corresponding  $E_2$ -term is zero.
- (2) If  $y = x$ , then  $x \in E_2^{s, t, *}$  subject to  $t - s = q(p^2 + 2p + 2) - 2$ . Similarly, from

$$q(p^2 + 2p + 2) - 2 \equiv 6p - 2 \pmod{qp - 2}$$

we compute that the total degree  $t - s \pmod{qp - 2}$  of the generators in

$$\mathbb{Z}/p\{1, h_{20}h_{10}, h_{20}h_{11}, h_{12}h_{10}, h_{21}h_{11}\} \otimes [b_{20}, b_{11}]$$

and find none of them is  $6p - 2$ . Thus the corresponding  $E_2$ -term is zero.

The Lemma follows.  $\square$

It is easily showed that the following theorem holds from the lemma above.

**Theorem 4.2.** *For  $p \geq 5$ ,  $s \geq 1$ , the Toda bracket  $\langle \alpha_1\beta_1, p, \gamma_s \rangle = 0$ .*

*Proof.* Let  $\tilde{v}_3$  be the composition of the following maps

$$S^{q(p^2+p+1)} \xrightarrow{\tilde{i}} \Sigma^{q(p^2+p+1)}V(2) \xrightarrow{v_3} V(2),$$

where the first map is inclusion to the bottom cell.

It is known that  $\tilde{v}_3$  is an order  $p$  element in  $\pi_{q(p^2+p+1)}(V(2))$ . Thus the Toda bracket  $\langle \alpha_1\beta_1, p, \tilde{v}_3 \rangle$  is well defined and  $\langle \alpha_1\beta_1, p, \tilde{v}_3 \rangle \in \pi_{q(p^2+2p+2)-2}(V(2)) = 0$ . It follows that the Toda bracket  $\langle \alpha_1\beta_1, p, \tilde{v}_3 \rangle = 0$ .

Let  $\tilde{j} : V(2) \rightarrow S^{q(p+2)+3}$  be the collapsing lower cells map from  $V(2)$ , then  $\gamma_s = \tilde{v}_3 \cdot v_3^{s-1} \cdot \tilde{j}$ . As a result,

$$\langle \alpha_1\beta_1, p, \gamma_s \rangle = \langle \alpha_1\beta_1, p, \tilde{v}_3 \cdot v_3^{s-1} \cdot \tilde{j} \rangle = \langle \alpha_1\beta_1, p, \tilde{v}_3 \rangle \cdot v_3^{s-1} \cdot \tilde{j} = 0$$

because  $\langle \alpha_1\beta_1, p, \tilde{v}_3 \rangle = 0 \in \pi_{q(p^2+2p+2)-2}V(2) = 0$ .  $\square$

**Proposition 4.3.** *(see also [22] 7.5.11) Let  $p \geq 5$  be an odd prime. Then in  $\pi_*(S^0)$ , the Toda bracket  $\langle \alpha_1\beta_1^{p-1}, \alpha_1\beta_1, p, \gamma_s \rangle$  is well defined and*

$$\alpha_1\beta_1^{p-1}h_{20}\gamma_s = \langle \alpha_1\beta_1^{p-1}, \alpha_1\beta_1, p, \gamma_s \rangle = \beta_{p/p-1}\gamma_s.$$

*Proof.* From  $\langle \beta_1^{p-1}, \alpha_1\beta_1, p \rangle = 0$ ,  $\langle \alpha_1\beta_1, p, \alpha_1 \rangle = 0$ ,  $\langle \alpha_1, \alpha_1\beta_1, p \rangle = 0$  and  $\langle \alpha_1\beta_1, p, \gamma_s \rangle = 0$ , we know that the following 4-fold Toda bracket is well defined and

$$\beta_{p/p-1} = \langle \beta_1^{p-1}, \alpha_1\beta_1, p, \alpha_1 \rangle; \quad \alpha_1h_{20}\gamma_s = \langle \alpha_1, \alpha_1\beta_1, p, \gamma_s \rangle.$$

On the other hand, one has

$$\begin{aligned} \beta_1^{p-1}\alpha_1h_{20}\gamma_s &= \beta_1^{p-1}\langle \alpha_1, \alpha_1\beta_1, p, \gamma_s \rangle \\ &= \langle \alpha_1\beta_1^{p-1}, \alpha_1\beta_1, p, \gamma_s \rangle \\ &= \alpha_1\langle \beta_1^{p-1}, \alpha_1\beta_1, p, \gamma_s \rangle \\ &= \langle \beta_1^{p-1}, \alpha_1\beta_1, p, \alpha_1\gamma_s \rangle \\ &= \langle \beta_1^{p-1}, \alpha_1\beta_1, p, \alpha_1 \rangle \cdot \gamma_s \\ &= \beta_{p/p-1}\gamma_s \end{aligned}$$

The proposition follows.  $\square$

**Theorem 4.4.** *Let  $p \geq 5$  be an odd prime and  $2 \leq s \leq p-2$ . Then in the ANSS, we have the following Adams-Novikov differential*

$$d_{2p-1}(h_{2,0}b_{1,1}\gamma_s) = \alpha_1\beta_1^p h_{2,0}\gamma_s.$$

*Proof.* Note that  $b_{11} = \beta_{p/p}$ . Then from (3.1) one has the differential in the small descent spectral sequence

$$d_2(h_{20}b_{11}) = \beta_1\beta_{p/p-1},$$

which could be read as  $d(h_{20}\beta_{p/p}) = \beta_1\beta_{p/p-1}$  and  $d(h_{20}\beta_{p/p}\gamma_s) = \beta_1\beta_{p/p-1}\gamma_s$  in the cobar complex of  $BP_*$  or equivalently the first Adams-Novikov differential in the ANSS. Then from the relation  $\beta_{p/p-1}\gamma_s = \alpha_1\beta_1^{p-1}h_{20}\gamma_s$  in  $\pi_*(S^0)$  and  $\beta_{p/p-1}\gamma_s = 0$  in  $Ext_{BP_*BP}^{5,*}(BP_*, BP_*)$ , we get the Adams differential in the ANSS

$$d_{2p-1}(h_{2,0}b_{1,1}\gamma_s) = \beta_1 \cdot \beta_1^{p-1} \alpha_1 h_{20}\gamma_s = \alpha_1 \beta_1^p h_{20}\gamma_s.$$

The theorem follows.  $\square$

## 5. THE PROOF OF THEOREM A

In this section we prove our main theorem by showing that  $\beta_{p^2/p^2-1}$  survives to  $E_\infty$  in the ANSS. Note that  $\beta_{p^2/p^2-1}$  has too low dimension to be the target of an Adams-Novikov differential, we will do this by showing that all the Adams-Novikov differentials  $d_r(\beta_{p^2/p^2-1})$  are trivial.

**Lemma 5.1.** *Let  $p \geq 5$  and  $i \not\equiv 0 \pmod p$ . In the ANSS, one has the following Adams-Novikov differential*

$$d_{2p-1}(\eta_i) = \beta_1^p \beta_{i+1}$$

*Proof.* Recall from [22] 7.3.11 Theorem (e), in the SDSS

$$E_1 = Ext_{BP_*BP}^{s,t}(BP_*, BP_*(X^{p^2-1})) \otimes E[h_{11}] \otimes P[b_{11}] \implies Ext_{BP_*BP}^{s,t}(BP_*, BP_*(X)),$$

where  $BP_*(X^{p^2-1}) = BP_*[t_1]/\langle t_1^{p^2} \rangle$  (cf. [22] 7.3.8 Theorem), one has  $d_2(h_{20}\mu_{i-1}) = ib_{11}\beta_{i+1}$ . And from its definition we know that  $\eta_i = h_{11}\mu_{i-1}$  is represented by

$$\delta\delta \left( \frac{v_2^{p+i-1}t_2 + v_2^i t_2^p - v_2^i t_1^{p^2+p} - v_2^{i-1}v_3 t_1^p}{pv_1} \right)$$

(cf. [22] p.288) which is also denoted by  $\delta\delta \left( \frac{v_2^{p+i}}{pv_1} \zeta_2 \right)$  in [10, 29]. In the cobar complex of  $N_0^2 = BP_*/(p^\infty, v_1^\infty)$ , a straightforward computation shows that the coboundary of

$$\frac{v_2^i(t_3 - t_1 t_2^p - t_2 t_1^{p^2} + t_1^{p^2+p+1}) + v_2^{p+i-1}(t_1 t_2 - t_1^{p+2}) - v_2^{i-1}v_3(t_2 - t_1^{p+1})}{pv_1}$$

$$+ \frac{2v_2^{p+i}}{(p+i)p^2 v_1} t_1 - \frac{v_2^{p+i}}{(p+i)pv_1^2} t_1^2$$

is  $\frac{(v_2^{p+i-1}t_2 + v_2^i t_2^p - v_2^i t_1^{p^2+p} - v_2^{i-1}v_3 t_1^p) \otimes t_1}{pv_1} + \frac{v_2^{i+1}}{pv_1} b_{11}$ . This shows that in  $Ext_{BP_*BP}^{2,*}(BP_*, N_0^2)$  the cohomology class

$$\left[ \frac{(v_2^{p+i-1}t_2 + v_2^i t_2^p - v_2^i t_1^{p^2+p} - v_2^{i-1}v_3 t_1^p) \otimes t_1}{pv_1} \right] = - \left[ \frac{v_2^{i+1}}{pv_1} b_{11} \right].$$

Applying the connecting homomorphism  $\delta\delta$ , we get  $\alpha_1\eta_i = \beta_{i+1}\beta_{p/p}$ .

From  $\alpha_1 \eta_i = \beta_{i+1} \beta_{p/p}$  and the Toda differential, one has:

$$\alpha_1 d_{2p-1}(\eta_i) = d_{2p-1}(\alpha_1 \eta_i) = d_{2p-1}(\beta_{i+1} \beta_{p/p}) = \alpha_1 \beta_1^p \beta_{i+1}$$

The lemma follows from  $\alpha_1 d_{2p-1}(\eta_i) = \alpha_1 \beta_1^p \beta_{i+1}$ .  $\square$

**Proof of Theorem A** From  $\beta_{p^2/p^2-1} \in Ext_{BP_*BP}^{2,q(p^3+1)}(BP_*, BP_*)$ , we know that  $d_r(\beta_{p^2/p^2-1}) \in Ext_{BP_*BP}^{s,t}(BP_*, BP_*)$  subject to  $t - s = q(p^3 + 1) - 3$ . From Theorem 3.2 we know that the corresponding  $Ext_{BP_*BP}^{s,t}(BP_*, BP_*)$  is the  $\mathbb{Z}/p$ -module generated by  $\mathfrak{g}_1, \mathfrak{g}_3, \mathfrak{g}_4, \mathfrak{g}_6$  and  $\mathfrak{g}_7, \mathfrak{g}_8$ .

$\mathfrak{g}_7 = \alpha_1 \beta_{(p-1)p+1}$  and  $\mathfrak{g}_8 = \alpha_1 \beta_{p^2/p^2}$  have too low dimension to be the target of  $d_r(\beta_{p^2/p^2-1})$ . From the Toda differential  $d_{2p-1}(b_{11}) = \alpha_1 \beta_1^p$  we have


$$\begin{aligned} d_{2p-1}(\beta_1^{p^2-p-1} b_{11} \beta_2) &= \alpha_1 \beta_1^{p^2-1} \beta_2 = \mathfrak{g}_1 \\ d_{2p-1}(\mathfrak{g}_4) &= d_{2p-1}(\beta_1^{\frac{p^2-6p+1}{2}} b_{11}^2 \gamma_{\frac{p+1}{2}}) = 2\alpha_1 \beta_1^{\frac{p^2-4p+1}{2}} b_{11} \gamma_{\frac{p+1}{2}}. \end{aligned}$$

From  $d_{2p-1}(h_{20} b_{11} \gamma_s) = \alpha_1 \beta_1^p h_{20} \gamma_s$  (cf. Theorem 4.4), we have

$$d_{2p-1} \left( \beta_1^{\frac{p^2-4p-1}{2}} h_{20} b_{11} \gamma_{\frac{p+1}{2}} \right) = \alpha_1 \beta_1^{\frac{p^2-2p-1}{2}} h_{20} \gamma_{\frac{p+1}{2}} = \mathfrak{g}_3.$$

From Lemma 5.1, we have

$$d_{2p-1}(\mathfrak{g}_6) = d_{2p-1}(\beta_1^{p-1} \eta_{(p-3)p+3}) = \beta_1^{2p-1} \beta_{(p-3)p+4}.$$



Theorem A follows.  $\square$

## 6. A CONJECTURE

Consider the cofiber sequence

$$S^0 \xrightarrow{p} S^0 \longrightarrow M$$

and the induced short exact sequence of  $BP$ -homologies

$$0 \longrightarrow BP_*(S^0) \xrightarrow{p} BP_*(S^0) \longrightarrow BP_*(M) \longrightarrow 0,$$

which induces a long exact sequence of  $Ext$  groups

$$\begin{array}{ccccccc} \cdots \longrightarrow & Ext^{1,t}(BP_*(S^0)) & \longrightarrow & Ext^{1,t}(BP_*(S^0)) & \longrightarrow & Ext^{1,t}(BP_*(M)) & \xrightarrow{\delta} Ext^{2,t}(BP_*(S^0)) \longrightarrow \cdots \\ & \downarrow d_{2p-1} & & \downarrow d_{2p-1} & & \downarrow d_{2p-1} & \downarrow d_{2p-1} \\ \cdots \longrightarrow & Ext^{2p,*}(BP_*(S^0)) & \longrightarrow & Ext^{2p,*}(BP_*(S^0)) & \longrightarrow & Ext^{2p,*}(BP_*(M)) & \xrightarrow{\delta} Ext^{2p+1,*}(BP_*(S^0)) \longrightarrow \cdots \end{array}$$

For the connecting homomorphism  $\delta$ , one has

$$\delta(h_{i+2}) = \beta_{p^{i+1}/p^{i+1}}, \quad \delta(v_1 h_{i+2}) = \beta_{p^{i+1}/p^{i+1}-1} \quad \text{and} \quad \delta(v_1^i) = i\alpha_i.$$

From the Toda differential  $d_{2p-1}(\beta_{p/p}) = \alpha_1 \beta_1^p$ , one can get a non-trivial differential in the ANSS for the Moore spectrum  $M$

$$d_{2p-1}(h_2) = v_1 \beta_1^p.$$

Then from the relation  $h_{i+1} \beta_{p/p}^{p^i} = h_{i+2} \beta_1^p$  (cf. [19] and [22] 6.4.7), we get the following Adams-Novikov differential by induction

$$\begin{aligned} d_{2p-1}(h_{i+2}) \beta_1^{p^i} &= d_{2p-1}(h_{i+2} \beta_1^p) = d_{2p-1}(h_{i+1} \beta_{p/p}^{p^i}) \\ &= d_{2p-1}(h_{i+1}) \beta_{p/p}^{p^i} \\ &= v_1 \beta_{p^{i-1}/p^{i-1}} \beta_{p/p}^{p^i} \\ &= v_1 (\beta_{p^{i-1}/p^{i-1}} \beta_{p/p}^{p^{i-1}})^p \\ &= v_1 \beta_{p^i/p^i}^p \beta_1^{p^i}, \end{aligned}$$

which implies  $d_{2p-1}(h_{i+2}) = v_1 \beta_{p^i/p^i}^p$  in the ANSS for the Moore spectrum  $M$ . Then from the convergence of  $v_1$  in the ANSS for the Moore spectrum one has

$$d_{2p-1}(v_1 h_{i+2}) = v_1^2 \beta_{p^i/p^i}^p$$

Applying the connecting homomorphism  $\delta$ , we have the Adams-Novikov differential for the sphere

$$(5.2) \quad d_{2p-1}(\beta_{p^{i+1}/p^{i+1}-1}) = d_{2p-1}(\delta(v_1 h_{i+2})) = \delta(d_{2p-1}(v_1 h_{i+2})) = \delta(v_1^2 \beta_{p^i/p^i}^p) = 2\alpha_2 \beta_{p^i/p^i}^p.$$

So one can prove the non-existence of  $\beta_{p^{i+1}/p^{i+1}-1}$  from the non-triviality of

$$\alpha_2 \beta_{p^i/p^i}^p \neq 0 \in Ext_{BP_*BP}^{2p+1,*}(BP_*, BP_*).$$

(1)  $\beta_{p/p-1}$  exists and  $\alpha_2 \beta_1^p = 0$  because  $\alpha_2 \beta_1 = 0$ .

(2)  $\beta_{p^2/p^2-1}$  exists, this implies  $\alpha_2 \beta_{p/p}^p = 0$ .

As we know that  $\beta_{p/p}^p \neq 0$  in  $Ext_{BP_*BP}^{2p,qp^3}(BP_*, BP_*)$  [19, 22]. But we could not find its representative element  $b_{11}^p$  in  $Ext_{BP_*BP}^{2p,qp^3}(BP_*, BP_*(X))$  (cf. [22] 7.3.12 (b) and the ABC Theorem) because of the differential in the SDSS.

$$d(h_{11} b_{20}^{p-1}) = b_{11}^p$$

(1) At the prime  $p = 5$ ,  $\beta_1 x_{952}$  converges to  $\beta_{5/5}^5$ , where  $x_{952} = h_{11} b_{20}^{p-3} \gamma_2$ . This implies  $\alpha_2 \beta_{5/5}^5 = \alpha_2 \beta_1 x_{952} = 0$  (cf. [22] 7.5.5 stem 990) because  $\alpha_2 \beta_1 = 0$ .



(2) At the prime  $p \geq 5$ , we compute  $Ext_{BP_*BP}^{2p, qp^3}(BP_*, BP_*)$  by the SDSS. The  $E_1$ -term

$$E_1^{s,t,u} = Ext_{BP_*BP}^{s,*}(BP_*, BP_*(X)) \otimes E[\alpha_1] \otimes P[\beta_1]$$

subject to  $s + u = 2p$ ,  $t = qp^3$  is the  $\mathbb{Z}/p$  module generated by

$$\beta_1 h_{11} b_{20}^{p-3} \gamma_2, \quad \alpha_1 \beta_1 b_{20}^{p-3} \eta_p, \quad \alpha_1 \beta_1^{\frac{p-1}{2}} h_{20} b_{11}^{\frac{p-5}{2}} b_{20} \mu_{\frac{p-3}{2}p+p-2}.$$

In any case, we can conclude  $\beta_{p/p}^p$  is divisible by  $\beta_1$ . Here we believe that it is  $\beta_1 h_{11} b_{20}^{p-3} \gamma_2$  converges to  $\beta_{p/p}^p$ . So we have conjectures for the behavior of  $\beta_{p^i/p^i}^p$  in general as summarized in Conjecture C.

## REFERENCES

- [1] Adams, J. F., *On the structure and applications of the Steenrod algebra*, Comm. Math. Helv. **32** (1958), 180-214.
- [2] Cohen, R., *Odd primary infinite families in stable homotopy theory*, Mem. Amer. Math. Soc. **30** (1981) no. 242 VIII +92pp.
- [3] Cohen, R. and Goerss, P., *Secondary cohomology operations that detect homotopy classes*. Topology **23** (1984), no. 2, 177-194.
- [4] Hovey, M., *Algebraic topology problem list*, <http://claudio.math.wesleyan.edu/~mhovey/problems/index.html>.
- [5] Kato, R., Shimomura, K., *Products of greek letter elements dug up from the third morava stabilizer algebra*, Algebr. Geom. Topol. **12** (2012), 951-961.
- [6] Liulevicius, A., *The factorization of cyclic reduced powers by secondary cohomology operations*, Mem. Amer. Math. Soc. **42** (1962).
- [7] Liu, X., Wang, X., *A four-filtered May spectral sequence and its applications*, Acta Math. Sin., (Engl. Ser.) **24** (2008), 1507-1524.
- [8] May, J. P., *The cohomology of restricted Lie algebras and of Hopf algebras; Applications to the Steenrod algebra (Theses)*, Princeton (1964).
- [9] May, J. P., *The cohomology of restricted Lie algebras and of Hopf algebras*, J. Algebra **3** (1966), 123-146.
- [10] Miller, H., Ravenel, D. C., Wilson, S., *Periodic phenomena in the Adams-Novikov spectral sequence*, Ann. of Math. **106** (1977), 469-516.
- [11] Nakai, H., *The chromatic  $E_1$ -term  $H^0 M_1^2$  for  $p > 3$* , New York J. Math. **6** (2000), 21-54.
- [12] Novikov, S. P., *The methods of algebraic topology from the viewpoint of cobordism theories*, Izv. Akad. Nauk. SSSR. Ser. Mat. **31** (1967), 855-951 (Russian).
- [13] Oka, S., *A new family in the stable homotopy groups of sphere I*, Hiroshima Math. J. **5** (1975), 87-114.
- [14] Oka, S., *A new family in the stable homotopy groups of sphere II*, Hiroshima Math. J. **6** (1976), 331-342.
- [15] Oka, S., *Realizing some cyclic  $BP_*$ -modules and applications to stable homotopy of spheres*, Hiroshima Math. J. **7** (1977), 427-447.
- [16] Oka, S., *Ring spectra with few cells*, Japan. J. Math. **5** (1979), 81-100.
- [17] Oka, S., *Multiplicative structure of finite ring spectra and stable homotopy of spheres*, Algebraic Topology (Aarhus 1982) 418-41. Lect. Notes in Math. **1051** Springer-Verlag 1984.
- [18] Oka, S., *Small ring spectra and  $p$ -rank of the stable homotopy of spheres*, Contemp. Math. **19** (1983), 267-308.
- [19] Ravenel, D. C., *The nonexistence of odd primary Arf invariant elements in stable homotopy theory*, Math. Proc. Cambridge Phil. Soc. **83** (1978), 429-443.
- [20] Ravenel, D. C., *The Adams-Novikov  $E_2$ -term for a complex with  $p$ -cells*, Amer. J. Math. **107**(4) (1978), 933-968.
- [21] Ravenel, D. C., *The method of infinite descent in stable homotopy theory. I* Recent progress in homotopy theory (Baltimore, MD, 2000), Contemp. Math., vol. 293, Amer. Math. Soc., Providence, RI, 2002, pp. 251-284.
- [22] Ravenel, D. C., *Complex Cobordism and Stable Homotopy Groups of Spheres*, Academic Press, New York, 1986.
- [23] Ravenel, D. C., *Complex Cobordism and Stable Homotopy Groups of Spheres*, A. M. S. Chelsea Publishing, Providence, 2004.
- [24] Ravenel, D. C., *A novice's guide to the Adams-Novikov spectral sequence*, Proc. Evanston Homotopy Theory Conf. Lect. Notes in Math., **658**, 404-475.
- [25] Shimomura, K., *The beta elements  $\beta_{tp^2/r}$  in the homotopy of spheres*, Algebr. Geom. Topol. **10** (2010) 2079-2090.

- [26] Toda, H., *An important relation in the homotopy groups of spheres*, Proc. Japan Acad. **43** (1967), 893-942.
- [27] Toda, H., *Extended  $p$ -th powers of complexes and applications to homotopy theory*, Proc. Japan Acad., **44** (1968), 198-203.
- [28] Toda, H., *On spectra realizing exterior parts of Steenrod algebra*, Topology **10** (1971), 55-65.
- [29] Wang, X., *The secondary differentials on the third line of the Adams spectral sequence*, Topology. Appl. **156** (2009), 477-499.

JIANGUO HONG, SCIENCE COLLEGE, SHIJIAZHUANG UNIVERSITY, SHIJIAZHUANG 050035, P. R. CHINA  
Email address: `jghong66@163.com`

XIANGJUN WANG, SCHOOL OF MATHEMATICAL SCIENCE AND LPMC, NANKAI UNIVERSITY, TIANJIN 300071, P. R. CHINA  
Email address: `xjwang@nankai.edu.cn`

YU ZHANG, SCHOOL OF MATHEMATICAL SCIENCE, NANKAI UNIVERSITY, TIANJIN 300071, P. R. CHINA  
Email address: `zhang.4841@osu.edu`