

APPLIED NUMERICAL METHODS

Homework # 1

Yuhe Guo Student ID:

2021.11.01

Question 1

[Range, Linear Mapping and Matrix] Let f_1, \dots, f_8 be a set of functions defined on the interval $[1, 8]$ with the property that for any numbers d_1, \dots, d_8 , there exists a set of coefficients c_1, \dots, c_8 such that

$$\sum_{j=1}^8 c_j f_j(i) = d_i, \quad i = 1, \dots, 8$$

- (a) Show by appealing to the theorems of this lecture that d_1, \dots, d_8 determine c_1, \dots, c_8 uniquely.
(b) Let A be the 8×8 matrix representing the linear mapping from data d_1, \dots, d_8 to coefficients c_1, \dots, c_8 . What is the i, j entry of A^{-1} ?

Answer

- (a). For proof, we will use two theorems from APPLIED NUMERICAL ANALYSIS below:

Theorem 1. *Range(A) is the space spanned by the columns of A .*

Theorem 2. *A matrix $A \in \mathbb{C}^{m \times n}$ has full rank if and only if maps no two distinct vectors to the same vector.*

Proof. Denote $F = \begin{bmatrix} f_1(1) & \dots & f_8(1) \\ \dots & \dots & \dots \\ f_1(8) & \dots & f_8(8) \end{bmatrix}$. We can re-write the mapping f on c_1, \dots, c_8 to d_1, \dots, d_8 in a matrix-vector-product way:

$$F \begin{bmatrix} c_1 \\ \dots \\ c_8 \end{bmatrix} = \begin{bmatrix} d_1 \\ \dots \\ d_8 \end{bmatrix}.$$

Note that the value of d_i can be any number, and we can view $[d_i]_{i \in [1, 8]}$ to be spanned by the columns of F , so $\text{Range}(F) = 8$. By applying Theorem 1, we get that F is full-rank.

So, d_1, \dots, d_8 determine c_1, \dots, c_8 uniquely. Otherwise, we'll find two different vectors that map the columns of full-rank matrix F to a same vector, which conflicts Theorem 2. Until now We have finished sub-question (a). \square

- (b). As F is full-rank, $()$ can be written as:

$$\begin{bmatrix} c_1 \\ \dots \\ c_8 \end{bmatrix} = F^{-1} \begin{bmatrix} d_1 \\ \dots \\ d_8 \end{bmatrix}.$$

So $A = F^{-1} \rightarrow A^{-1} = F \rightarrow A_{i,j}^{-1} = F_{ij} = f_j(i)$. Until now We have finished sub-question (b).

Question 2

[Rank-one perturbation of the identity] 2.6. If u and v are m -vectors, the matrix $A = I + uv^*$ is known as a rank-one perturbation of the identity.

- Show that if A is nonsingular, then its inverse has the form $A^{-1} = I + a uv^*$ for some scalar a , and give an expression for a .
- For what u and v is A singular? If it is singular, what is $\text{null}(A)$?

The first part of proof

Suppose that A is non-singular, let $B = I + a uv^*$, we will prove that for any A , we can find a corresponding a , and thus a corresponding B , so that $B = A^{-1}$.

If $AB = I$, we get:

$$\begin{aligned} AB = I &\rightarrow (I + uv^*)(I + a uv^*) = I \\ &\rightarrow I + a uv^* + uv^* + a u(v^*u)v^* = I \\ &\rightarrow (a + 1 + a v^*u)uv^* = 0 \\ &\rightarrow a(1 + v^*u) + 1 = 0 \text{ or } uv^* = 0 \end{aligned}$$

As been given, $\text{rank}(u) = \text{rank}(v) = 1 > 0$, so the a we need satisfies:

$$a(1 + v^*u) = -1 \quad (1)$$

Next we show that $v^*u \neq -1$. Assume that $v^*u = -1$, then $Au = (I + uv^*)u = u + u * (-1) = \mathbf{0}$, conflicting that $u \neq \mathbf{0}$ and A is non-singular.

So from (1) we further get:

$$a = -\frac{1}{1 + v^*u} \quad (2)$$

The second part of proof

In this part, we show that $v^*u = -1$ is the necessary and sufficient condition when A is singular.

sufficiency

As been shown in the former part, when $v^*u = -1$, $Au = (I + uv^*)u = u + u * (-1) = \mathbf{0}$, then A is singular.

necessity

Once $v^*u \neq -1$, according to the former part, we can construct a matrix $B = I - \frac{1}{1 + v^*u} uv^*$ that $AB = I$, which indicates that A is non-singular. So we have proved the necessity.

The Null Space of A

$$\begin{aligned} Ax &= 0 \\ &\rightarrow (I + uv^*)x = 0 \\ &\rightarrow x = -(v^*x)u \\ &\rightarrow x // u \rightarrow \text{Null}(A) = \text{span}([u]) \end{aligned}$$

Question 3

[Singular Values of Rotated Matrix] Suppose A is an $m \times n$ matrix and B is the $n \times m$ matrix obtained by rotating A ninety degrees clockwise on paper (not exactly a standard mathematical transformation!). Do A and B have the same singular values? Prove that the answer is yes or give a counterexample.

Answer

Yes.

Proof

Notation 1. We use A^R as a notation for the ninety-degree clock-wisely rotated A , and A^T for the transpose of A .

For proof, we will use the theorem and corollary below:

Theorem 3. The nonzero singular values of A are the square roots of the nonzero eigenvalues of A^*A or AA^* . [NLA Theorem 5.4]

We will also use the immediate corollary of 3 below:

Corollary 1. For any matrix $M \in \mathbb{R}^{m \times n}$, AA^* and A^*A share the same singular values.

Lemma 1 (Relation between A^R and A^T). For any matrix $A \in \mathbb{R}^{m \times n}$, $A_{i,j}^T = A_{i,m-j+1}^R$

Proof. $A_{i,j}^T = A_{j,i} = A_{i,m-j+1}^R$. □

Lemma 2 (Singular value of A^R and A^T). A^R and A^T share the same singular values.

Proof. Denote A^T to be C , A^R to be D . According to 1, the elements in each row i of C and D are the same, they are just permuted (more precisely, permuted in reversed orders), and the permutation are parallel among each row. So the inner product of any two rows in C and D are the same:

$$\langle C[i, :], C[j, :] \rangle = \langle D[i, :], D[j, :] \rangle$$

So:

$$CC^T = [\langle C[i, :], C[j, :] \rangle]_{ij} = \langle D[i, :], D[j, :] \rangle = DD^T \quad (3)$$

By Theorem 3 and (3) we derive that C and D share the same singular values. Thus, we have finished the proof of Lemma 2. □

Proof. Back to the question. By 2, B shares the same singular values with A^T ; By 1, A^T shares the same singular values with A . So A and B have the same singular values. □

Question 4

[Practice with SVD]

Consider the matrix

$$A = \begin{bmatrix} -2 & 11 \\ -10 & 5 \end{bmatrix}$$

1. Determine, on paper, a real SVD of A in the form $A = U\Sigma V^T$. The SVD is not unique, so find the one that has the minimal number of minus signs in U and V ;
2. List the singular values, left singular vectors, and right singular vectors of A . Draw a careful, labeled picture of the unit ball in \mathbb{R}^2 and its image under A , together with the singular vectors, with the coordinates of their vertices marked;
3. What are the 1-, 2-, ∞ -, and Frobenius norms of A ?
4. Find A^{-1} not directly, but via the SVD;
5. Find the eigenvalues λ_1, λ_2 of A ;
6. Verify that $\det A = \lambda_1 \lambda_2$ and $|\det A| = \sigma_1 \sigma_2$;
7. What is the area of the ellipsoid onto which A maps the unit ball in \mathbb{R}^2 ?

Answer

1. SVD Calculation

$$\begin{aligned} A &= U\Sigma V^T = \begin{bmatrix} -2 & 11 \\ -10 & 5 \end{bmatrix} \\ \rightarrow AA^T &= U\Sigma^2 U^T = \begin{bmatrix} 125 & 75 \\ 75 & 125 \end{bmatrix}, \\ A^T A &= V\Sigma^2 V^T = \begin{bmatrix} 104 & -72 \\ -72 & 146 \end{bmatrix} \end{aligned}$$

Conduct eigen decomposition on AA^T and $A^T A$, we get:

$$U = \begin{bmatrix} -\sqrt{2}/2 & \sqrt{2}/2 \\ \sqrt{2}/2 & \sqrt{2}/2 \end{bmatrix}, \quad \Sigma = \begin{bmatrix} 5\sqrt{2} & 0 \\ 0 & 10\sqrt{2} \end{bmatrix}, \quad V = \begin{bmatrix} -4/5 & -3/5 \\ -3/5 & 4/5 \end{bmatrix}$$

We adjust U and V so that we get the minimal number of minus signs, we turn the sign of U 's first column and V 's first column, and get:

$$U = \begin{bmatrix} \sqrt{2}/2 & -\sqrt{2}/2 \\ \sqrt{2}/2 & \sqrt{2}/2 \end{bmatrix}, \quad \Sigma = \begin{bmatrix} 5\sqrt{2} & 0 \\ 0 & 10\sqrt{2} \end{bmatrix}, \quad V = \begin{bmatrix} 4/5 & 3/5 \\ -3/5 & 4/5 \end{bmatrix} \quad (4)$$

2. Visualization of SVD.

From (4) we directly get:

- Singular values:

$$\sigma_1 = 5\sqrt{2}, \quad \sigma_2 = 10\sqrt{2}; \quad (5)$$

- Left singular vectors: $[-\sqrt{2}/2, \sqrt{2}/2]^T, [\sqrt{2}/2, \sqrt{2}/2]^T$;
- Right singular vectors : $[4/5, -3/5]^T, [3/5, 4/5]^T$.

The unit ball's image under A is show as follow:




Figure 1: The unit ball's image under A

3. Matrix's Norms

$$\begin{aligned} \|A\|_1 &= 28, \\ \|A\|_2 &= 5\sqrt{10}, \\ \|A\|_\infty &= 11, \\ \|A\|_F &= 5\sqrt{10} \end{aligned}$$

4. A^{-1} By SVD **[TODO: Check again!]**

$$A^{-1} = (U\Sigma V^T)^{-1} = V\Sigma^{-1}U^T = \begin{bmatrix} 0.05 & 0.11 \\ -0.1 & -0.02 \end{bmatrix}$$

5. Eigen Values of A^{-1}

The characteristic polynomial of A is:

$$p_A(\lambda) = \det(A - \lambda I) = \lambda^2 - 3\lambda + 100,$$

so that we obtain:

$$\begin{cases} \lambda_1 = \frac{3 + \sqrt{-391}}{2} \\ \lambda_2 = \frac{3 - \sqrt{-391}}{2} \end{cases} \quad (6)$$

6. Relationship of Eigen Values and Matrix's Determinant and Trace

From (6) and (5), we get:

$$\begin{cases} \lambda_1 \lambda_2 = 100 = \det A \\ \sigma_1 \sigma_2 = 100 = |\det A| \end{cases}$$

7. Ellipsoid Mapped A^{-1}

Figure 1 has been shown in sub-question 2, now we describe the mapped-to ellipsoid analytically:
The unit ball before mapped by A is :

$$\{ (x, y) \mid [x, y] \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} \leq 1 \}.$$

After being mapped by A , the set of points comes to:

$$\{ (x, y) \mid ([x, y] A^T) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} (A \begin{bmatrix} x \\ y \end{bmatrix}) \leq 1 \},$$

which can be written as:

$$\{ (x, y) \mid [x, y] A^T A \begin{bmatrix} x \\ y \end{bmatrix} \leq 1 \}. \quad (7)$$

Equation (7) is just the analytical solution of obtain ellipsoid.

Question 5

[\[\(Orthogonal\) Projector onto Column Range\]](#) Consider the matrices:

$$A = \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 1 & 0 \end{bmatrix}, \quad B = \begin{bmatrix} 1 & 2 \\ 0 & 1 \\ 1 & 0 \end{bmatrix}$$

Answer the following questions by hand calculation.

- What is the orthogonal projector P onto $\text{range}(A)$, and what is the image under P of vector $[1, 2, 3]^*$?
- Same question for B .

Answer

- The projector is:

$$P = A(A^T A)^{-1} A^T = \begin{bmatrix} \frac{1}{2} & 0 & \frac{1}{2} \\ 0 & 1 & 0 \\ \frac{1}{2} & 0 & \frac{1}{2} \end{bmatrix}.$$

The image under P of $[1, 2, 3]^*$ is:

$$P[1, 2, 3]^* = [2, 2, 2]^*.$$

- The projector is:

$$P = B(B^T B)^{-1} B^T = \begin{bmatrix} 5/6 & 1/3 & 1/6 \\ 1/3 & 1/3 & -1/3 \\ 1/6 & -1/3 & 5/6 \end{bmatrix}.$$

The image under P of $[1, 2, 3]^*$ is:

$$P[1, 2, 3]^* = [2, 0, 2]^*.$$

Question 6

[QR Factorization] Consider again the matrices A and B of question 6.

(a) Using any method you like, determine (on paper) a reduced QR factorization $A = \hat{Q}\hat{R}$ and a full QR factorization $A = QR$.

(b) Again using any method you like, determine reduced and full QR factorizations $B = \hat{Q}\hat{R}$ and $B = QR$.

Answer

(a1)[Reduced QR factorization of A]. Denote the reduced QR factorization of A as $A = \hat{Q}\hat{R}$, $\hat{Q} \in \mathbb{R}^{3 \times 2}$, $\hat{R} \in \mathbb{R}^{2 \times 2}$,

$$A = [\mathbf{a}_1, \mathbf{a}_2], \quad \hat{Q} = [\mathbf{q}_1, \mathbf{q}_2], \quad \hat{R} = \begin{bmatrix} r_{11} & r_{12} \\ 0 & r_{22} \end{bmatrix},$$

$$\mathbf{a}_1, \mathbf{a}_2 \in \mathbb{R}^{3 \times 1}, \quad \mathbf{q}_1, \mathbf{q}_2 \in \mathbb{R}^{3 \times 1},$$

$$r_{11}, r_{12}, r_{22} \in \mathbb{R}.$$

As been given, $\mathbf{a}_1 \perp \mathbf{a}_2$, so we directly derive that:

$$\mathbf{q}_1 = \mathbf{a}_1 / \|\mathbf{a}_1\|, \quad r_{11} = \|\mathbf{a}_1\| \quad (8)$$

$$\mathbf{q}_2 = \mathbf{a}_2 / \|\mathbf{a}_2\|, \quad r_{22} = \|\mathbf{a}_2\| \quad (9)$$

, and

$$\begin{aligned} \mathbf{a}_2 &= r_{12}\mathbf{q}_1 + r_{22}\mathbf{q}_2 \\ \Rightarrow \langle \mathbf{a}_2, \mathbf{q}_1 \rangle &= r_{12}\langle \mathbf{q}_1, \mathbf{q}_1 \rangle + r_{22}\langle \mathbf{q}_2, \mathbf{q}_1 \rangle \\ &\Rightarrow r_{12} = 0. \end{aligned} \quad (10)$$

Compounding equations (8) (9) and (10), we get we the reduced QR factorization of A:

$$\hat{Q} = \begin{bmatrix} \sqrt{2}/2 & 0 \\ 0 & 1 \\ \sqrt{2}/2 & 0 \end{bmatrix}, \quad \hat{R} = \begin{bmatrix} \sqrt{2} & 0 \\ 0 & 1 \end{bmatrix}$$

(a2)[Full QR factorization of A]. Denote the full QR factorization of A as $A = QR$, $Q \in \mathbb{R}^{3 \times 3}$, $R \in \mathbb{R}^{3 \times 2}$, note that in the equations below, all the values except for \mathbf{q}_3 keep the same with in reduced QR factorization:

$$A = [\mathbf{a}_1, \mathbf{a}_2], \quad Q = [\mathbf{q}_1, \mathbf{q}_2, \mathbf{q}_3], \quad R = \begin{bmatrix} r_{11} & r_{12} \\ 0 & r_{22} \\ 0 & 0 \end{bmatrix},$$

As $\mathbf{q}_3 \perp \mathbf{q}_1$ and $\mathbf{q}_3 \perp \mathbf{q}_2$, $\mathbf{q}_3 \in \text{Null}(\hat{Q})$,

$$\begin{aligned} \hat{Q}^T \mathbf{q}_3 &= 0, \quad \|\mathbf{q}_3\| = 1 \\ \Rightarrow \mathbf{q}_3 &= \pm[\sqrt{2}/2, 0, -\sqrt{2}/2]^T \end{aligned}$$

Let $\mathbf{q}_3 = [\sqrt{2}/2, 0, -\sqrt{2}/2]^T$, we get the full QR factorization of A :

$$Q = \begin{bmatrix} \sqrt{2}/2 & 0 & \sqrt{2}/2 \\ 0 & 1 & 0 \\ \sqrt{2}/2 & 0 & -\sqrt{2}/2 \end{bmatrix}, \quad R = \begin{bmatrix} \sqrt{2} & 0 \\ 0 & 1 \\ 0 & 0 \end{bmatrix}$$

(b1)[Reduced QR factorization of B]. Denote the reduced QR factorization of B as $B = \hat{Q}\hat{R}$, $\hat{Q} \in \mathbb{R}^{3 \times 2}$, $\hat{R} \in \mathbb{R}^{2 \times 2}$,

$$B = [\mathbf{b}_1, \mathbf{b}_2], \quad \hat{Q} = [\mathbf{q}_1, \mathbf{q}_2], \quad \hat{R} = \begin{bmatrix} r_{11} & r_{12} \\ 0 & r_{22} \end{bmatrix},$$

$$\mathbf{b}_1, \mathbf{b}_2 \in \mathbb{R}^{3 \times 1}, \mathbf{q}_1, \mathbf{q}_2 \in \mathbb{R}^{3 \times 1}$$

$$r_{11}, r_{12}, r_{22} \in \mathbb{R}.$$

By the inherent property of QR factorization,

$$\begin{aligned} \mathbf{q}_1 &\perp \mathbf{q}_2, \\ \text{span}[\mathbf{q}_1] &= \text{span}[\mathbf{b}_1], \\ \text{span}[\mathbf{q}_1, \mathbf{q}_2] &= \text{span}[\mathbf{b}_1, \mathbf{b}_2] \end{aligned}$$

we derive that:

$$\begin{aligned} \mathbf{q}_1 &= \mathbf{b}_1 / \|\mathbf{b}_1\|, \quad r_{11} = \|\mathbf{b}_1\| \\ \Rightarrow \mathbf{q}_1 &= [\sqrt{2}/2, 0, \sqrt{2}/2]^T, \quad r_{11} = \sqrt{2} \end{aligned} \tag{11}$$

, and

$$\begin{aligned} \mathbf{b}_2 &= r_{12}\mathbf{q}_1 + r_{22}\mathbf{q}_2 \\ \Rightarrow \langle \mathbf{b}_2, \mathbf{q}_1 \rangle &= r_{12}\langle \mathbf{q}_1, \mathbf{q}_1 \rangle + r_{22}\langle \mathbf{q}_2, \mathbf{q}_1 \rangle \end{aligned} \tag{12}$$

$$\Rightarrow r_{12} = \langle \mathbf{b}_2, \mathbf{q}_1 \rangle = \sqrt{2}. \tag{13}$$

Feed (13) into (12), and apply $\|\mathbf{q}_2\| = 1$ we get:

$$\begin{aligned} r_{22}\mathbf{q}_2 &= \mathbf{b}_2 - r_{12}\mathbf{q}_1 = [1, 1, -1]^T \\ \Rightarrow r_{22} &= \sqrt{3}, \mathbf{q}_2 = [\sqrt{3}/3, \sqrt{3}/3, -\sqrt{3}/3] \end{aligned} \tag{14}$$

Compounding equations (11) to (14), we get the reduced QR factorization of B :

$$B = \hat{Q}\hat{R}, \quad \hat{Q} = \begin{bmatrix} \sqrt{2}/2 & \sqrt{3}/3 \\ 0 & \sqrt{3}/3 \\ \sqrt{2}/2 & -\sqrt{3}/3 \end{bmatrix}, \quad \hat{R} = \begin{bmatrix} \sqrt{2} & \sqrt{2} \\ 0 & \sqrt{3} \end{bmatrix}.$$

(b2)[Full QR factorization of B]. Denote the full QR factorization of B as $B = QR$, $Q \in \mathbb{R}^{3 \times 3}$, $R \in \mathbb{R}^{3 \times 2}$, note that in the equations below, all the values except for \mathbf{q}_3 keep the same with in reduced QR factorization:

$$B = [\mathbf{b}_1, \mathbf{b}_2], \quad Q = [\mathbf{q}_1, \mathbf{q}_2, \mathbf{q}_3], \quad R = \begin{bmatrix} r_{11} & r_{12} \\ 0 & r_{22} \\ 0 & 0 \end{bmatrix},$$

As $\mathbf{q}_3 \perp \mathbf{q}_1$ and $\mathbf{q}_3 \perp \mathbf{q}_2$, $\mathbf{q}_3 \in \text{Null}(\hat{Q})$,

$$\begin{aligned} \hat{Q}^T \mathbf{q}_3 &= 0, \quad \|\mathbf{q}_3\| = 1 \\ \Rightarrow \mathbf{q}_3 &= \pm[-\sqrt{6}/6, \sqrt{6}/3, \sqrt{6}/6]^T \end{aligned}$$

Let $\mathbf{q}_3 = [-\sqrt{6}/6, \sqrt{6}/3, \sqrt{6}/6]^T$, we get the full QR factorization of B :

$$B = QR, \quad Q = \begin{bmatrix} \sqrt{2}/2 & \sqrt{3}/3 & -\sqrt{6}/6 \\ 0 & \sqrt{3}/3 & \sqrt{6}/3 \\ \sqrt{2}/2 & -\sqrt{3}/3 & \sqrt{6}/6 \end{bmatrix}, \quad R = \begin{bmatrix} \sqrt{2} & \sqrt{2} \\ 0 & \sqrt{3} \\ 0 & 0 \end{bmatrix}.$$

Question 7

[QR Factorization with MATLAB Householder]

(a) Write a Matlab function $[W, R] = \text{house}(A)$ that computes an implicit representation of a full QR factorization $A = QR$ of an $m \times n$ matrix A with $m > n$ using Householder reflections. The output variables are a lower-triangular matrix $W \in \mathbb{C}^{m \times n}$ whose columns are the vectors v_k defining the successive Householder reflections, and a triangular matrix $R \in \mathbb{C}^{n \times n}$.

(b) Write a Matlab function $Q = \text{formQ}(W)$ that takes the matrix W produced by house as input and generates a corresponding $m \times m$ orthogonal matrix Q .

Answers / Matlab Codes

Listing 1: Code for $\text{house}(A)$

```
1 function [W, R] = house(A)
2     [m, n] = size(A)
3     W = zeros(m, n)
4     e1 = eye(m, 1)
5     for k=1:n
6         x = A(k:m, k)
7         % We use e1_sub together with x to decide reflector,
8         % as the dimension of subspace (submatrix) is reduced in each loop,
9         % e1_sub comes shorter and shorter.
10        e1_sub = e1(1:m-k+1)
11        % Reflector; Normal plane
12        vk = sign(x(1))*norm(x,2) * e1_sub + x
13        vk = vk / norm(vk,2)
14        % Map the submatrix. To improve efficiency,
15        % it's important to calculate (vk'*A(k:m,k:n)) first.
16        A(k:m,k:n) = A(k:m,k:n) - 2*vk*(vk'*A(k:m,k:n))
17        % Record the sub-vectors who define the mappings
18        W(k:m,k) = vk
19    end
20 end
```

Listing 2: Test Code for $\text{house}(A)$

```
1 m = 8
2 n = 4
3 x = (-m/2:m/2-1)'/(m/2)
4 A = [x.^0 x.^1 x.^2 x.^3]
5 W, R = house(A)
```

Listing 3: Code for $\text{formQ}(Q)$

```
1 function [Q] = formQ(W)
2     [m, m] = size(W)
3     QT = eye(m)
4     for k=1:n
5         vk = W(:,k)
6         % REDO the functions A
7         QT = QT - 2*vk*(vk'*QT)
8     end
9     Q = QT.'
```

Question 8

[MATLAB]

11.3. Take $m = 50$, $n = 12$. Using Matlab's `linspace`, define t to be the m -vector corresponding to linearly spaced grid points from 0 to 1. Using Matlab's `vander` and `fliplr`, define A to be the $m \times n$ matrix associated with least squares fitting on this grid by a polynomial of degree $n - 1$. Take b to be the function $\cos(4t)$ evaluated on the grid. Now, calculate and print (to sixteen-digit precision) the least squares coefficient vector x by six methods: (a) Formation and solution of the normal equations, using Matlab's (b) QR factorization computed by `mgs` (modified Gram-Schmidt, Exercise 8.2), (c) QR factorization computed by `house` (Householder triangularization, Exercise 10.2), (d) QR factorization computed by Matlab's `qr` (also Householder triangularization), (e) $x = A \backslash b$ in Matlab (also based on QR factorization), (f) SVD, using Matlab's `svd`. (g) The calculations above will produce six lists of twelve coefficients. In each list, shade with red pen the digits that appear to be wrong (affected by rounding error). Comment on what differences you observe. Do the normal equations exhibit instability? You do not have to explain your observations.

Question 9

[\[Hadmad\]](#)