

A
MGH259, oligodendroglioma G3
cohort 1
IDHA17, astrocytoma G2
cohort 2
C1b (z)
2
1
0
-1
-2
Gi* (FDR<0.05)
Hotspot
Coldspot
B
Cohort 1 (GSE237183)
Cohort 2 (Hoefflin)
Z=3.3
Z=7.0
Z=-3.0
Z=-1.5
Moran's I vs 1,000 matched
random gene sets (z)
10
5
0
-5
C1b
C1b | TAM
C
Mac
Reactive-Ast
Vasc
MES
Inflammatory-Mac
Oligo
MES-Hyp
MES-Ast
chromatin-reg
AC
OPC
Prolif-Metab
Neuron
NPC
Cohort 1
Cohort 2
Specific
co-loc. -
avoidance
1.0
0.5
0.0
-0.5
-1.0
Fraction of
sections
0.00
0.25
0.50
0.75
1.00
D
q < 0.05
n.s.
C1Q+ TAM
Endothelium
Pericytes
Microglia-like TAM
SPP1+/lipid TAM
Inflammation
Type I IFN
IFN-gamma
Monocyte-like TAM
Hypoxia
0
5
10
Enrichment inside C1b hotspots
(Stouffer z; MSR null, depth-adjusted)
E
Endothelium
Pericytes
C1Q+ TAM
Hypoxia
Target score (z, mean ± SE)
0.9
0.6
0.3
0.0
0 1 2 3 4 5 ≥6
Distance to nearest C1b hotspot (spots; 0 = inside)
F
C1b (full)
MHC-II arm | TAM
Mesenchymal arm | TAM
Cohort 2 IDHm
Cohort 1 GBM
Pericytes
Endothelium
IFN-gamma
Type I IFN
Inflammation
Hypoxia
-4
0
4
8
Enrichment inside hotspots (Stouffer z)