

Next-Generation Polarized ^3He Targets for Electron Scattering Experiments

by

© Yunxiao Wang

A thesis submitted to the
School of Graduate Studies
in partial fulfilment of the
requirements for the degree of
Doctor of Philosophy

Department of Physics
University of Virginia

November 2016

Abstract

Historically, ^3He targets for electron scattering experiments have been polarized through spin-exchange optical pumping (SEOP). Polarized laser light passes its circular polarization to alkali metal vapor, which then transfers its polarization to ^3He through spin-exchange collisions.

This thesis discusses the basics of SEOP and the polarimetry techniques used in our lab. Narrowband laser and alkali-hybrid SEOP have improved the performance of targets significantly. In alkali-hybrid SEOP, potassium is used together with rubidium for transferring polarization to ^3He nuclei. We discussed the data collected over many pure-rubidium targets and alkali-hybrid targets. In the course of analyzing the data, we also studied the “X factor” which limits the highest achievable polarization of ^3He .

Because the experiments planned for the 12GeV era in Jefferson National Laboratory (JLAB) will use much higher electron beam current, we are exploring the possibility of using metal (instead of glass) as the entry points (commonly referred to as “end windows”) for future targets. We established the metal composition and developed the techniques to incorporate metal to targets without introducing significant spin-relaxation rates. We have successfully demonstrated that future targets can be constructed with metal end windows and are very close to making such targets.

Acknowledgements

First and foremost, I'd like to give my thanks to my advisor, Gordon Cates, for supporting me in the past five years in the group. Not only because I have learned a lot about physics from you, but also for your incredible enthusiasm that kept motivating me to push towards earning my degree. I am honored to have worked with you.

I'd like to thank Peter Dolph and Yuan Zheng, who have taught me the basics of spin-exchange optical pumping and ^3He and alkali polarimetry. Even though the time we spent working together was not long, it laid the very foundation for my work in the following years.

I would like to thank Dr. W. Al Tobias and Dr. Vladimir Nelyubin as well. Al has helped us fill almost every cell we have studied and also taught me the right way to do experiments and the importance of documentation. Vladimir always made sure our experiments could go as smoothly as possible with his laser expertise. We would not have been able to do it without the help from Al and Vladimir.

I also would like to thank all the other graduate students in the group. Maduka Kaluarachchi and Dan Matyas, the two of you made the hours in lab such a joy. I will always miss our talks and of course, the "Young Chicken" we shared at Taste of China. Graduate school was not an easy time for me, I could not have done it without the help from you guys. Sumudu Katugampola and Dave Keder, the two of you are already doing great work, I have no doubt that you will have tremendous amount of success in the future. I look forward to seeing what our lab will be able to achieve because of you.

I'd like to thank the members of my committee: Nilanga Liyanage, Don Crabb,

Wilson Miller, for helping me complete the final step towards my degree. I also owe my thanks to my parents and all my friends who have supported me through all these years. It means a lot to me knowing that no matter what comes next, I will always have your support.

Last but not least, I would like to thank my girlfriend. Shuangshuang, every bit of success I was able to achieve since I met you was at least partly because of your love and support. You gave me something to hold on to so I could pull myself through a tough time. I would not have been able to move on to the next chapter of my life the same way I did without you.

Contents

Abstract	ii
Acknowledgements	iii
List of Tables	vi
List of Figures	vii
1 Conclusions and Discussion	1
1.1 Overall Design of Next-Generation Polarized ^3He Targets	2
1.2 Incorporating Metal End Windows	4
1.3 Summary	5
Bibliography	7

List of Tables

List of Figures

1.1	Design of next-generation target for upcoming SBS G_E^n experiments. .	2
-----	--	---

Chapter 1

Conclusions and Discussion

The work presented in this thesis has made it possible to design and begin producing the next-generation polarized ^3He targets for use at Jefferson Laboratory. These next-generation targets are being developed and produced in two steps, which are referred to internally at JLab as the Stage-I and Stage-II designs. The Stage-I targets contain a volume of 3 STP liters of ^3He , the same quantity that was contained in the cell Antoinette, the results from which were described in Chapter 4. The Stage-II targets, while similar in geometry, are larger and will contain 6 STP liters of ^3He . At the time of this writing, the first Stage-I target cell has already been produced and is undergoing bench tests. When using only half the design laser power, the target has already achieved a polarization of 65%, higher than what is needed for actual running (although we note that this is without the depolarizing effects of an electron beam). These early tests are quite encouraging when extrapolated to the full laser power that will be used. Also, the conceptual design of the Stage-II target cells has been completed, and was evaluated in a review, conducted in March of 2016, of the




Figure 1.1: Design of next-generation target for upcoming SBS G_E^n experiments.

polarized ^3He target that is being built for the Hall A Super Bigbite Spectrometer (SBS) experiment to measure G_E^n , the electric form factor of the neutron. The design for the SBS G_E^n target is shown in Fig 1.1.

1.1 Overall Design of Next-Generation Polarized ^3He Targets

The target-cell design illustrated in Fig. 1.1 incorporates multiple features that distinguish it from earlier polarized ^3He targets based on spin-exchange optical pumping. The basic geometry is what we refer to as a “convection-style” cell, in which the pumping and target chambers are connected by two transfer tubes, one of which is heated, resulting in controllable convective flow between the target’s two principle chambers.

The development of convection-style cells was an important part of the thesis work of Peter Dolph, and is well documented in Ref. [32]. Subsequent to the work described in [32], our group constructed and tested two convection-style prototype target cells.

Another distinguishing feature of the design shown in Fig. 1.1 is that it is significantly larger than previous target cells, containing 6 STP liters of ^3He , and has a target chamber that is 60 cm in length, 50% longer than any target cells previously operated at JLab. The confidence that this target will achieve polarization $> 60\%$ in $60\mu\text{A}$ of beam came as a direct result of the work described in Chapter 4, and published in Ref. [60]. While much of the data presented in Ref. [60] were obtained prior to the work described here, most of the analysis was performed as part of this work, including a determination of the coefficient characterizing spin-exchange between potassium and ^3He .

Data included in ref. [60] that were specifically obtained as part of this thesis work included all of the studies of the cell Antoinette, which contained roughly 3 STP liters of ^3He . While target-cells containing 3 STP liters of ^3He were used during the first Hall A G_E^n experiment, commercial narrow-band high-power diode-laser arrays, which are critical to achieving high performance, were not yet available during both testing and the experiment itself. Thus, Antoinette was the first target cell tested with narrow-band lasers that contained both 3 STP liters of ^3He and an alkali-hybrid mixture for optical pumping. As such, Antoinette became a proof of principle for both of the above-mentioned prototype convection-style target cells, as well as the first of the Stage-I target cells that is undergoing testing at the time of this writing.

In short, the work presented here provided the basis for designing the high-performance Stage-I and Stage-II target cells that will be used in future JLab ex-

periments. The proof-of-principle began with the cell Antoinette, and continued with the two prototype convection-style target cells (both of which also contained roughly 3 STP liters of ^3He). The first actual Stage-I target cell looks very promising in early tests, providing further confidence in the design of the Stage-II target cells that will soon be produced.

1.2 Incorporating Metal End Windows

Prior to the work described here, there was only very limited experience with spin-polarized noble gases in cells containing metal. The reason is that the introduction of any new material generally causes spin relaxation greatly in excess to that caused by the walls of the glass container. The target-cell design shown in Fig. 1.1, however, incorporates metal end windows so that even a fairly intense electron beam will not cause the target cell to rupture, even after multiple weeks of operation. One of the important achievements of the current work was the development of a technique for producing metal surfaces that induce spin relaxation at an acceptably slow rate. We further have demonstrated a means for making transitions between glass and metal, even when operating at the high pressures used in our polarized ^3He targets. We describe next why metal end windows based on the techniques demonstrated here are likely to have an almost negligible effect on the overall performance of our target cells.

The spin relaxation caused by metal end windows in future target cells can be calculated using the equation $\Gamma_{metal} = \rho_{metal} S_{metal} / V_{total}$ (see Chapter 2.4). Furthermore, the relaxivity associated with metal can be extracted by comparing a glass-metal-glass

test cell (such as GoldRush in Table. ??) with an all-glass control cell (such as Pyrah in Table.??). The resulting relaxivity is 0.03144 cm/hr and 0.1231 cm/hr for metal. Using this relaxivity and the designed dimensions for the target end windows, the additional relaxation times would be $1/130.6 \text{ hr}^{-1}$ for Stage-I cells and $1/261.2 \text{ hr}^{-1}$ for Stage-II cells.

During the 6 GeV era where only 10-15 μA was used, glass end windows were already running into risk of rupturing after 4-6 weeks of being exposed to electron beam. If it was solely due to radiation damage, one can expect the glass windows to rupture after roughly a week of being used in electron beam of 60 μA , which will be far from enough for the experiments to complete. On the other hand, experience at JLab suggests that even very thin metal windows should still be able to survive the electron beams. As a an example, aluminum as thin as 2 mils has been routinely used in JLab without failing. The fact that metal end windows will conduct heat better further suggests that they will be more suitable for experiments planned for the 12 GeV era. Copper test pieces with thin hemispherical end caps using OFHC copper and aluminum will be made into test cells based on the design shown in Fig. 1.1 before the planned experiments are conducted.

1.3 Summary

We have confidence that convection style targets with metal end windows will not only give high ^3He polarization in both PC and TC, but also survive the high electron beam currents planned for the future experiments. The work done in this thesis has demonstrated that the additional spin-relaxation rate due to surfaces in metal end

windows will be negligible for our purposes and has provided techniques for connecting metal end windows to glass. All the tests so far have been performed on test cells with metal tubes. Before a target cell with next-generation design can be produced, tests exploring techniques for making hemispherical metal end caps should be carried out. However, with techniques established so far, we believe the next-generation design will be used to great success in the upcoming experiments planned for the 12 GeV era.

Bibliography

- [1] Nuclear relaxation of ^3He gas on various solid surfaces. *Canadian Journal of Physics*, 1971.
- [2] Physics of practical spin-exchange optical pumping. *PhD thesis, University of Wisconsin-Madison*, 2001.
- [3] Spin-exchange optical pumping with alkali-metal vapors. *PhD thesis, University of Wisconsin-Madison*, 2005.
- [4] Alkali-hybrid spin-exchange optically-pumped polarized ^3He targets used for studying neutron structure. *PhD thesis, University of Virginia*, 2010.
- [5] High-performance nuclear-polarized ^3He targets for electron scattering based on spin-exchange optical pumping. *PhD thesis, University of Virginia*, 2010.
- [6] A. Abragam. *Principles of Nuclear Magnetism*.
- [7] M. S. Albert, G. D. Cates, B. Driehuys, W. Happer, B. Saam, C. S. Springer, and A. Wishnia. Biological magnetic resonance imaging using laser-polarized ^{129}Xe . *Nature*, 370.

- [8] M. Amarian, L. Auerbach, T. Averett, J. Berthot, P. Bertin, W. Bertozzi, T. Black, E. Brash, D. Brown, E. Burtin, J. R. Calarco, G. D. Cates, Z. Chai, J.-P. Chen, S. Choi, E. Chudakov, E. Cisbani, C. W. de Jager, A. Deur, R. DiSalvo, S. Dieterich, P. Djawotho, M. Finn, K. Fissum, H. Fonvieille, S. Frullani, H. Gao, J. Gao, F. Garibaldi, A. Gasparian, S. Gilad, R. Gilman, A. Glamazdin, C. Glashauser, E. Goldberg, J. Gomez, V. Gorbenko, J.-O. Hansen, F. W. Hersman, R. Holmes, G. M. Huber, E. W. Hughes, T. B. Humensky, S. Incerti, M. Iodice, S. Jensen, X. Jiang, C. Jones, G. M. Jones, M. Jones, C. Jutier, A. Ketikyan, I. Kominis, W. Korsch, K. Kramer, K. S. Kumar, G. Kumbartzki, M. Kuss, E. Lakuriqi, G. Laveissiere, J. Lerose, M. Liang, N. Liyanage, G. Lolos, S. Malov, J. Marroncle, K. McCormick, R. McKeown, Z.-E. Meziani, R. Michaels, J. Mitchell, Z. Papandreou, T. Pavlin, G. G. Petratos, D. Pripstein, D. Prout, R. Ransome, Y. Roblin, D. Rowntree, M. Rvachev, F. Sabatie, A. Saha, K. Slifer, P. A. Souder, T. Saito, S. Strauch, R. Suleiman, K. Takahashi, S. Teiji, L. Todor, H. Tsubota, H. Ueno, G. Urciuoli, R. Van der Meer, P. Vernin, H. Voskanian, B. Wojtsekhowski, F. Xiong, W. Xu, J.-C. Yang, B. Zhang, and P. Zolnierczuk. Q^2 evolution of the generalized gerasimov-drell-hearn integral for the neutron using a ^3He target. *Phys. Rev. Lett.*, 89:242301, Nov 2002.
- [9] P. L. Anthony, R. G. Arnold, H. R. Band, H. Borel, P. E. Bosted, V. Breton, G. D. Cates, T. E. Chupp, F. S. Dietrich, J. Dunne, R. Erbacher, J. Fellbaum, H. Fonvieille, R. Gearhart, R. Holmes, E. W. Hughes, J. R. Johnson, D. Kawall, C. Keppel, S. E. Kuhn, R. M. Lombard-Nelsen, J. Marroncle, T. Maruyama, W. Meyer, Z.-E. Meziani, H. Middleton, J. Morgenstern, N. R. Newbury, G. G.

- Petratos, R. Pitthan, R. Prepost, Y. Roblin, S. E. Rock, S. H. Rokni, G. Shapiro, T. Smith, P. A. Souder, M. Spengos, F. Staley, L. M. Stuart, Z. M. Szalata, Y. Terrien, A. K. Thompson, J. L. White, M. Woods, J. Xu, C. C. Young, and G. Zapalac. Determination of the neutron spin structure function. *Phys. Rev. Lett.*, 71:959–962, Aug 1993.
- [10] S. Appelt, A. B.-A. Baranga, C. J. Erickson, M. V. Romalis, A. R. Young, and W. Happer. Theory of spin-exchange optical pumping of ^3He and ^{129}Xe . *Phys. Rev. A*, 58:1412–1439, Aug 1998.
- [11] E. Babcock, B. Chann, T. G. Walker, W. C. Chen, and T. R. Gentile. Limits to the polarization for spin-exchange optical pumping of ^3He . *Phys. Rev. Lett.*, 96:083003, Mar 2006.
- [12] E. Babcock, I. Nelson, S. Kadlecsek, B. Driehuys, L. W. Anderson, F. W. Hersman, and T. G. Walker. Hybrid spin-exchange optical pumping of ^3He . *Phys. Rev. Lett.*, 91:123003, Sep 2003.
- [13] E. Babcock, I. A. Nelson, S. Kadlecsek, and T. G. Walker. ^3He polarization-dependent epr frequency shifts of alkali-metal- ^3He pairs. *Phys. Rev. A*, 71:013414, Jan 2005.
- [14] R. M. Barrer. *Diffusion in and through Solids*.
- [15] A. Ben-Amar Baranga, S. Appelt, M. V. Romalis, C. J. Erickson, A. R. Young, G. D. Cates, and W. Happer. Polarization of ^3He by spin exchange with optically pumped rb and k vapors. *Phys. Rev. Lett.*, 80:2801–2804, Mar 1998.

- [16] F. Bloch. Nuclear induction. *Phys. Rev.*, 70:460–474, Oct 1946.
- [17] M. A. Bouchiat, T. R. Carver, and C. M. Varnum. Nuclear polarization in he^3 gas induced by optical pumping and dipolar exchange. *Phys. Rev. Lett.*, 5:373–375, Oct 1960.
- [18] G. Breit and I. I. Rabi. Measurement of nuclear spin. *Phys. Rev.*, 38:2082–2083, Dec 1931.
- [19] A. C., V. Itkin, and H. M.K. *Canadian Metallurgical Quarterly*, 23, 1984.
- [20] P. Callaghan. *Principles of Nuclear Magnetic Resonance Microscopy*.
- [21] G. D. Cates. Polarized targets in high energy physics. *Proceedings of the 1993 Summer Institute on Particle Physics: Spin Structure in High Energy Processes (SSI93)*, SLAC-R-444:185–207, 1993.
- [22] G. D. Cates, S. R. Schaefer, and W. Happer. Relaxation of spins due to field inhomogeneities in gaseous samples at low magnetic fields and low pressures. *Phys. Rev. A*, 37:2877–2885, Apr 1988.
- [23] G. D. Cates, D. J. White, T.-R. Chien, S. R. Schaefer, and W. Happer. Spin relaxation in gases due to inhomogeneous static and oscillating magnetic fields. *Phys. Rev. A*, 38:5092–5106, Nov 1988.
- [24] B. Chann, E. Babcock, L. Anderson, T. Walker, W. Chen, T. Smith, A. Thompson, and T. Gentile. Production of highly polarized ^3he using spectrally narrowed diode laser array bars. *Journal of applied physics*, 94:6908–6914.

- [25] B. Chann, E. Babcock, L. W. Anderson, and T. G. Walker. Measurements of ^3He spin-exchange rates. *Phys. Rev. A*, 66:032703, Sep 2002.
- [26] W. C. Chen, T. R. Gentile, T. G. Walker, and E. Babcock. Spin-exchange optical pumping of ^3He with rb-k mixtures and pure k. *Phys. Rev. A*, 75:013416, Jan 2007.
- [27] T. E. Chupp, M. E. Wagshul, K. P. Coulter, A. B. McDonald, and W. Happer. Polarized, high-density, gaseous ^3He targets. *Phys. Rev. C*, 36:2244–2251, Dec 1987.
- [28] F. D. Colegrove, L. D. Schearer, and G. K. Walters. Polarization of he^3 gas by optical pumping. *Phys. Rev.*, 132:2561–2572, Dec 1963.
- [29] K. Coulter, A. McDonald, W. Happer, T. Chupp, and M. Wagshul. *Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment*, 270:90–94, 1988.
- [30] I. Delstar Metal Finishing. <https://www.delstar.com/electropolishing>.
- [31] A. Deninger, W. Heil, W. E. Otten, M. Wolf, K. R. Kremer, and A. Simon. Paramagnetic relaxation of spin polarized ^3He at coated glass walls. *The European Physical Journal D - Atomic, Molecular, Optical and Plasma Physics*, 38(3):439–443, 2006.
- [32] P. A. M. Dolph, J. Singh, T. Averett, A. Kelleher, K. E. Mooney, V. Nelyubin, W. A. Tobias, B. Wojtsekhowski, and G. D. Cates. Gas dynamics in high-

- luminosity polarized ^3He targets using diffusion and convection. *Phys. Rev. C*, 84:065201, Dec 2011.
- [33] W. A. Fitzsimmons, L. L. Tankersley, and G. K. Walters. Nature of surface-induced nuclear-spin relaxation of gaseous He^3 . *Phys. Rev.*, 179:156–165, Mar 1969.
- [34] J. Frenkel. *Kinetic Theory of Liquids*.
- [35] R. L. Gamblin and T. R. Carver. Polarization and relaxation processes in He^3 gas. *Phys. Rev.*, 138:A946–A960, May 1965.
- [36] L. E. Glass. <http://www.larson-electronic-glass.com>.
- [37] W. Happer, G. Cates, M. Romalis, and C. Erickson. U.s. patent no. 6318092. 2001.
- [38] W. Heil, H. Humblot, E. Otten, M. Schafer, R. Sarkau, and M. Leduc. Very long nuclear relaxation times of spin-polarized ^3He in metal coated cells. *Phys. Rev. A*, 201:337–343, May 1995.
- [39] W. G. Houskeeper. The art of sealing base metals through glass. *Transactions of the American Institute of Electrical Engineers*, XLII:870–877, 1923.
- [40] M. F. Hsu, G. D. Cates, I. Kominis, I. A. Aksay, and D. M. Dabbs. Sol-gel coated glass cells for spin-exchange polarized ^3He . *Applied Physics Letters*, 77(13):2069–2071, 2000.

- [41] M. G. Huber, M. Arif, T. C. Black, W. C. Chen, T. R. Gentile, D. S. Hussey, D. A. Pushin, F. E. Wietfeldt, and L. Yang. Precision measurement of the n-³He incoherent scattering length using neutron interferometry. 102, 2009.
- [42] E. T. Inc. *<http://www.epner.com>*.
- [43] J. Jackson. *Classical Electrodynamics*.
- [44] C. Kittel. *Introduction to Solid State Physics*.
- [45] J. Korringa. Nuclear magnetic relaxation and resonance line shift in metals. *Physica*, 16, 1950.
- [46] B. Larson, O. Hausser, P. P. J. Delheij, D. M. Whittall, and D. Thiessen. Optical pumping of rb in the presence of high-pressure ³He buffer gas. *Phys. Rev. A*, 44:3108–3118, Sep 1991.
- [47] D. Matyas. Characterizing ³He nuclear spin relaxation in vessels of glass and metal. *Master thesis, University of Virginia*, 2016.
- [48] E. Merzbacher. *Quantum Mechanics*. 1998.
- [49] N. R. Newbury, A. S. Barton, P. Bogorad, G. D. Cates, M. Gatzke, B. Saam, L. Han, R. Holmes, P. A. Souder, J. Xu, and D. Benton. Laser polarized muonic helium. *Phys. Rev. Lett.*, 67:3219–3222, Dec 1991.
- [50] N. R. Newbury, A. S. Barton, G. D. Cates, W. Happer, and H. Middleton. Gaseous ³–³He magnetic dipolar spin relaxation. *Phys. Rev. A*, 48:4411–4420, Dec 1993.

- [51] X. Qian, K. Allada, C. Dutta, J. Huang, J. Katich, Y. Wang, Y. Zhang, K. Aniol, J. R. M. Annand, T. Averett, F. Benmokhtar, W. Bertozzi, P. C. Bradshaw, P. Bosted, A. Camsonne, M. Canan, G. D. Cates, C. Chen, J.-P. Chen, W. Chen, K. Chirapatpimol, E. Chudakov, E. Cisbani, J. C. Cornejo, F. Cusanno, M. M. Dalton, W. Deconinck, C. W. de Jager, R. De Leo, X. Deng, A. Deur, H. Ding, P. A. M. Dolph, D. Dutta, L. El Fassi, S. Frullani, H. Gao, F. Garibaldi, D. Gaskell, S. Gilad, R. Gilman, O. Glamazdin, S. Golge, L. Guo, D. Hamilton, O. Hansen, D. W. Higinbotham, T. Holmstrom, M. Huang, H. F. Ibrahim, M. Iodice, X. Jiang, G. Jin, M. K. Jones, A. Kelleher, W. Kim, A. Kolarkar, W. Korsch, J. J. LeRose, X. Li, Y. Li, R. Lindgren, N. Liyanage, E. Long, H.-J. Lu, D. J. Margaziotis, P. Markowitz, S. Marrone, D. McNulty, Z.-E. Meziani, R. Michaels, B. Moffit, C. Muñoz Camacho, S. Nanda, A. Narayan, V. Nelyubin, B. Norum, Y. Oh, M. Osipenko, D. Parno, J. C. Peng, S. K. Phillips, M. Posik, A. J. R. Puckett, Y. Qiang, A. Rakhman, R. D. Ransome, S. Riordan, A. Saha, B. Sawatzky, E. Schulte, A. Shahinyan, M. H. Shabestari, S. Širca, S. Stepanyan, R. Subedi, V. Sulkosky, L.-G. Tang, A. Tobias, G. M. Urciuoli, I. Vilardi, K. Wang, B. Wojtsekhowski, X. Yan, H. Yao, Y. Ye, Z. Ye, L. Yuan, X. Zhan, Y.-W. Zhang, B. Zhao, X. Zheng, L. Zhu, X. Zhu, and X. Zong. Single spin asymmetries in charged pion production from semi-inclusive deep inelastic scattering on a transversely polarized ^3He target at $Q^2 = 1.4\text{--}2.7 \text{ geV}^2$. *Phys. Rev. Lett.*, 107:072003, Aug 2011.
- [52] I. I. Rabi, N. F. Ramsey, and J. Schwinger. Use of rotating coordinates in magnetic resonance problems. *Rev. Mod. Phys.*, 26:167–171, Apr 1954.

- [53] M. L. R.Barbe and F. Laloe. Experimental verifications - measurement of the he3 self-diffusion coefficient. 35:935–951, 1974.
- [54] S. Riordan, S. Abrahamyan, B. Craver, A. Kelleher, A. Kolarkar, J. Miller, G. D. Cates, N. Liyanage, B. Wojtsekhowski, A. Acha, K. Allada, B. Anderson, K. A. Aniol, J. R. M. Annand, J. Arrington, T. Averett, A. Beck, M. Bellis, W. Boeglin, H. Breuer, J. R. Calarco, A. Camsonne, J. P. Chen, E. Chudakov, L. Coman, B. Crowe, F. Cusanno, D. Day, P. Degtyarenko, P. A. M. Dolph, C. Dutta, C. Ferdi, C. Fernández-Ramírez, R. Feuerbach, L. M. Fraile, G. Franklin, S. Frullani, S. Fuchs, F. Garibaldi, N. Gevorgyan, R. Gilman, A. Glamazdin, J. Gomez, K. Grimm, J.-O. Hansen, J. L. Herraiz, D. W. Higinbotham, R. Holmes, T. Holmstrom, D. Howell, C. W. de Jager, X. Jiang, M. K. Jones, J. Katich, L. J. Kaufman, M. Khandaker, J. J. Kelly, D. Kiselev, W. Korsch, J. LeRose, R. Lindgren, P. Markowitz, D. J. Margaziotis, S. M.-T. Beck, S. Mayilyan, K. McCormick, Z.-E. Meziani, R. Michaels, B. Moffit, S. Nanda, V. Nelyubin, T. Ngo, D. M. Nikolenko, B. Norum, L. Pentchev, C. F. Perdrisat, E. Piasetzky, R. Pomatsalyuk, D. Protopopescu, A. J. R. Puckett, V. A. Punjabi, X. Qian, Y. Qiang, B. Quinn, I. Rachek, R. D. Ransome, P. E. Reimer, B. Reitz, J. Roche, G. Ron, O. Rondon, G. Rosner, A. Saha, M. M. Sargsian, B. Sawatzky, J. Segal, M. Shabestari, A. Shahinyan, Y. Shestakov, J. Singh, S. Širca, P. Souder, S. Stepanyan, V. Stibunov, V. Sulkosky, S. Tajima, W. A. Tobias, J. M. Udias, G. M. Urciuoli, B. Vlahovic, H. Voskanyan, K. Wang, F. R. Wesselmann, J. R. Vignote, S. A. Wood, J. Wright, H. Yao, and X. Zhu. Measurements of the electric form factor of the neutron up to $Q^2 = 3.4 \text{ geV}^2$ using

- the reaction ${}^3\text{He}(\vec{e}, e'n)pp$. *Phys. Rev. Lett.*, 105:262302, Dec 2010.
- [55] M. V. Romalis and G. D. Cates. Accurate ${}^3\text{He}$ polarimetry using the rb zee-man frequency shift due to the Rb- ${}^3\text{He}$ spin-exchange collisions. *Phys. Rev. A*, 58:3004–3011, Oct 1998.
- [56] M. V. Romalis, E. Miron, and G. D. Cates. Pressure broadening of rb d_1 and d_2 lines by ${}^3\text{He}$, ${}^4\text{He}$, N_2 , and Xe : line cores and near wings. *Phys. Rev. A*, 56(6), 1997.
- [57] L. D. Schearer, F. D. Colegrove, and G. K. Walters. Large He^3 nuclear polarization. *Phys. Rev. Lett.*, 10:108–110, Feb 1963.
- [58] L. D. Schearer and G. K. Walters. Nuclear spin-lattice relaxation in the presence of magnetic-field gradients. *Phys. Rev.*, 139:A1398–A1402, Aug 1965.
- [59] J. Schmiedeskamp, W. Heil, W. E. Otten, K. R. Kremer, A. Simon, and J. Zimmer. Paramagnetic relaxation of spin polarized ${}^3\text{He}$ at bare glass surfaces. *The European Physical Journal D - Atomic, Molecular, Optical and Plasma Physics*, 38(3):427–438, 2006.
- [60] J. T. Singh, P. A. M. Dolph, W. A. Tobias, T. D. Averett, A. Kelleher, K. E. Mooney, V. V. Nelyubin, Y. Wang, Y. Zheng, and G. D. Cates. Development of high-performance alkali-hybrid polarized ${}^3\text{He}$ targets for electron scattering. *Phys. Rev. C*, 91:055205, May 2015.
- [61] C. P. Slichter. *Principles of Magnetic Resonance*.
- [62] W. Smythe. *Static and Dynamic Electricity*.

- [63] A. E. A. M. I. Technologies. <http://www.ableelectropolishing.com>.
- [64] T. V. Tscherbul, P. Zhang, H. R. Sadeghpour, and A. Dalgarno. Anisotropic hyperfine interactions limit the efficiency of spin-exchange optical pumping of ^3He nuclei. *Phys. Rev. Lett.*, 107:023204, Jul 2011.
- [65] M. Wagshul and T. Chupp. Optical pumping of high-density rb with a broadband dye laser and gaalas diode laser arrays: Application to ^3he polarization. *Phys. Rev. A.*, 40, 1989.
- [66] M. Wagshul and T. Chupp. Laser optical pumping of high-density rb in polarized ^3he targets. *Phys. Rev. A*, 49:3854–3869, 1994.
- [67] T. G. Walker and W. Happer. Spin-exchange optical pumping of noble-gas nuclei. *Rev. Mod. Phys.*, 69:629–642, Apr 1997.
- [68] T. G. Walker, I. A. Nelson, and S. Kadlecik. Method for deducing anisotropic spin-exchange rates. *Phys. Rev. A*, 81:032709, Mar 2010.
- [69] T. G. Walker, J. H. Thywissen, and W. Happer. Spin-rotation interaction of alkali-metal \sim he-atom pairs. *Phys. Rev. A*, 56:2090–2094, Sep 1997.
- [70] D. K. Walter, W. Happer, and T. G. Walker. Estimates of the relative magnitudes of the isotropic and anisotropic magnetic-dipole hyperfine interactions in alkali-metal \sim noble-gas systems. *Phys. Rev. A*, 58:3642–3653, Nov 1998.
- [71] Y. Zheng. Low field mri and the development of polarized nuclear imaging (pni)-a new imaging modality, 2015.