

Chapter 1

Introduction

Nuclear-polarized noble gases have been proven to be very useful in various applications, such as polarized targets for electron scattering experiments [9], magnetic resonance imaging [7] and neutron scattering experiments [41]. Polarized ^3He has been particularly useful for studying spin-dependent interactions involving neutrons because, to first-order approximation, a ^3He nucleus has a pair of protons with paired spins and a single neutron that carries the most of the nuclear spin. Free neutrons are not used as targets because they decay with a lifetime of about 14 minutes, 42 seconds.

1.1 Proposed JLab Experiments

The neutron electromagnetic form factors, G_E^n and G_M^n , play essential roles for understanding nucleon structure. They are the Fourier transforms of the electric charge and magnetic moment distributions at non-relative energies. However, this interpretation suffers in that the momentum transfers are too large to ignore relativistic

Even at relativistic energies, the elastic form factors provide unique information on the transverse structure of the nucleon [Miller paper, Carlson].

Why are these ref. #'s not in order?

can be viewed as

Double-polarization experiments on the proton showed that

nearly linearly

effects. Recent results on proton show that the ratio G_E^p/G_M^p declines sharply as Q^2 increases. We expect the mechanism causing this behavior to affect neutron in similar way, but our knowledge of G_E^n at high Q^2 is poor compared to that of proton. Double polarization experiments provide an important tool to study G_E^n . The planned experiments are enabled by both the equipments including the large acceptance Super BigBite spectrometer and the high degree of neutron polarization of ^3He targets with luminosity that exceeds other polarized targets.

Current understanding of the nucleon spin is that the total spin is composed of contributions from valence quarks, $q\bar{q}$ sea quarks, their orbital angular momenta, and gluons. To further study the nucleon spin, it is important to know how it decomposes into the three components in the full x and Q^2 region and to understand data from stand model of strong interactions - Quantum Chromodynamics (QCD). However, it is extremely difficult to make absolute predictions from QCD on the nucleon spin. The neutron asymmetry A_1^n can be related to the unpolarized and the polarized structure functions, which have explicit implication in the quark-parton model. Precision measurements of A_1^n in the deep inelastic scattering region $0.3 \leq x \leq 0.77$ and $3 \leq Q^2 \leq 10 \text{ (GeV/c)}^2$ has been proposed. Polarized ^3He targets, the high momentum spectrometer (HMS) and the planned Super HMS (SHMS) will be used to explore possible Q^2 -dependence of A_1^n . This measurement will provide the first precision data in the region above $x = 0.6$ and test various predictions.

See
email.

1.2 Overview of Recent Target Development

An early use of polarized ^3He target³ in electron scattering experiment was at Stanford Linear Accelerator Center (SLAC) in the year of 1992. The experiment was known as E142 and investigated the spin structure of neutrons [9]. Recent experiments were conducted at Jefferson Laboratory (JLAB) in Newport News, Virginia, such as the aforementioned E02-013, also known as "Measurement of the Neutron Electric Form Factor G_E^n at High Q^2 ", also known as E02-013 [54]. E06-10, E06-014 and E05-015 were experiments that investigated single-spin asymmetries [29]. included

The ^3He targets used in these experiments were polarized with the technique Spin-Exchange Optical Pumping (SEOP). Fig. 1.1 shows schematically what a typical target looks like, they are also referred to as "target cells" very often. These cells were made of the GE180 glass and use a two-chambered design. The top chamber, known as the pumping chamber, is where ^3He is polarized through SEOP. The bottom chamber, known as the target chamber, is where electron scattering occurs. The two ends of the target chamber where electron beam enters and exits are known as the "end windows". Great effort has been made in our lab to develop this generation of cells. Alkali-hybrid SEOP together with narrowband laser diode arrays have increased the ^3He polarization from 37% to 70%. Among other things, we also carefully studied an additional spin relaxation mechanism that limits the maximum achievable ^3He polarization, which is referred to as the "X Factor". Analysis of data accumulated through developing this generation of target cells were thoroughly discussed in Ref. [60], part of which will be presented in chapter 4.




Figure 1.1: A schematic representation of a target cell. The dimensions of different parts of the cell are not to scale.

1.3 New Generation Target Cells

The future experiments planned during the 12GeV era after the upgrade will be much more demanding in terms of target cell performance. In particular, there is a desire to run experiments with higher luminosity where luminosity is the product of gas density in the target, interaction length and beam current. Increased luminosity will lead to more interactions that depolarize the target. We have designed and tested a new style cell that utilizes convection instead of diffusion to increase the rate at which the polarization in the target chamber is replenished by polarized gas from pumping chamber [32]. We have obtained over 50% polarization with controllable convection

speed so far.

An additional problem that comes with higher beam current is that the glass end windows of traditional design are not likely to survive the experiments. Our group started exploring the option of using metal end windows since eight years ago. Fig ?? shows an example configuration of such a target. The first problem to solve ~~is to~~ find out the correct material and the proper technique to incorporate metal without introducing significant spin relaxation and still being able to hold high pressure gas (12 atm) inside. This is a brand new technique that may have a profound impact on future cell designs once fully developed. Although no metal end windows have been tested so far, through carefully examining multiple glass cells with different kinds metal tubes (much larger in area compared to the end windows that will be used in JLAB experiments) attached, we have developed a reliable way of incorporating metal to target cells without introducing significant spin relaxation rates. We believe the next generation target cells used in the 12GeV era will be able to utilize metal end windows. In ~~these~~ ^{our} test cells, the metals tubes were connected to Pyrex glass with Houskeeper seals [39] and stayed intact through high pressure tests. After exploring options such as pure copper, gold coated copper, titanium, stainless steel, gold coated titanium, we have established that electroplating gold on copper substrate yields the best result so far. ~~We~~ ^{we} have achieved a 15.6 h relaxation time with a Pyrex cell that had a 5" long by 1" gold coated copper tube attached horizontally. The additional relaxation rate introduced by the metal surface is proportional to the area of the gold surface. By comparing relaxation rate ^s of test metal cells with pure-glass control cells, the relaxation rate due to ~~gold~~ ^{the} surface was extrapolated. With this result, we believe the relaxation rate introduced by small metal windows in a target will be less than




Figure 1.2: A diagram of convection style target cell with metal end windows.

$1/135 \text{ hr}^{-1}$. To the best of our knowledge, our group was the first to have proved the potential of incorporating metal to target cells in the presence of alkali vapor.

need
to make
consistent
with
conclusion.

1.4 Structure of This Thesis

This thesis focuses on both discussion on the development of high-performance polarized ^3He targets that utilize spin-exchange optical pumping (SEOP) and the development of future target cells that incorporate metal end windows. Chapter 2 gives a general description of SEOP. Chapter 3 introduces polarimetry techniques used in our lab for target cell characterization. Chapter 4 discusses the result collected in our lab from the over-a-decade development of ^3He target cells, in which the spin-exchange rate constant for K and ^3He is calculated and the so-called "X Factor" is studied. Chapter 5 presents the development process of target cells with metal parts that aims to incorporate metal end windows to future cells for the 12 GeV era