

AN APPROXIMATION METHOD TO PRICE VIX OPTIONS

by

You Wang

A THESIS

Submitted to the Faculty of the Stevens Institute of Technology
in partial fulfillment of the requirements for the degree of

MASTER OF SCIENCE - FINANCIAL ENGINEERING

You Wang, Candidate

ADVISORY COMMITTEE

Zhenyu Cui, Advisor Date

Ionut Florescu, Reader Date

STEVENS INSTITUTE OF TECHNOLOGY

Castle Point on Hudson

Hoboken, NJ 07030

2021

AN APPROXIMATION METHOD TO PRICE VIX OPTIONS

ABSTRACT

We propose an approximation method to value VIX options. This method is based on choosing models with closed form solution as an auxiliary model, and derive a mis-pricing formula between the true price and the auxiliary one, then apply Ito-Taylor expansions on the mis-pricing formula to create increasingly improved refinements. We propose an approach to evaluate VIX options under mean-reverting models, in which auxiliary model selection and expansion methods are explained. Method in this paper is applied to mean-reverting Constant elasticity of variance(CEV) model and double CEV models. Numerical results show that the proposed method is accurate and efficient.

Author: You Wang

Advisor: Zhenyu Cui

Date: Nov 17, 2021

Department: Financial Engineering

Degree: Master of Science - Financial Engineering

Acknowledgments

The acknowledgements section recognizes anyone that provided significant help in producing your thesis or dissertation. Frequently acknowledged people are your advisor, colleagues, and family. Sometimes companies or outside groups have contributed to the research done for a dissertation, and they can be thanked here as well. The Acknowledgements page is optional.

Table of Contents

Abstract	iii
Dedication	iv
Acknowledgments	iv
List of Tables	vi
List of Figures	vii
1 Introduction	1
2 Method Description	2
2.1 The DOI Variance Reduction Method	2
2.2 Approximation Method based on the DOI method	5
3 Approximations of VIX options	7
3.1 Approximating options under mean-reverting CEV model	7
3.1.1 Drawbacks of using Black-Scholes model as an auxiliary model	7
3.1.2 Using square root mean-reverting model as auxiliary model	9
3.1.3 Derivatives of non-central chi-square distribution function	11
3.2 Approximating options under double Heston model	15
A Appendix A Title	1
Bibliography	2

List of Tables

List of Figures

3.1	Call option price with regard to time to maturity	8
3.2	Call option price with regard to volatility	9

Chapter 1

Introduction

Intro to be added

Chapter 2

Method Description

In this section, the origin DOI method, JDOI method and approximation method based on DOI method are described. In section 2.1,, we introduce the origin DOI method, seeHeath and Platen (Heath and Platen). In section 2.2, we illustrate the approximation method proposed by Kristensen and Mele (2011). In section ??, we discuss our estimator to price American options based on JDOI method, Auster et al. (2021).

2.1 The DOI Variance Reduction Method

Consider a multi-factor model, in which a d -dimensional vector of state variables $X(t)$ on a filtered probability space $(\Omega, \mathcal{F}, \mathbb{Q})$ satisfies the following Stochastic Differential Equations(SDEs)

$$dX(t) = \mu(t, X(t))dt + \sigma(t, X(t))dW(t) \quad (2.1.1)$$

where $\mu(t, X(t))$ and $\sigma(t, X(t))$ are drift and diffusion functions under the risk-neutral measure \mathbb{Q} , which also satisfies appropriate growth and Lipschitz conditions such that equation(2.1.1) admits a unique strong solution and is Markovian; $W(t)$ is a d -dimensional standard Brownian Motion and $t \in [0, T]$.

Let $w(t, x)$ be the value function of European option written on $X(T)$ with current state $X(t) = x$, $G(t, x)$ be the payoff function. We define the infinitesimal generator \mathcal{L} associated with equation(2.1.1)to be

$$\mathcal{L}w(t, x) = \frac{\partial w}{\partial t} + \sum_{i=1}^d \mu_i(t, x) \frac{\partial w}{\partial x_i} + \frac{1}{2} \sum_{i=1}^d \sum_{j=1}^d (\sigma(t, x) \sigma^\top(t, x))_{i,j} \frac{\partial^2 w}{\partial x_i \partial x_j} \quad (2.1.2)$$

Let $R(t, x)$ be the instantaneous short-term interest rate, combining with equation(2.1.1) and equation(2.1.2), the price of European option V is a solution to the following partial differential equation(PDE)

$$\mathcal{L}w(x, t) = R(x, t)w(x, t) \quad (2.1.3)$$

with boundary condition $w(T, x(T)) = G(T, X(T))$. It's easily seen that under risk neutral measure \mathbb{Q} , the instantaneous option price change is equal to the price gain in saving account.

Next we consider to use a m -dimensional($m \leq d$) process $\bar{X}(t)$ which is a simpler auxiliary model to approximate the price of option. $\bar{X}(t)$ satisfies the following SDE

$$d\bar{X}(t) = \begin{cases} \bar{\mu}_i(t, \bar{X}(t))dt + \bar{\sigma}_i(t, \bar{X}(t))dW(t) & 1 \leq i \leq m \\ \bar{\mu}_i(t, \bar{X}(t)) = 0, \bar{\sigma}_i(t, \bar{X}(t)) = 0 & m < i \leq d \end{cases} \quad (2.1.4)$$

where $\bar{\mu}(t, \bar{X}(t))$ and $\bar{\sigma}(t, \bar{X}(t))$ are drift and diffusion functions, and they are also assumed to satisfy appropriate conditions such that equation(2.1.4) admits a unique strong solution and is Markovian.

Let $\bar{w}(t, x)$ be the option price written on process $\bar{X}(t)$ and assume \bar{w} has closed form solution under this new process, the infinitesimal generator $\bar{\mathcal{L}}$ for option price \bar{w} is the same as equation(2.1.2) but replacing $\mu(t, x)$, $\sigma(t, x)$ by $\bar{\mu}(t, x)$ and $\bar{\sigma}(t, x)$. Therefore $\bar{w}(t, x)$ is a solution to

$$\bar{\mathcal{L}}\bar{w}(x, t) = R(x, t)\bar{w}(x, t) \quad (2.1.5)$$

Denote the price difference $\Delta w(t, x) = w(t, x) - \bar{w}(t, x)$, by subtract equation(2.1.5 from equation(2.1.3)), $\Delta w(t, x)$ satisfies the following equation

$$\mathcal{L}\Delta w(t, x) + (\mathcal{L} - \bar{\mathcal{L}})w(t, x) = R(x, t)\Delta w(t, x) \quad (2.1.6)$$

with boundary condition $\Delta w(T, x) = G(T, x) - \bar{G}(T, x)$. We can find that the price difference arises from two parts:

- The use of a wrong payoff function $\bar{G}(t, x)$, it can be eliminated once we use the same payoff function in auxiliary model as it in general model
- The discrepancies between the auxiliary model and general model.

Define $\delta(t, x) = (\mathcal{L} - \bar{\mathcal{L}})w(t, x)$, $d(t, x) = G(t, x) - \bar{G}(t, x)$, under standard regularity conditions¹, we can derive the following formula by using the Feynman-Kac representation.

$$\begin{aligned} w(t, x) = & \bar{w}(t, x) + \mathbb{E}_{t,x} \left[\exp \left(- \int_t^T R(s, X(s)) ds \right) d(T, X(T)) \right] \\ & + \int_t^T \mathbb{E}_{t,x} \left[\exp \left(- \int_t^s R(u, X(u)) du \right) \delta(s, X(s)) \right] ds \end{aligned} \quad (2.1.7)$$

Finally, under the initial condition $Z_0 = w(0, x)$, the DOI estimator

$$\begin{aligned} Z_t = & \bar{w}(t, x) + \exp \left(- \int_t^T R(s, X(s)) ds \right) d(T, X(T)) \\ & + \int_t^T \exp \left(- \int_t^s R(x(u), u) du \right) \delta(s, X(s)) ds \end{aligned} \quad (2.1.8)$$

¹See Appendix A in Kristensen and Mele (2011)

is an unbiased estimator for Z_0 . And if a good auxiliary model is chosen, the variance of Z_t will be small.

2.2 Approximation Method based on the DOI method

Recall equation(2.1.7), instead of using it as an estimator to do simulations, Kristensen and Mele (2011) make some additional assumptions and use Ito-Taylor expansion to get closed form approximation formula.

For sufficiently smooth function $f(t, x)$, Ito-Taylor expansion is given by

$$\mathbb{E}^{t,x}[f(s, X(s))] = \sum_{N=0}^J \frac{(s-t)^N}{N!} (\mathcal{L})^N f(t, x) + \mathcal{R}_J \quad (2.2.1)$$

where the remainder term \mathcal{R}_J is given by

$$\mathcal{R}_J = \mathbb{E}^{t,x} \left[\int_t^s du_1 \int_t^{u_1} du_2 \cdots \int_t^{u_J} (\mathcal{L})^{J+1} f(u_{J+1}, X(u_{J+1})) du_{J+1} \right] \quad (2.2.2)$$

The process $X(t)$ here is defined in equation(2.1.1), and the infinitesimal generator \mathcal{L} is defined in equation(2.1.2).

Assume closed form solution of option price \bar{V} under auxiliary model and the difference of payoff function $d(t, x)$ is sufficiently smooth. In other words, for $N \geq 1$, assume $\delta(t, x)$ and $d(t, x)$ to be $2N$ times differentiable with respect to x , $\delta(t, x)$ to be N times differentiable with respect to t . By applying Ito-Taylor expansion to equation(2.1.7)

$$\begin{aligned}
V(t, x) = & \bar{V}(t, x) + \mathbb{E}_{t,x} \left[\exp \left(- \int_t^T R(s, X(s)) ds \right) d(T, X(T)) \right] \\
& + \int_t^T \mathbb{E}_{t,x} \left[\exp \left(- \int_t^s R(u, X(u)) du \right) \delta(s, X(s)) \right] ds
\end{aligned} \tag{2.2.3}$$

We can get a closed-form approximation formula

$$V_N(t, x) = \bar{V}(t, x) + \sum_{n=0}^N \frac{(T-t)^n}{n!} d_n(t, x) + \sum_{n=0}^N \frac{(T-t)^{n+1}}{(n+1)!} \delta_n(t, x) \tag{2.2.4}$$

where $d_0(t, x) = d(x)$, $\delta_0(t, x) = \delta(t, x)$, and

$$\begin{aligned}
d_n(t, x) &= L d_{n-1}(t, x) - R(t, x) d_{n-1}(t, x) \\
\delta_n(t, x) &= L \delta_{n-1}(t, x) - R(t, x) \delta_{n-1}(t, x)
\end{aligned} \tag{2.2.5}$$

Note that the terms in equation((2.2.4)) can be calculated once for all, meaning that it be computed much faster than simulation methods using estimator.

Chapter 3

Approximations of VIX options

3.1 Approximating options under mean-reverting CEV model

3.1.1 Drawbacks of using Black-Scholes model as an auxiliary model

Chan et al. (1992) proposes the mean-reversion CEV model, in which volatility follows

$$dV_t = (\alpha + \beta V_t) dt + \sigma V_t^\gamma dW_t$$

when β is negative, this model has mean-reverting property. We can rewrite it to be

$$dV_t = \kappa(m - V_t)dt + \sigma V_t^\gamma dW_t \quad (3.1.1)$$

where κ is the speed of mean-reversion, m is the long-run mean. A natural idea is to use Black-Scholes model as auxiliary model as mentioned in Kristensen and Mele (2011), then apply their method to approximate the VIX option price under mean-reverting CEV model. Denote \mathcal{L} and \mathcal{L}^{BS} to be infinitesimal generators of mean-reverting CEV model and Black-Scholes model respectively

$$\begin{aligned} \mathcal{L}w &= \frac{\partial w}{\partial t} + \kappa(m - V) \frac{\partial w}{\partial V} + \frac{1}{2} \sigma^2 V^{2\gamma} \frac{\partial^2 w}{\partial V^2} \\ \mathcal{L}^{\text{BS}}w &= \frac{\partial w}{\partial t} + rV \frac{\partial w}{\partial V} + \frac{1}{2} \sigma^2 V^2 \frac{\partial^2 w}{\partial V^2} \end{aligned}$$

The mis-pricing term for using Black-Scholes model is then

$$\delta^{\text{BS}} = (\mathcal{L} - \mathcal{L}^{\text{BS}})w^{\text{BS}} = (\kappa - r)V \frac{\partial w^{\text{BS}}}{\partial t} + \kappa m \frac{\partial w^{\text{BS}}}{\partial V} + \sigma^2 (V - V^{2\gamma}) \frac{\partial^2 w^{\text{BS}}}{\partial V^2}$$

with the solution of Black-Scholes model w^{BS} . Note that δ^{BS} contains theta and gamma of option. Their differences in Black-Scholes model and mean-reverting model determines that we have to use other auxiliary models.

Take call option prices under $\gamma = \frac{1}{2}$ in model(3.1.1) as an example. This model is known as mean square root mean-reverting model proposed by Grünbichler and Longstaff (1996).




Figure 3.1: Call option price with regard to time to maturity

From figure 3.1, we can find that in contrast to Black-Scholes model, the value of call option price under mean-reverting model is not always increasing as time to maturity increases; From figure 3.2, by contrast, the call option price does not converge to zero as volatility goes to zero. In addition, Grünbichler and Longstaff (1996) also shows that V has less influence of the current value of the call option than in Black-Scholes model. For these reasons, we conclude that Black-Scholes model is not an appropriate auxiliary model and in the next section, we discuss that using the square root mean-reverting model as the auxiliary model.




Figure 3.2: Call option price with regard to volatility

3.1.2 Using square root mean-reverting model as auxiliary model

Recall the mean-reverting CEV model with $\gamma = \frac{1}{2}$

$$dV_t = \kappa(m - V_t)dt + \sigma\sqrt{V_t}dW_t \quad (3.1.2)$$

We are going to use it as our auxiliary model as it captures the mean-reverting property of general mean-reverting CEV models. Grünbichler and Longstaff (1996) gives an explicit solution to this model. Denote the call option price \bar{w} with strike K , constant risk-free rate r , time to maturity T and no expected premium for volatility risk is paid, its price is given by

$$\begin{aligned} \bar{w} = & e^{-(\kappa+r)T} V Q(xK; \nu + 4, \lambda) \\ & + m e^{-rT} (1 - e^{-\kappa T}) Q(xK; \nu + 2, \lambda) \\ & - e^{-rT} K Q(xK; \nu, \lambda) \end{aligned} \quad (3.1.3)$$

where

$$\begin{aligned}
x &= \frac{4\kappa}{\sigma^2(1 - e^{-\kappa T})} \\
\nu &= \frac{4\kappa m}{\sigma^2}, \\
\lambda &= e^{-\kappa T} x V
\end{aligned}$$

and $Q(xK; \nu + i, \lambda)$ is the complementary distribution function for the non-central chi-squared density with $\nu + i$ degrees of freedom and non-centrality parameter λ .

Define the infinitesimal generators $\bar{\mathcal{L}}$ for square root mean-reverting model and \mathcal{L} for mean-reverting CEV model

$$\begin{aligned}
\mathcal{L}w &= \frac{\partial w}{\partial t} + \kappa(m - V)\frac{\partial w}{\partial V} + \frac{1}{2}\sigma^2 V^{2\gamma} \frac{\partial^2 w}{\partial V^2} \\
\bar{\mathcal{L}}w &= \frac{\partial w}{\partial t} + \kappa(m - V)\frac{\partial w}{\partial V} + \frac{1}{2}\sigma^2 V \frac{\partial^2 w}{\partial V^2}
\end{aligned} \tag{3.1.4}$$

Subtract infinitesimal generators in equation(3.1.4), we get the mis-pricing formula for using square root mean-reverting model

$$\delta = (\mathcal{L} - \bar{\mathcal{L}})\bar{w} = \frac{1}{2}\sigma^2(V^{2\gamma} - V)\frac{\partial^2 w}{\partial V^2}$$

We can then use the approximation formula discussed in 2.2 to price call options under mean-reverting CEV model¹

$$w_N(t, x) = \bar{w}(t, x) + \sum_{n=0}^N \frac{(T - t)^{n+1}}{(n + 1)!} \delta_n(t, x) \tag{3.1.5}$$

where

$$\begin{aligned}
\delta_0 &= \delta = \frac{1}{2}\sigma^2(V^{2\gamma} - V)\frac{\partial^2 w}{\partial V^2} \\
\delta_n(t, x) &= L\delta_{n-1}(t, x) - r\delta_{n-1}(t, x)
\end{aligned} \tag{3.1.6}$$

¹Put options can be priced easily in the same way

Finally we get a closed form approximating formula for call options under mean-reverting CEV model. But notice that the call price (3.1.3) contains non-square chi square distribution functions, applying infinitesimal generator \mathcal{L} on it can be a hard point and in the next section we are going to talk about how to derive partial derivatives of distribution function $Q(\cdot|\nu + i, \lambda)$.

3.1.3 Derivatives of non-central chi-square distribution function

In this section, methods² are given to calculate the derivatives of call option price \bar{w} to time t and volatility index V . We have already discussed in 3.1.3 the call option prices \bar{w} under square root mean-reverting model, where

$$\begin{aligned} xK &= \frac{4\kappa}{\sigma^2(1 - e^{-\kappa T})}K \\ \lambda &= xe^{-\kappa T}V = \frac{4\kappa e^{-\kappa T}}{\sigma^2(1 - e^{-\kappa T})}V \end{aligned} \quad (3.1.7)$$

Based on equation(3.1.7), we can calculate partial derivatives of x and λ with regard to V and t

$$\begin{aligned} \frac{\partial(xK)}{\partial V} &= 0 \\ \frac{\partial\lambda}{\partial V} &= xe^{-\kappa T} = \frac{4\kappa e^{-\kappa T}}{\sigma^2(1 - e^{-\kappa T})} \end{aligned} \quad (3.1.8)$$

$$\begin{aligned} \frac{\partial(xK)}{\partial t} &= \frac{-\kappa e^{-\kappa T}}{1 - e^{-\kappa T}} \cdot xK \\ \frac{\partial\lambda}{\partial t} &= \frac{-\kappa e^{-\kappa T}}{1 - e^{-\kappa T}} \cdot xV \end{aligned} \quad (3.1.9)$$

Besides, using the relationship between the complementary cumulative distribution function(ccdf) $Q(xK; \nu + i, \lambda)$, the cumulative distribution function(cdf)

²Methods are inspired by Hossain et al. (2019), who use to calculate the Greeks of options under CEV model

$F(xK; \nu + i, \lambda)$ and the probability density function(pdf) $p(xK; \nu + i, \lambda)$ we can derive

$$\begin{aligned} \frac{\partial Q(xK; \nu, \lambda)}{\partial x} &= \frac{\partial[1 - F(xK; \nu, \lambda)]}{\partial x} \\ &= -\frac{\partial F(xK; \nu, \lambda)}{\partial x} \\ &= -p(xK; \nu, \lambda) \end{aligned} \quad (3.1.10)$$

$$\begin{aligned} \frac{\partial Q(xK; \nu, \lambda)}{\partial \lambda} &= \frac{\partial[1 - F(xK; \nu, \lambda)]}{\partial \lambda} \\ &= -\frac{\partial F(xK; \nu, \lambda)}{\partial \lambda} \\ &= -p(xK; \nu + 2, \lambda) \end{aligned} \quad (3.1.11)$$

Combining equation(3.1.9), (3.1.10), (3.1.11) to use chain rule we can get partial derivative of $Q(xK; \nu, \lambda)$ to time t

$$\begin{aligned} \frac{\partial Q(xK; \nu, \lambda)}{\partial t} &= \frac{\partial Q}{\partial(xK)} \frac{\partial(xK)}{\partial t} + \frac{\partial Q}{\partial \lambda} \frac{\partial \lambda}{\partial t} \\ &= p(xK; \nu, \lambda) \cdot \frac{\kappa e^{-\kappa T}}{1 - e^{-\kappa T}} \cdot xK + p(x; \nu + 2, \lambda) \cdot \frac{\kappa e^{-\kappa T}}{1 - e^{-\kappa T}} \cdot xV \quad (3.1.12) \\ &= \frac{\kappa x e^{-\kappa T}}{1 - e^{-\kappa T}} (Kp(x, \nu, \lambda) + Vp(x; \nu + 2, \lambda)) \end{aligned}$$

Similarly combining equation(3.1.8), (3.1.10), (3.1.11) we can get partial derivative of $Q(w, \nu, \lambda)$ to volatility V

$$\begin{aligned} \frac{\partial Q(xK; \nu, \lambda)}{\partial V} &= \frac{\partial Q}{\partial x} \frac{\partial x}{\partial V} + \frac{\partial Q}{\partial \lambda} \frac{\partial \lambda}{\partial V} \\ &= -x e^{-\kappa T} p(x; \nu + 2, \lambda) \end{aligned} \quad (3.1.13)$$

Thus we can calculate delta of call option

$$\begin{aligned}
\Delta_{\bar{w}} &= \frac{\partial \bar{w}}{\partial V} \\
&= e^{-(\kappa+r)T} [Q(xK; \nu + 4, \lambda) - xV e^{-\kappa T} p(xK; \nu + 6, \lambda) \\
&\quad - mx(1 - e^{-\kappa T})p(xK; \nu + 4, \lambda) + xKp(xK; \nu + 2, \lambda)]
\end{aligned} \tag{3.1.14}$$

As is seen that pdf in previous equation are all with 2 higher degrees of freedom. However, pdf with higher degrees of freedom require us to re-calculate pdf; Besides, result of approximating pdf with higher degrees of freedom may be inaccurate. To let degrees of freedom λ of pdf bounded in some range, we consider the following relationship proposed by Cohen (1988)

$$p(xK; \nu + 2, \lambda) = \frac{xK}{\lambda} p(xK; \nu - 2, \lambda) - \frac{\nu - 2}{\lambda} p(xK; \nu, \lambda) \tag{3.1.15}$$

Using (3.1.15) we can rewrite $\Delta_{\bar{w}}$ as

$$\begin{aligned}
\Delta_{\bar{w}} &= \frac{\partial \bar{w}}{\partial V} \\
&= e^{-(\kappa+r)T} [Q(xK; \nu + 4, \lambda) \\
&\quad - xV e^{-\kappa T} (\frac{K}{e^{-\kappa T} V} p(xK; \nu + 2, \lambda) - \frac{\nu + 2}{\lambda} p(xK; \nu + 4, \lambda)) \\
&\quad - mx(1 - e^{-\kappa T})p(xK; \nu + 4, \lambda) + xKp(xK; \nu + 2, \lambda)] \\
&= e^{-(\kappa+r)T} [Q(xK; \nu + 4, \lambda) - mx(1 - e^{-\kappa T})p(xK; \nu + 4, \lambda) + (\nu + 2)p(xK; \nu + 4, \lambda)]
\end{aligned} \tag{3.1.16}$$

Next we calculate the partial derivatives of pdf $p(x; \nu, \lambda)$ to V and t in order to calculate gamma and apply infinitesimal generator on the mis-pricing formula. Cohen (1988) proposes the recurrence relations for p , it's also been used in Hossain et al.

(2019) to calculate the Greeks of options under CEV models, which is given by

$$\begin{aligned}\frac{\partial p(xK; \nu, \lambda)}{\partial \lambda} &= \frac{1}{2}[-p(xK; v, \lambda) + p(xK; v + 2, \lambda)] \\ \frac{\partial p(xK; \nu, \lambda)}{\partial(xK)} &= \frac{1}{2}[-p(xK; v, \lambda) + p(xK; v - 2, \lambda)]\end{aligned}\tag{3.1.17}$$

Combining it with equation(3.1.8), (3.1.9) gives

$$\begin{aligned}\frac{\partial p(xK; \nu, \lambda)}{\partial t} &= \frac{\partial p}{\partial(xK)} \frac{\partial(xK)}{\partial t} + \frac{\partial p}{\partial \lambda} \frac{\partial \lambda}{\partial t} \\ &= \frac{-\kappa x e^{-\kappa T}}{2(1 - e^{-\kappa T})} [Vp(xK; v + 2, \lambda) - (K + V)p(xK; v, \lambda) + Kp(xK; v - 2, \lambda)] \\ \frac{\partial p}{\partial V} &= \frac{\partial p}{\partial(xK)} \frac{\partial(xK)}{\partial V} + \frac{\partial p}{\partial \lambda} \frac{\partial \lambda}{\partial V} \\ &= \frac{x e^{-\kappa T}}{2} [-p(xK; v, \lambda) + p(xK; v + 2, \lambda)]\end{aligned}\tag{3.1.18}$$

Besides, by using the transformation of (3.1.15)

$$p(xK; \nu - 2, \lambda) = \frac{\lambda}{xK} p(xK; \nu + 2, \lambda) + \frac{v - 2}{xK} p(xK; \nu, \lambda)\tag{3.1.19}$$

$p(xK; \nu - 2, \lambda)$ in (3.1.18) can be substituted by

$$\begin{aligned}\frac{\partial p(xK; \nu, \lambda)}{\partial(xK)} &= \frac{1}{2}[-p(xK; v, \lambda) + p(xK; v - 2, \lambda)] \\ &= \frac{1}{2} \left[\left(\frac{v - 2}{xK} - 1 \right) p(xK; \nu, \lambda) + \frac{\lambda}{xK} p(xK; \nu + 2, \lambda) \right]\end{aligned}\tag{3.1.20}$$

Thus gamma of call option under square root mean-reverting model is then

$$\begin{aligned}
\Gamma_{\bar{w}} &= \frac{\partial \Delta_{\bar{w}}}{\partial V} \\
&= \frac{x e^{-(2\kappa+r)T}}{2} [m x (1 - e^{-\kappa T}) (p(xK; \nu + 4, \lambda) - p(xK + 6; \nu, \lambda)) \\
&\quad + \nu p(xK + 6; \nu, \lambda) - (\nu + 2) p(xK + 4; \nu, \lambda)]
\end{aligned} \tag{3.1.21}$$

From (3.1.6) we know that $\delta = \frac{1}{2}\sigma^2(V^{2\gamma} - V)\Gamma_{\bar{w}}$, all terms have been solved from above. In essence, during expansion, we can solve partial derivatives of cdf $Q(xK; \nu, \lambda)$ and pdf $p(xK; \nu, \lambda)$ explicitly. And for those pdf with degrees out of bounds, we use (3.1.19) and (3.1.15) to adjust them, making degrees of freedom within our desired range. Finally we can get a closed-form approximation formula for all mean-reverting CEV model. The expansions in approximating formula can be computed once for all, we can solve it manually or use symbolic language.

3.2 Approximating options under double Heston model

Gatheral (2008) proposes volatility with double mean-reverting dynamics

$$\begin{aligned}
dV_t &= -\kappa (V_t - V'(t)) dt + \eta_1 V_t'^{\alpha} dW_1(t) \\
dV_t' &= -c (V_t' - m) dt + \eta_2 V_t'^{\beta} dW_2(t)
\end{aligned}$$

where $\alpha, \beta \in [\frac{1}{2}, 1]$.

- It's called Double Heston model in the case $\alpha = \beta = \frac{1}{2}$.
- The case $\alpha = \beta = 1$ Double Log-normal model.
- And the general Double CEV model.

From our previous work we can price options with V_t following heston dynamics and V' following any mean-reverting CEV process, the model is thus

$$\begin{aligned}
dV_t &= -\kappa(V_t - V'(t))dt + \eta_1 \sqrt{V_t} dW_1(t) \\
dV'_t &= -c(V'_t - m)dt + \eta_2 V'^{\beta}_t dW_2(t)
\end{aligned} \tag{3.2.1}$$

Define infinitesimal generator \mathcal{L} for (3.2.1) and we still use square root mean-reverting model as our auxiliary model

$$\begin{aligned}
\mathcal{L}w &= \frac{\partial w}{\partial t} + \kappa(V' - V)\frac{\partial w}{\partial V} + \frac{1}{2}\eta_1^2 V \frac{\partial^2 w}{\partial V^2} \\
&\quad + \frac{\partial w}{\partial t} + c(m' - V')\frac{\partial w}{\partial V'} + \frac{1}{2}\eta_2^2 V' \frac{\partial^2 w}{\partial V'^2} \\
\bar{\mathcal{L}}w &= \frac{\partial w}{\partial t} + \kappa(m - V)\frac{\partial w}{\partial V} + \frac{1}{2}\eta_1^2 V \frac{\partial^2 w}{\partial V^2}
\end{aligned} \tag{3.2.2}$$

Mis-pricing formula for it is then

$$\begin{aligned}
\delta &= (\mathcal{L} - \bar{\mathcal{L}})\bar{w} = \kappa(V' - m)\frac{\partial w}{\partial V} \\
&= \kappa(V' - m)\Gamma_{\bar{w}}
\end{aligned}$$

where $\Gamma_{\bar{w}}$ is given in (3.1.21).

Appendix A

Appendix A Title

Appendices at the end of a dissertation are optional, and depend on the content of the dissertation. There can be one or more appendices, however they should retain the page numbering requirements for dissertations. Any concerns about the formatting of an appendix should be brought to Doris Oliver, who can direct you how to format your appendix if you have questions.

Theoretical Dissertation Timeline		
Taskt	Time to Finish	Notes
Problem statement	10 hours	Initially very upbeat.
Research	3 days	Literature search to very previous studies.
Reformulation	4 hours	Presented and accepted by advisor
Research	20 days	Literature search to very previous studies.
Experiments	14 days	Do some experiments and get results.
Format	1 day	Understand format guidelines for paper.
Write	years	Write the paper.
Revise	not too long	Proof read, etc.
Format	1-3 days	Verify correct report format is used.
See Library	1 hour	Meet with Doris to verify formatting.
Defend	1 day	Defend your research.
Revise	0 hours	It was perfect the first time.
Submit	1 day	Submit final dissertation to the library.

Bibliography

- Auster, J., L. Mathys, and F. Maeder (2021, May). JDOI Variance Reduction Method and the Pricing of American-Style Options. *arXiv:2104.01365 [q-fin]*. arXiv: 2104.01365.
- Chan, K. C., G. A. Karolyi, F. A. Longstaff, and A. B. Sanders (1992, July). An Empirical Comparison of Alternative Models of the Short-Term Interest Rate. *The Journal of Finance* 47(3), 1209–1227.
- Cohen, J. D. (1988, May). Noncentral Chi-Square: Some Observations on Recurrence. *The American Statistician* 42(2), 120.
- Gatheral, J. (2008). Consistent Modeling of SPX and VIX options. pp. 75.
- Grünbichler, A. and F. A. Longstaff (1996, July). Valuing futures and options on volatility. *Journal of Banking & Finance* 20(6), 985–1001.
- Heath, D. and E. Platen. A Monte Carlo Method using PDE Expansions for a Diversified Equity Index Model. pp. 31.
- Hossain, A. S., M. T. Noushin, and K. Hasan (2019, July). A Comparison of the Black-Scholes Option Pricing Model and Its Alternatives. *Dhaka University Journal of Science* 67(2), 105–110.
- Kristensen, D. and A. Mele (2011, November). Adding and subtracting Black-Scholes: A new approach to approximating derivative prices in continuous-time models. *Journal of Financial Economics* 102(2), 390–415.