

[1]Highlights

Neural trajectories in the hippocampus exhibited greater variability during a working memory (WM) task compared to

- G. Buzski, Hippocampal sharp wave-ripple: A cognitive biomarker for episodic memory and planning, *Hippocampus* 25 (1995) 3–18. URL <https://onlinelibrary.wiley.com/doi/abs/10.1002/hipo.22488>
- S. P. Jadhav, C. Kemere, P. W. German, L. M. Frank, Awake Hippocampal Sharp-Wave Ripples Support Spatial Memory, *Neuron* 82 (2014) 236–249. URL <https://www.science.org/doi/abs/10.1126/science.1217230>
- M. A. Wilson, B. L. McNaughton, Reactivation of hippocampal ensemble memories during sleep, *Science (New York, N.Y.)* 261 (1992) 955–958.
- Z. Nadasdy, H. Hirase, A. Czurk, J. Csicsvari, G. Buzski, Replay and Time Compression of Recurring Spike Sequences in the Hippocampus, *Neuron* 26 (1999) 991–1002. URL <https://www.jneurosci.org/content/19/21/9497>
- A. K. Lee, M. A. Wilson, Memory of sequential experience in the hippocampus during slow wave sleep, *Neuron* 36 (6) (2002) 1005–1019.
- K. Diba, G. Buzski, Forward and reverse hippocampal place-cell sequences during ripples, *Nature Neuroscience* 10 (10) (2007) 1202–1208. URL <https://www.nature.com/articles/nn1961>
- T. J. Davidson, F. Kloosterman, M. A. Wilson, Hippocampal replay of extended experience, *Neuron* 63 (4) (2009) 497–507.
- G. Girardeau, K. Benchenane, S. I. Wiener, G. Buzski, M. B. Zugaro, Selective suppression of hippocampal ripples impairs spatial memory, *Nature Neuroscience* 13 (2010) 1542–1548. URL <http://www.nature.com/articles/nn.2384>
- V. Ego-Stengel, M. A. Wilson, Disruption of ripple-associated hippocampal activity during rest impairs spatial learning in the rat, *Neuroscience* 129 (2005) 572–581.
- A. Fernandez-Ruiz, A. Oliva, E. Fermino de Oliveira, F. Rocha-Almeida, D. Tingley, G. Buzski, Long-duration hippocampal ripples support spatial memory, *Science* 335 (6078) (2011) 1017–1020. URL <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6693581/>
- J. Kim, A. Joshi, L. Frank, K. Ganguly, Corticalhippocampal coupling during manifold exploration in motor cortex, *Nature Neuroscience* 15 (2012) 1039–1047. URL <https://www.nature.com/articles/s41586-022-05533-z>
- C.-T. Wu, D. Haggerty, C. Kemere, D. Ji, Hippocampal awake replay in fear memory retrieval, *Nature Neuroscience* 20 (2017) 1537–1545.
- Y. Norman, E. M. Yeagle, S. Khuvis, M. Harel, A. D. Mehta, R. Malach, Hippocampal sharp-wave ripples linked to visual memory, *Science* 335 (6078) (2011) 1021–1024. URL <https://www.sciencemag.org/lookup/doi/10.1126/science.aax1030>
- Y. Norman, O. Raccach, S. Liu, J. Parvizi, R. Malach, Hippocampal ripples and their coordinated dialogue with the default network, *Neuron* 73 (2012) 1025–1037. URL [https://www.cell.com/neuron/abstract/S0896-6273\(21\)00461-X](https://www.cell.com/neuron/abstract/S0896-6273(21)00461-X)
- C. J. Behrens, L. P. van den Boom, L. de Hoz, A. Friedman, U. Heinemann, Induction of sharp waveripple complexes in the rat hippocampus, *Neuron* 50 (2006) 663–674. URL <https://www.nature.com/articles/nn1571>
- H. Norimoto, K. Makino, M. Gao, Y. Shikano, K. Okamoto, T. Ishikawa, T. Sasaki, H. Hioki, S. Fujisawa, Y. Ikegaya, H. Okabe, J. O’Keefe, J. Dostrovsky, The hippocampus as a spatial map: Preliminary evidence from unit activity in the freely-moving rat, *Neuroscience* 30 (1980) 593–601.
- J. O’Keefe, Place units in the hippocampus of the freely moving rat, *Experimental Neurology* 51 (1) (1976) 78–109. doi:10.1016/0014-7039(76)90051-0. URL <https://www.sciencedirect.com/science/article/pii/0014488676900558>
- A. D. Ekstrom, M. J. Kahana, J. B. Caplan, T. A. Fields, E. A. Isham, E. L. Newman, I. Fried, Cellular networks underlying human spatial memory, *Nature* 427 (2004) 429–434. URL <https://www.nature.com/articles/nature01964>
- K. B. Kjelstrup, T. Solstad, V. H. Brun, T. Hafting, S. Leutgeb, M. P. Witter, E. I. Moser, M.-B. Moser, Finite Scale of Spatial Representation in the Adult Rat Hippocampus, *Science* 325 (5945) (2008) 91–98. URL <https://www.science.org/doi/abs/10.1126/science.1157086>
- C. D. Harvey, F. Collman, D. A. Dombeck, D. W. Tank, Intracellular dynamics of hippocampal place cells during virtual navigation, *Nature* 457 (2009) 394–399. URL <https://www.nature.com/articles/nature08499>
- H. Zhang, P. D. Rich, A. K. Lee, T. O. Sharpee, Hippocampal spatial representations exhibit a hyperbolic geometry that is independent of the environment, *Neuron* 73 (2012) 1038–1050. URL <https://www.nature.com/articles/s41593-022-01212-4>
- P. A. Naber, F. H. Lopes da Silva, M. P. Witter, Reciprocal connections between the entorhinal cortex and hippocampus in the rat, *Hippocampus* 12 (2002) 1002–1015. URL <https://onlinelibrary.wiley.com/doi/abs/10.1002/hipo.1028>
- N. M. van Strien, N. L. M. Cappaert, M. P. Witter, The anatomy of memory: an interactive overview of the parahippocampal-gyrus complex, *Neuroscience* 127 (2005) 127–148. URL <https://www.nature.com/articles/nrn2614>
- B. A. Strange, M. P. Witter, E. S. Lein, E. I. Moser, Functional organization of the hippocampal longitudinal axis, *Nature Neuroscience* 1 (1998) 202–205. URL <https://www.nature.com/articles/nrn3785>
- R. J. Gardner, E. Hermansen, M. Pachitariu, Y. Burak, N. A. Baas, B. A. Dunn, M.-B. Moser, E. I. Moser, Toroidal topology of the hippocampal theta rhythm, *Nature Neuroscience* 15 (2012) 1039–1047. URL <https://www.nature.com/articles/s41586-021-04268-7>
- Y. Watanabe, M. Okada, Y. Ikegaya, Towards threshold invariance in defining hippocampal ripples, *Journal of Neural Engineering* 14 (2017) 060101. URL <https://dx.doi.org/10.1088/1741-2552/ac3266>
- E. Boran, T. Fedele, A. Steiner, P. Hilfiker, L. Stieglitz, T. Grunwald, J. Sarnthein, Dataset of human medial temporal lobe sharp wave-ripples, *Neuroscience* 335 (2017) 1025–1037. URL <https://www.nature.com/articles/s41597-020-0364-3>
- B. M. Yu, J. P. Cunningham, G. Santhanam, S. I. Ryu, K. V. Shenoy, M. Sahani, Gaussian-Process Factor Analysis for Robust Latent Variable Modeling of Population Activity, *Neuron* 62 (2009) 920–934. URL <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2712272/>
- J. Niediek, J. Bostrom, C. E. Elger, F. Mormann, Reliable Analysis of Single-Unit Recordings from the Human Brain using a Bayesian Approach, *PLOS ONE* 13 (2018) e0166598. URL <https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0166598>
- A. A. Liu, S. Henin, S. Abbaspoor, A. Bragin, E. A. Buffalo, J. S. Farrell, D. J. Foster, L. M. Frank, T. Gedankien, J. C. Magee, Hippocampal sharp-wave ripples are generated by a single population of cells, *Nature Neuroscience* 15 (2012) 1039–1047. URL <https://www.nature.com/articles/s41467-022-33536-x>
- K. Kay, M. Sosa, J. E. Chung, M. P. Karlsson, M. C. Larkin, L. M. Frank, A hippocampal network for spatial coding during awake behavior, *Neuron* 73 (2012) 1038–1050.
- L. McInnes, J. Healy, N. Saul, L. Groberger, UMAP: Uniform Manifold Approximation and Projection, *Journal of Open Source Software* 1 (2018) 00861. URL <https://joss.theoj.org/papers/10.21105/joss.00861>
- P. J. Rousseeuw, Silhouettes: A graphical aid to the interpretation and validation of cluster analysis, *Journal of Computational Graphics* 7 (1987) 223–231. URL <https://www.sciencedirect.com/science/article/pii/0377042787901257>
- P. Virtanen, R. Gommers, T. E. Oliphant, M. Haberland, T. Reddy, D. Cournapeau, E. Burovski, P. Peterson, W. Weckstein, J. Virtanen, SciPy: Open Source Scientific Data Analysis, *Proceedings of the 9th Python Conference* (2007) 261–272. URL <https://ui.adsabs.harvard.edu/abs/2020NatMe...17..261V>
- G. Buzski, Two-stage model of memory trace formation: a role for “noisy” brain states, *Neuroscience* 31 (3) (1989) 551–570.
- M. L. V. Quyen, A. Bragin, R. St

Subject ID# of sessions									
AHL	AHR	PHL	PHR	CLE	CRA	LAR			
1	4	n.a.				n.a.			
lightgray 2	7								
3	3								n.a.
*Tables [1]Table	tables [2]ID 01id ₀ 1	[htbp]	lightgray 4	2					AHL
5	3	n.a.	n.a.	n.a.	n.a.	n.a.			
lightgray 6	6								AHL
7	4								
lightgray 8	5								
9	2								

This figure denotes the placements of electrodes and seizure onset zones. Regions marked with were included in the

Subject	AHL	AHR	PHL	PHR					
1	0.60	0.14	n.a.	n.a.	0.1	0			
lightgray 20.21	0.16	0.17	0.21	0.18	0.22	0.20	0.15		
3	0.40	0.42	0.83	0.12	n.a.	n.a.			
[2]ID 02id ₀ 2	[htbp]	lightgray 40.10	0.00	0.10	0.00	0.90	0.00	0.10	0.14
5	n.a.	n.a.	n.a.	n.a.					
lightgray 60.63	0.06	n.a.	n.a.	0.27	0.06				
7	0.10	0.00	0.35	0.37	0.47	0.10	0.00		
lightgray 80.13	0.10	n.a.	0.28	0.49	n.a.				
9	n.a.	0.85	0.07	0.15	0.07	n.a.			

The silhouette scores (mean \pm SD across sessions per subject) for UMAP clustering of SWR^+ candidates and SWR^- candidates

Subject ID# of sessions# of trialsROI# of SWRsSWR incidence [Hz]							
[2]ID 03id ₀ 3[htbp]	#1	2	100	AHL	274	0.34	width = Summ
	lightgray #3	2	97	AHR	325	0.42	
	#4	2	99	PHL	202	0.26	
	lightgray #6	2	100	AHL	297	0.37	
	#9	2	97	AHR	72	0.09	
lightgray	Total = 10	Total = 493	Total = 1170	0.30	0.13 (mean	SD)	

The table provides statistics of presumptive CA1 regions and SWR events. Only the initial two sessions (sessions 1 and 2) are shown.

*Figures [1]Figuresfigures

[ht] [2]ID 01figure_id01[width = 1]./src/figures/.png/Figure_ID01.png **Local Field Potentials, Multiunit Activity**
A. Representative wideband LFP signals for intracranial EEG recording from the left hippocampal head are presented.

[ht] [2]ID 02figure_id02[width = 0.5]./src/figures/.png/Figure_ID02.png **State-Dependent Neural Trajectory** of
A. Neural trajectories (NTs) depicted as a point cloud within the first three-dimensional factors derived from GPF

[ht] [2]ID 03figure_id03[width = 1]./src/figures/.png/Figure_ID03.png **Positive Correlation between Memory**
A. The relationship between set size (number of letters to be encoded) and accuracy in the working memory task (

[ht] [2]ID 04figure_id04[width = 1]./src/figures/.png/figure_ID04.png**Detection of SWRs in Putative CA1 R**
A. Two-dimensional UMAP [0] projection displays multi-unit spikes during SWR⁺ candidates (*purple*) and SWR⁻

[ht] [2]ID 05figure_id05[width = 1]./src/figures/.png/Figure_ID05.png**Transient Change in Neural Trajectory**
A. The distance from origin (*O*) of the peri-sharp-wave-ripple neural trajectory (mean $\pm 95\%$ confidence interval).

[ht] [2]ID 06figure_id06[width = 1]./src/figures/.png/Figure_ID06.png **Visualization of Neural Trajectory Du**
The panels depict hippocampal neural trajectories (NTs) during SWR projected onto two-dimensional spaces. **A.** S

[ht] [2]ID 07figure_id07[width = 0.5]./src/figures/.png/Figure_ID07.png **Direction of Neural Trajectory During**
A–B The kernel density estimation distributions of $\overrightarrow{\text{eSWR}} \cdot \overrightarrow{\text{rSWR}}$ (*pink circles*), $\overrightarrow{\text{eSWR}} \cdot \overrightarrow{\text{gEGR}}$ (*blue triangles*)