

Finite subgroups of derived automorphisms of generic K3 surfaces

Yu-Wei Fan (Tsinghua)
Joint work with Kuan-Wen Lai (Bonn)

July 2023
1st International Congress of Basic Science

Slogan of main results

Let X be a Calabi–Yau manifold. It is expected that

$$D := D^b\mathrm{Coh}(X) \cong D^\pi\mathrm{Fuk}(X^\vee).$$

This triangulated category has autoequivalences arising from both complex geometry of X and symplectic geometry of X^\vee , for instance:

$$\mathrm{Aut}(X), \quad L \otimes -; \quad \mathrm{Symp}(X^\vee).$$

Slogan: When X is a K3 surface of Picard number one,

- there is no finite order autoequivalence arising from either $\mathrm{Aut}(X)$, $L \otimes -$, or $\mathrm{Symp}(X^\vee)$;
- there are interesting finite order autoequivalences given by **mixings** of complex and symplectic geometric autoequivalences;
- we give full **classification** and **counting** of finite subgroups of $\mathrm{Aut}(D)$ and $\mathrm{Aut}(D)/[2]$ up to conjugations.

Slogan of main results

Let X be a Calabi–Yau manifold. It is expected that

$$D := D^b\mathrm{Coh}(X) \cong D^\pi\mathrm{Fuk}(X^\vee).$$

This triangulated category has autoequivalences arising from both complex geometry of X and symplectic geometry of X^\vee , for instance:

$$\mathrm{Aut}(X), \quad L \otimes -; \quad \mathrm{Symp}(X^\vee).$$

Slogan: When X is a K3 surface of Picard number one,

- there is no finite order autoequivalence arising from either $\mathrm{Aut}(X)$, $L \otimes -$, or $\mathrm{Symp}(X^\vee)$;
- there are interesting finite order autoequivalences given by **mixings** of complex and symplectic geometric autoequivalences;
- we give full **classification** and **counting** of finite subgroups of $\mathrm{Aut}(D)$ and $\mathrm{Aut}(D)/[2]$ up to conjugations.

Slogan of main results

Let X be a Calabi–Yau manifold. It is expected that

$$D := D^b\mathrm{Coh}(X) \cong D^\pi\mathrm{Fuk}(X^\vee).$$

This triangulated category has autoequivalences arising from both complex geometry of X and symplectic geometry of X^\vee , for instance:

$$\mathrm{Aut}(X), \quad L \otimes -; \quad \mathrm{Symp}(X^\vee).$$

Slogan: When X is a K3 surface of Picard number one,

- there is no finite order autoequivalence arising from either $\mathrm{Aut}(X)$, $L \otimes -$, or $\mathrm{Symp}(X^\vee)$;
- there are interesting finite order autoequivalences given by **mixings** of complex and symplectic geometric autoequivalences;
- we give full **classification** and **counting** of finite subgroups of $\mathrm{Aut}(D)$ and $\mathrm{Aut}(D)/[2]$ up to conjugations.

Slogan of main results

Let X be a Calabi–Yau manifold. It is expected that

$$D := D^b\mathrm{Coh}(X) \cong D^\pi\mathrm{Fuk}(X^\vee).$$

This triangulated category has autoequivalences arising from both complex geometry of X and symplectic geometry of X^\vee , for instance:

$$\mathrm{Aut}(X), \quad L \otimes -; \quad \mathrm{Symp}(X^\vee).$$

Slogan: When X is a K3 surface of Picard number one,

- there is no finite order autoequivalence arising from either $\mathrm{Aut}(X)$, $L \otimes -$, or $\mathrm{Symp}(X^\vee)$;
- there are interesting finite order autoequivalences given by **mixings** of complex and symplectic geometric autoequivalences;
- we give full **classification** and **counting** of finite subgroups of $\mathrm{Aut}(D)$ and $\mathrm{Aut}(D)/[2]$ up to conjugations.

Slogan of main results

Let X be a Calabi–Yau manifold. It is expected that

$$D := D^b\mathrm{Coh}(X) \cong D^\pi\mathrm{Fuk}(X^\vee).$$

This triangulated category has autoequivalences arising from both complex geometry of X and symplectic geometry of X^\vee , for instance:

$$\mathrm{Aut}(X), \quad L \otimes -; \quad \mathrm{Symp}(X^\vee).$$

Slogan: When X is a K3 surface of Picard number one,

- there is no finite order autoequivalence arising from either $\mathrm{Aut}(X)$, $L \otimes -$, or $\mathrm{Symp}(X^\vee)$;
- there are interesting finite order autoequivalences given by **mixings** of complex and symplectic geometric autoequivalences;
- we give full **classification** and **counting** of finite subgroups of $\mathrm{Aut}(D)$ and $\mathrm{Aut}(D)/[2]$ up to conjugations.

Slogan of main results

Let X be a Calabi–Yau manifold. It is expected that

$$D := D^b\mathrm{Coh}(X) \cong D^\pi\mathrm{Fuk}(X^\vee).$$

This triangulated category has autoequivalences arising from both complex geometry of X and symplectic geometry of X^\vee , for instance:

$$\mathrm{Aut}(X), \quad L \otimes -; \quad \mathrm{Symp}(X^\vee).$$


Slogan: When X is a K3 surface of Picard number one,

- there is no finite order autoequivalence arising from either $\mathrm{Aut}(X)$, $L \otimes -$, or $\mathrm{Symp}(X^\vee)$;
- there are interesting finite order autoequivalences given by **mixings** of complex and symplectic geometric autoequivalences;
- we give full **classification** and **counting** of finite subgroups of $\mathrm{Aut}(D)$ and $\mathrm{Aut}(D)/[2]$ up to conjugations.

Finite order autoequivalences and Gepner points

Consider an one-parameter family of Calabi–Yau manifolds:

$$\{x_0^{n+1} + x_1^{n+1} + \cdots + x_n^{n+1} + t \cdot x_0 x_1 \cdots x_n = 0\} \subseteq \mathbb{CP}^n.$$



The monodromies correspond to autoequivalences of $D^b\mathrm{Coh}(X)$:


- large complex structure limit point: $\mathcal{O}(1) \otimes -$;
- conifold point: $T_{\mathcal{O}_X}(-)$;
- Gepner point: $\Phi := T_{\mathcal{O}_X} \circ (- \otimes \mathcal{O}(1))$, where $\Phi^{n+1} = [2]$.

In particular, when X is a K3 surface, we have $\Phi^4 = [2]$, therefore $\Phi^2[-1]$ is of order two in $\mathrm{Aut}(D^b(X))$.

Finite order autoequivalences and Gepner points

Consider an one-parameter family of Calabi–Yau manifolds:

$$\{x_0^{n+1} + x_1^{n+1} + \cdots + x_n^{n+1} + t \cdot x_0 x_1 \cdots x_n = 0\} \subseteq \mathbb{CP}^n.$$



The monodromies correspond to autoequivalences of $D^b\mathrm{Coh}(X)$:


- large complex structure limit point: $\mathcal{O}(1) \otimes -$;
- conifold point: $T_{\mathcal{O}_X}(-)$;
- Gepner point: $\Phi := T_{\mathcal{O}_X} \circ (- \otimes \mathcal{O}(1))$, where $\Phi^{n+1} = [2]$.

In particular, when X is a K3 surface, we have $\Phi^4 = [2]$, therefore $\Phi^2[-1]$ is of order two in $\mathrm{Aut}(D^b(X))$.

Finite order autoequivalences and Gepner points

Consider an one-parameter family of Calabi–Yau manifolds:

$$\{x_0^{n+1} + x_1^{n+1} + \cdots + x_n^{n+1} + t \cdot x_0 x_1 \cdots x_n = 0\} \subseteq \mathbb{CP}^n.$$



The monodromies correspond to autoequivalences of $D^b\mathrm{Coh}(X)$:


- large complex structure limit point: $\mathcal{O}(1) \otimes -$;
- conifold point: $T_{\mathcal{O}_X}(-)$;
- Gepner point: $\Phi := T_{\mathcal{O}_X} \circ (- \otimes \mathcal{O}(1))$, where $\Phi^{n+1} = [2]$.

In particular, when X is a K3 surface, we have $\Phi^4 = [2]$, therefore $\Phi^2[-1]$ is of order two in $\mathrm{Aut}(D^b(X))$.

Finite order autoequivalences and Gepner points

Consider an one-parameter family of Calabi–Yau manifolds:

$$\{x_0^{n+1} + x_1^{n+1} + \cdots + x_n^{n+1} + t \cdot x_0 x_1 \cdots x_n = 0\} \subseteq \mathbb{CP}^n.$$



The monodromies correspond to autoequivalences of $D^b\mathrm{Coh}(X)$:


- large complex structure limit point: $\mathcal{O}(1) \otimes -$;
- conifold point: $T_{\mathcal{O}_X}(-)$;
- Gepner point: $\Phi := T_{\mathcal{O}_X} \circ (- \otimes \mathcal{O}(1))$, where $\Phi^{n+1} = [2]$.

In particular, when X is a K3 surface, we have $\Phi^4 = [2]$, therefore $\Phi^2[-1]$ is of order two in $\mathrm{Aut}(D^b(X))$.

Finite order autoequivalences and Gepner points

Consider an one-parameter family of Calabi–Yau manifolds:

$$\{x_0^{n+1} + x_1^{n+1} + \cdots + x_n^{n+1} + t \cdot x_0 x_1 \cdots x_n = 0\} \subseteq \mathbb{CP}^n.$$



The monodromies correspond to autoequivalences of $D^b\mathrm{Coh}(X)$:


- large complex structure limit point: $\mathcal{O}(1) \otimes -$;
- conifold point: $T_{\mathcal{O}_X}(-)$;
- Gepner point: $\Phi := T_{\mathcal{O}_X} \circ (- \otimes \mathcal{O}(1))$, where $\Phi^{n+1} = [2]$.

In particular, when X is a K3 surface, we have $\Phi^4 = [2]$, therefore $\Phi^2[-1]$ is of order two in $\mathrm{Aut}(D^b(X))$.

Finite order autoequivalences and Gepner points

Consider an one-parameter family of Calabi–Yau manifolds:

$$\{x_0^{n+1} + x_1^{n+1} + \cdots + x_n^{n+1} + t \cdot x_0 x_1 \cdots x_n = 0\} \subseteq \mathbb{CP}^n.$$



The monodromies correspond to autoequivalences of $D^b\mathrm{Coh}(X)$:

- large complex structure limit point: $\mathcal{O}(1) \otimes -$;
- conifold point: $T_{\mathcal{O}_X}(-)$;
- Gepner point: $\Phi := T_{\mathcal{O}_X} \circ (- \otimes \mathcal{O}(1))$, where $\Phi^{n+1} = [2]$.

In particular, when X is a K3 surface, we have $\Phi^4 = [2]$, therefore $\Phi^2[-1]$ is of order two in $\mathrm{Aut}(D^b(X))$.

Automorphisms of K3 surfaces

As a warm-up, let us sketch the proof of the following result.

Theorem (Nikulin)

Let X be a complex projective K3 surface with $NS(X) \cong \langle H \rangle$. Then

$$\mathrm{Aut}(X) = \begin{cases} \{id\} & \text{if } H^2 > 2 \\ \mathbb{Z}/2\mathbb{Z} & \text{if } H^2 = 2 \end{cases}$$

- There is an injective map $\mathrm{Aut}(X) \hookrightarrow O(H^2(X, \mathbb{Z}))$.
- $f \in \mathrm{Aut}(X)$ acts trivially on $NS(X)$: pullback of H is still ample.
- f acts as $\pm id$ on $T(X) := NS(X)^\perp$: true for any odd Picard number.
- Its induced actions on the discriminant groups $T(X)^*/T(X)$ and $NS(X)^*/NS(X) \cong \mathbb{Z}/(H^2)\mathbb{Z}$ coincide $\implies f = id$ unless $H^2 = 2$.
- When $H^2 = 2$, X is a double cover over \mathbb{P}^2 and the covering involution gives the nontrivial automorphism.

Automorphisms of K3 surfaces

As a warm-up, let us sketch the proof of the following result.

Theorem (Nikulin)

Let X be a complex projective K3 surface with $NS(X) \cong \langle H \rangle$. Then

$$\mathrm{Aut}(X) = \begin{cases} \{id\} & \text{if } H^2 > 2 \\ \mathbb{Z}/2\mathbb{Z} & \text{if } H^2 = 2 \end{cases}$$

- There is an injective map $\mathrm{Aut}(X) \hookrightarrow O(H^2(X, \mathbb{Z}))$.
- $f \in \mathrm{Aut}(X)$ acts trivially on $NS(X)$: pullback of H is still ample.
- f acts as $\pm id$ on $T(X) := NS(X)^\perp$: true for any odd Picard number.
- Its induced actions on the discriminant groups $T(X)^*/T(X)$ and $NS(X)^*/NS(X) \cong \mathbb{Z}/(H^2)\mathbb{Z}$ coincide $\implies f = id$ unless $H^2 = 2$.
- When $H^2 = 2$, X is a double cover over \mathbb{P}^2 and the covering involution gives the nontrivial automorphism.

Automorphisms of K3 surfaces

As a warm-up, let us sketch the proof of the following result.

Theorem (Nikulin)

Let X be a complex projective K3 surface with $NS(X) \cong \langle H \rangle$. Then

$$\mathrm{Aut}(X) = \begin{cases} \{id\} & \text{if } H^2 > 2 \\ \mathbb{Z}/2\mathbb{Z} & \text{if } H^2 = 2 \end{cases}$$

- There is an injective map $\mathrm{Aut}(X) \hookrightarrow O(H^2(X, \mathbb{Z}))$.
- $f \in \mathrm{Aut}(X)$ acts trivially on $NS(X)$: pullback of H is still ample.
- f acts as $\pm id$ on $T(X) := NS(X)^\perp$: true for any odd Picard number.
- Its induced actions on the discriminant groups $T(X)^*/T(X)$ and $NS(X)^*/NS(X) \cong \mathbb{Z}/(H^2)\mathbb{Z}$ coincide $\implies f = id$ unless $H^2 = 2$.
- When $H^2 = 2$, X is a double cover over \mathbb{P}^2 and the covering involution gives the nontrivial automorphism.

Automorphisms of K3 surfaces

As a warm-up, let us sketch the proof of the following result.

Theorem (Nikulin)

Let X be a complex projective K3 surface with $NS(X) \cong \langle H \rangle$. Then

$$\mathrm{Aut}(X) = \begin{cases} \{id\} & \text{if } H^2 > 2 \\ \mathbb{Z}/2\mathbb{Z} & \text{if } H^2 = 2 \end{cases}$$

- There is an injective map $\mathrm{Aut}(X) \hookrightarrow O(H^2(X, \mathbb{Z}))$.
- $f \in \mathrm{Aut}(X)$ acts trivially on $NS(X)$: pullback of H is still ample.
- f acts as $\pm id$ on $T(X) := NS(X)^\perp$: true for any odd Picard number.
- Its induced actions on the discriminant groups $T(X)^*/T(X)$ and $NS(X)^*/NS(X) \cong \mathbb{Z}/(H^2)\mathbb{Z}$ coincide $\implies f = id$ unless $H^2 = 2$.
- When $H^2 = 2$, X is a double cover over \mathbb{P}^2 and the covering involution gives the nontrivial automorphism.

Automorphisms of K3 surfaces

As a warm-up, let us sketch the proof of the following result.

Theorem (Nikulin)

Let X be a complex projective K3 surface with $NS(X) \cong \langle H \rangle$. Then

$$\mathrm{Aut}(X) = \begin{cases} \{id\} & \text{if } H^2 > 2 \\ \mathbb{Z}/2\mathbb{Z} & \text{if } H^2 = 2 \end{cases}$$

- There is an injective map $\mathrm{Aut}(X) \hookrightarrow O(H^2(X, \mathbb{Z}))$.
- $f \in \mathrm{Aut}(X)$ acts trivially on $NS(X)$: pullback of H is still ample.
- f acts as $\pm id$ on $T(X) := NS(X)^\perp$: true for any odd Picard number.
- Its induced actions on the discriminant groups $T(X)^*/T(X)$ and $NS(X)^*/NS(X) \cong \mathbb{Z}/(H^2)\mathbb{Z}$ coincide $\implies f = id$ unless $H^2 = 2$.
- When $H^2 = 2$, X is a double cover over \mathbb{P}^2 and the covering involution gives the nontrivial automorphism.

Automorphisms of K3 surfaces

As a warm-up, let us sketch the proof of the following result.

Theorem (Nikulin)

Let X be a complex projective K3 surface with $NS(X) \cong \langle H \rangle$. Then

$$\mathrm{Aut}(X) = \begin{cases} \{id\} & \text{if } H^2 > 2 \\ \mathbb{Z}/2\mathbb{Z} & \text{if } H^2 = 2 \end{cases}$$

- There is an injective map $\mathrm{Aut}(X) \hookrightarrow O(H^2(X, \mathbb{Z}))$.
- $f \in \mathrm{Aut}(X)$ acts trivially on $NS(X)$: pullback of H is still ample.
- f acts as $\pm id$ on $T(X) := NS(X)^\perp$: true for any odd Picard number.
- Its induced actions on the discriminant groups $T(X)^*/T(X)$ and $NS(X)^*/NS(X) \cong \mathbb{Z}/(H^2)\mathbb{Z}$ coincide $\implies f = id$ unless $H^2 = 2$.
- When $H^2 = 2$, X is a double cover over \mathbb{P}^2 and the covering involution gives the nontrivial automorphism.

Automorphisms of K3 surfaces

As a warm-up, let us sketch the proof of the following result.

Theorem (Nikulin)

Let X be a complex projective K3 surface with $NS(X) \cong \langle H \rangle$. Then

$$\mathrm{Aut}(X) = \begin{cases} \{id\} & \text{if } H^2 > 2 \\ \mathbb{Z}/2\mathbb{Z} & \text{if } H^2 = 2 \end{cases}$$

- There is an injective map $\mathrm{Aut}(X) \hookrightarrow O(H^2(X, \mathbb{Z}))$.
- $f \in \mathrm{Aut}(X)$ acts trivially on $NS(X)$: pullback of H is still ample.
- f acts as $\pm id$ on $T(X) := NS(X)^\perp$: true for any odd Picard number.
- Its induced actions on the discriminant groups $T(X)^*/T(X)$ and $NS(X)^*/NS(X) \cong \mathbb{Z}/(H^2)\mathbb{Z}$ coincide $\implies f = id$ unless $H^2 = 2$.
- When $H^2 = 2$, X is a double cover over \mathbb{P}^2 and the covering involution gives the nontrivial automorphism.

Some difficulties of generalizing to $\text{Aut}(D^b(X))$

- If $\Phi \in \text{Aut}(D^b(X))$ is of finite order, then its induced actions on

$$T(X) \quad \text{and} \quad N(X) = H^0(X) \oplus \text{NS}(X) \oplus H^4(X) \cong \mathbb{Z}^3$$

are of finite order.

- Φ still acts as $\pm \text{id}$ on $T(X)$; but its action on $N(X)$ can be more complicated.

Strategy: We show that any finite order Φ fixes a **Bridgeland stability condition** on $D^b(X)$. In particular, it fixes a 2-plane in $N(X)_{\mathbb{R}}$ pointwisely, therefore the action of Φ on $N(X)$ is either id or a reflection.

- In order to fully classify the finite subgroups of $\text{Aut}(D^b(X))$, one needs to determine the conditions on finite subgroups of $T(X)$ and $N(X)$ of which there exists finite order lifts in $\text{Aut}(D^b(X))$.

Some difficulties of generalizing to $\text{Aut}(D^b(X))$

- If $\Phi \in \text{Aut}(D^b(X))$ is of finite order, then its induced actions on

$$T(X) \quad \text{and} \quad N(X) = H^0(X) \oplus \text{NS}(X) \oplus H^4(X) \cong \mathbb{Z}^3$$

are of finite order.

- Φ still acts as $\pm \text{id}$ on $T(X)$; but its action on $N(X)$ can be more complicated.

Strategy: We show that any finite order Φ fixes a **Bridgeland stability condition** on $D^b(X)$. In particular, it fixes a 2-plane in $N(X)_{\mathbb{R}}$ pointwisely, therefore the action of Φ on $N(X)$ is either id or a reflection.

- In order to fully classify the finite subgroups of $\text{Aut}(D^b(X))$, one needs to determine the conditions on finite subgroups of $T(X)$ and $N(X)$ of which there exists finite order lifts in $\text{Aut}(D^b(X))$.

Some difficulties of generalizing to $\text{Aut}(D^b(X))$

- If $\Phi \in \text{Aut}(D^b(X))$ is of finite order, then its induced actions on

$$T(X) \quad \text{and} \quad N(X) = H^0(X) \oplus \text{NS}(X) \oplus H^4(X) \cong \mathbb{Z}^3$$

are of finite order.

- Φ still acts as $\pm \text{id}$ on $T(X)$; but its action on $N(X)$ can be more complicated.

Strategy: We show that any finite order Φ fixes a **Bridgeland stability condition** on $D^b(X)$. In particular, it fixes a 2-plane in $N(X)_{\mathbb{R}}$ pointwisely, therefore the action of Φ on $N(X)$ is either id or a reflection.

- In order to fully classify the finite subgroups of $\text{Aut}(D^b(X))$, one needs to determine the conditions on finite subgroups of $T(X)$ and $N(X)$ of which there exists finite order lifts in $\text{Aut}(D^b(X))$.

Some difficulties of generalizing to $\text{Aut}(D^b(X))$

- If $\Phi \in \text{Aut}(D^b(X))$ is of finite order, then its induced actions on

$$T(X) \quad \text{and} \quad N(X) = H^0(X) \oplus \text{NS}(X) \oplus H^4(X) \cong \mathbb{Z}^3$$

are of finite order.

- Φ still acts as $\pm \text{id}$ on $T(X)$; but its action on $N(X)$ can be more complicated.

Strategy: We show that any finite order Φ fixes a **Bridgeland stability condition** on $D^b(X)$. In particular, it fixes a 2-plane in $N(X)_{\mathbb{R}}$ pointwisely, therefore the action of Φ on $N(X)$ is either id or a reflection.

- In order to fully classify the finite subgroups of $\text{Aut}(D^b(X))$, one needs to determine the conditions on finite subgroups of $T(X)$ and $N(X)$ of which there exists finite order lifts in $\text{Aut}(D^b(X))$.

Some difficulties of generalizing to $\text{Aut}(D^b(X))$

- If $\Phi \in \text{Aut}(D^b(X))$ is of finite order, then its induced actions on

$$T(X) \quad \text{and} \quad N(X) = H^0(X) \oplus \text{NS}(X) \oplus H^4(X) \cong \mathbb{Z}^3$$

are of finite order.

- Φ still acts as $\pm \text{id}$ on $T(X)$; but its action on $N(X)$ can be more complicated.

Strategy: We show that any finite order Φ fixes a **Bridgeland stability condition** on $D^b(X)$. In particular, it fixes a 2-plane in $N(X)_{\mathbb{R}}$ pointwisely, therefore the action of Φ on $N(X)$ is either id or a reflection.

- In order to fully classify the finite subgroups of $\text{Aut}(D^b(X))$, one needs to determine the conditions on finite subgroups of $T(X)$ and $N(X)$ of which there exists finite order lifts in $\text{Aut}(D^b(X))$.

Autoequivalences act on stability conditions

A natural way of studying a group is via considering its actions on various spaces. The group of autoequivalences $\text{Aut}(D)$ of a triangulated category admits a natural action on the space of its **Bridgeland stability conditions**.

This action $(\text{Aut}(D) \curvearrowright \text{Stab}(D))$ can be used to:

- define complexity (e.g. categorical entropy) of autoequivalences;
- provide classifications of autoequivalences (e.g. finite order, “reducible”, “pseudo-Anosov”, etc.);
- provide analogy with Teichmüller theory –

One of our key results, any finite order $\Phi \in \text{Aut}(D^b(X))$ fixes a stability condition on $D^b(X)$ (when X is a K3 surface of Picard number one), actually is motivated from the analogy with Teichmüller theory.

Autoequivalences act on stability conditions

A natural way of studying a group is via considering its actions on various spaces. The group of autoequivalences $\text{Aut}(D)$ of a triangulated category admits a natural action on the space of its **Bridgeland stability conditions**.

This action ($\text{Aut}(D) \curvearrowright \text{Stab}(D)$) can be used to:

- define complexity (e.g. categorical entropy) of autoequivalences;
- provide classifications of autoequivalences (e.g. finite order, “reducible”, “pseudo-Anosov”, etc.);
- provide analogy with Teichmüller theory –

One of our key results, any finite order $\Phi \in \text{Aut}(D^b(X))$ fixes a stability condition on $D^b(X)$ (when X is a K3 surface of Picard number one), actually is motivated from the analogy with Teichmüller theory.

Autoequivalences act on stability conditions

A natural way of studying a group is via considering its actions on various spaces. The group of autoequivalences $\text{Aut}(D)$ of a triangulated category admits a natural action on the space of its **Bridgeland stability conditions**.

This action ($\text{Aut}(D) \curvearrowright \text{Stab}(D)$) can be used to:

- define complexity (e.g. categorical entropy) of autoequivalences;
- provide classifications of autoequivalences (e.g. finite order, “reducible”, “pseudo-Anosov”, etc.);
- provide analogy with Teichmüller theory –

One of our key results, any finite order $\Phi \in \text{Aut}(D^b(X))$ fixes a stability condition on $D^b(X)$ (when X is a K3 surface of Picard number one), actually is motivated from the analogy with Teichmüller theory.

Autoequivalences act on stability conditions

A natural way of studying a group is via considering its actions on various spaces. The group of autoequivalences $\text{Aut}(D)$ of a triangulated category admits a natural action on the space of its **Bridgeland stability conditions**.

This action ($\text{Aut}(D) \curvearrowright \text{Stab}(D)$) can be used to:

- define complexity (e.g. categorical entropy) of autoequivalences;
- provide classifications of autoequivalences (e.g. finite order, “reducible”, “pseudo-Anosov”, etc.);
- provide analogy with Teichmüller theory –

One of our key results, any finite order $\Phi \in \text{Aut}(D^b(X))$ fixes a stability condition on $D^b(X)$ (when X is a K3 surface of Picard number one), actually is motivated from the analogy with Teichmüller theory.

Autoequivalences act on stability conditions

A natural way of studying a group is via considering its actions on various spaces. The group of autoequivalences $\text{Aut}(D)$ of a triangulated category admits a natural action on the space of its **Bridgeland stability conditions**.

This action ($\text{Aut}(D) \curvearrowright \text{Stab}(D)$) can be used to:

- define complexity (e.g. categorical entropy) of autoequivalences;
- provide classifications of autoequivalences (e.g. finite order, “reducible”, “pseudo-Anosov”, etc.);
- provide analogy with Teichmüller theory –

One of our key results, any finite order $\Phi \in \text{Aut}(D^b(X))$ fixes a stability condition on $D^b(X)$ (when X is a K3 surface of Picard number one), actually is motivated from the analogy with Teichmüller theory.

A dictionary of analogy

(after Gaiotto, Moore, Neitzke; Bridgeland, Smith; Dimitrov, Haiden, Katzarkov, Kontsevich, etc.)

Riemann surface Σ	Triangulated category \mathcal{D}
curve C	object E
$C_1 \cap C_2$	$\mathrm{Hom}(E_1, E_2)$
metric g	Bridgeland stability condition σ
geodesics	semistable objects
length $\ell_g(C)$	mass $m_\sigma(E)$
$\mathrm{MCG}(\Sigma)$	$\mathrm{Aut}(\mathcal{D})$
$\mathrm{Teich}(\Sigma)$	$\mathrm{Stab}(\mathcal{D})$
...	...

Many analogous statements in Teichmüller theory can be proved in the categorical setting for $\mathcal{D} = D^b\mathrm{Coh}(\text{elliptic curve})$. An interesting general question is whether some of these can be generalized to $\dim \geq 2$.

A dictionary of analogy

(after Gaiotto, Moore, Neitzke; Bridgeland, Smith; Dimitrov, Haiden, Katzarkov, Kontsevich, etc.)

Riemann surface Σ	Triangulated category \mathcal{D}
curve C	object E
$C_1 \cap C_2$	$\mathrm{Hom}(E_1, E_2)$
metric g	Bridgeland stability condition σ
geodesics	semistable objects
length $\ell_g(C)$	mass $m_\sigma(E)$
$\mathrm{MCG}(\Sigma)$	$\mathrm{Aut}(\mathcal{D})$
$\mathrm{Teich}(\Sigma)$	$\mathrm{Stab}(\mathcal{D})$
...	...

Many analogous statements in Teichmüller theory can be proved in the categorical setting for $\mathcal{D} = D^b\mathrm{Coh}(\text{elliptic curve})$. An interesting general question is whether some of these can be generalized to $\dim \geq 2$.

Nielsen realization problem

- Nielsen asked (1923): Let $G \subseteq \text{MCG}(\Sigma)$ be a finite subgroup. Does there always exist a lifting $G \subseteq \text{Diff}(\Sigma)$? (Recall that $\text{MCG}(\Sigma) = \text{Diff}(\Sigma)/\text{isotopy}$).
- Kerckhoff (1983): Yes! Moreover, there exists a metric g such that $G \subseteq \text{Isom}(\Sigma, g)$. Or equivalently, G fixes a point in $\text{Teich}(\Sigma)$. (There is a natural action of $\text{MCG}(\Sigma)$ on $\text{Teich}(\Sigma)$, e.g. $\text{MCG}(T^2) = \text{SL}(2, \mathbb{Z})$ acts on $\text{Teich}(T^2) = \mathbb{H}$.) (Rephrase: any finite subgroup of $\text{MCG}(\Sigma)$ can be realized as symmetries with respect to a metric on Σ .)
- Farb–Looijenga (2021) also proved similar statements for K3 surfaces (under certain conditions), where g is replaced by complex structure or Ricci-flat metric on the K3 surface.

Nielsen realization problem

- Nielsen asked (1923): Let $G \subseteq \text{MCG}(\Sigma)$ be a finite subgroup. Does there always exist a lifting $G \subseteq \text{Diff}(\Sigma)$? (Recall that $\text{MCG}(\Sigma) = \text{Diff}(\Sigma)/\text{isotopy}$).
- Kerckhoff (1983): Yes! Moreover, there exists a metric g such that $G \subseteq \text{Isom}(\Sigma, g)$. Or equivalently, G fixes a point in $\text{Teich}(\Sigma)$. (There is a natural action of $\text{MCG}(\Sigma)$ on $\text{Teich}(\Sigma)$, e.g. $\text{MCG}(T^2) = \text{SL}(2, \mathbb{Z})$ acts on $\text{Teich}(T^2) = \mathbb{H}$.) (Rephrase: any finite subgroup of $\text{MCG}(\Sigma)$ can be realized as symmetries with respect to a metric on Σ .)
- Farb–Looijenga (2021) also proved similar statements for K3 surfaces (under certain conditions), where g is replaced by complex structure or Ricci-flat metric on the K3 surface.

Nielsen realization problem

- Nielsen asked (1923): Let $G \subseteq \text{MCG}(\Sigma)$ be a finite subgroup. Does there always exist a lifting $G \subseteq \text{Diff}(\Sigma)$?
(Recall that $\text{MCG}(\Sigma) = \text{Diff}(\Sigma)/\text{isotopy}$).
- Kerckhoff (1983): Yes! Moreover, there exists a metric g such that $G \subseteq \text{Isom}(\Sigma, g)$. Or equivalently, G fixes a point in $\text{Teich}(\Sigma)$.
(There is a natural action of $\text{MCG}(\Sigma)$ on $\text{Teich}(\Sigma)$, e.g. $\text{MCG}(T^2) = \text{SL}(2, \mathbb{Z})$ acts on $\text{Teich}(T^2) = \mathbb{H}$.)
(Rephrase: any finite subgroup of $\text{MCG}(\Sigma)$ can be realized as symmetries with respect to a metric on Σ .)
- Farb–Looijenga (2021) also proved similar statements for K3 surfaces (under certain conditions), where g is replaced by complex structure or Ricci-flat metric on the K3 surface.

Nielsen realization problem

- Nielsen asked (1923): Let $G \subseteq \text{MCG}(\Sigma)$ be a finite subgroup. Does there always exist a lifting $G \subseteq \text{Diff}(\Sigma)$?
(Recall that $\text{MCG}(\Sigma) = \text{Diff}(\Sigma)/\text{isotopy}$).
- Kerckhoff (1983): Yes! Moreover, there exists a metric g such that $G \subseteq \text{Isom}(\Sigma, g)$. Or equivalently, G fixes a point in $\text{Teich}(\Sigma)$.
(There is a natural action of $\text{MCG}(\Sigma)$ on $\text{Teich}(\Sigma)$,
e.g. $\text{MCG}(T^2) = \text{SL}(2, \mathbb{Z})$ acts on $\text{Teich}(T^2) = \mathbb{H}$.)
(Rephrase: any finite subgroup of $\text{MCG}(\Sigma)$ can be realized as symmetries with respect to a metric on Σ .)
- Farb–Looijenga (2021) also proved similar statements for K3 surfaces (under certain conditions), where g is replaced by complex structure or Ricci-flat metric on the K3 surface.

Categorical realization problems

Let \mathcal{D} be a triangulated category, and $G \subseteq \mathrm{Aut}(\mathcal{D})$ be a finite subgroup. Does there exist $\sigma \in \mathrm{Stab}(\mathcal{D})$ such that $\Phi \cdot \sigma = \sigma$ for all $\Phi \in G$?

(Rephrase: any finite subgroup of $\mathrm{Aut}(\mathcal{D})$ can be realized as symmetries with respect to a stability condition on \mathcal{D} .)

- When $\mathcal{D} = D^b(X)$, stability conditions on \mathcal{D} are roughly Kähler structures on X ; so this problem is similar to (but not quite the same) the mirror problem of Farb–Looijenga.

Let $G \subseteq \mathrm{Aut}(\mathcal{D})/[1]$ be a finite subgroup.

Does there exist $\sigma \in \mathrm{Stab}(\mathcal{D})/\mathbb{C}$ such that $\Phi \cdot \sigma = \sigma$ for all $\Phi \in G$?

- There are various examples of \mathcal{D} where there are not many interesting finite order elements in $\mathrm{Aut}(\mathcal{D})$, but there are many interesting finite order elements in $\mathrm{Aut}(\mathcal{D})/[1]$.
- The shift functor $[1]$ (or rather $[2]$) has no classical counterpart in Teichmüller theory, so it is natural to consider this stronger question.

Categorical realization problems

Let \mathcal{D} be a triangulated category, and $G \subseteq \operatorname{Aut}(\mathcal{D})$ be a finite subgroup. Does there exist $\sigma \in \operatorname{Stab}(\mathcal{D})$ such that $\Phi \cdot \sigma = \sigma$ for all $\Phi \in G$?

(Rephrase: any finite subgroup of $\operatorname{Aut}(\mathcal{D})$ can be realized as symmetries with respect to a stability condition on \mathcal{D} .)

- When $\mathcal{D} = D^b(X)$, stability conditions on \mathcal{D} are roughly Kähler structures on X ; so this problem is similar to (but not quite the same) the mirror problem of Farb–Looijenga.

Let $G \subseteq \operatorname{Aut}(\mathcal{D})/[1]$ be a finite subgroup.

Does there exist $\sigma \in \operatorname{Stab}(\mathcal{D})/\mathbb{C}$ such that $\Phi \cdot \sigma = \sigma$ for all $\Phi \in G$?

- There are various examples of \mathcal{D} where there are not many interesting finite order elements in $\operatorname{Aut}(\mathcal{D})$, but there are many interesting finite order elements in $\operatorname{Aut}(\mathcal{D})/[1]$.
- The shift functor $[1]$ (or rather $[2]$) has no classical counterpart in Teichmüller theory, so it is natural to consider this stronger question.

Categorical realization problems

Let \mathcal{D} be a triangulated category, and $G \subseteq \mathrm{Aut}(\mathcal{D})$ be a finite subgroup. Does there exist $\sigma \in \mathrm{Stab}(\mathcal{D})$ such that $\Phi \cdot \sigma = \sigma$ for all $\Phi \in G$?

(Rephrase: any finite subgroup of $\mathrm{Aut}(\mathcal{D})$ can be realized as symmetries with respect to a stability condition on \mathcal{D} .)

- When $\mathcal{D} = D^b(X)$, stability conditions on \mathcal{D} are roughly Kähler structures on X ; so this problem is similar to (but not quite the same) the mirror problem of Farb–Looijenga.

Let $G \subseteq \mathrm{Aut}(\mathcal{D})/[1]$ be a finite subgroup.

Does there exist $\sigma \in \mathrm{Stab}(\mathcal{D})/\mathbb{C}$ such that $\Phi \cdot \sigma = \sigma$ for all $\Phi \in G$?

- There are various examples of \mathcal{D} where there are not many interesting finite order elements in $\mathrm{Aut}(\mathcal{D})$, but there are many interesting finite order elements in $\mathrm{Aut}(\mathcal{D})/[1]$.
- The shift functor $[1]$ (or rather $[2]$) has no classical counterpart in Teichmüller theory, so it is natural to consider this stronger question.

Categorical realization problems

Let \mathcal{D} be a triangulated category, and $G \subseteq \operatorname{Aut}(\mathcal{D})$ be a finite subgroup. Does there exist $\sigma \in \operatorname{Stab}(\mathcal{D})$ such that $\Phi \cdot \sigma = \sigma$ for all $\Phi \in G$?

(Rephrase: any finite subgroup of $\operatorname{Aut}(\mathcal{D})$ can be realized as symmetries with respect to a stability condition on \mathcal{D} .)

- When $\mathcal{D} = D^b(X)$, stability conditions on \mathcal{D} are roughly Kähler structures on X ; so this problem is similar to (but not quite the same) the mirror problem of Farb–Looijenga.

Let $G \subseteq \operatorname{Aut}(\mathcal{D})/[1]$ be a finite subgroup.

Does there exist $\sigma \in \operatorname{Stab}(\mathcal{D})/\mathbb{C}$ such that $\Phi \cdot \sigma = \sigma$ for all $\Phi \in G$?

- There are various examples of \mathcal{D} where there are not many interesting finite order elements in $\operatorname{Aut}(\mathcal{D})$, but there are many interesting finite order elements in $\operatorname{Aut}(\mathcal{D})/[1]$.
- The shift functor $[1]$ (or rather $[2]$) has no classical counterpart in Teichmüller theory, so it is natural to consider this stronger question.

Categorical realization problems

Let \mathcal{D} be a triangulated category, and $G \subseteq \mathrm{Aut}(\mathcal{D})$ be a finite subgroup. Does there exist $\sigma \in \mathrm{Stab}(\mathcal{D})$ such that $\Phi \cdot \sigma = \sigma$ for all $\Phi \in G$?

(Rephrase: any finite subgroup of $\mathrm{Aut}(\mathcal{D})$ can be realized as symmetries with respect to a stability condition on \mathcal{D} .)

- When $\mathcal{D} = D^b(X)$, stability conditions on \mathcal{D} are roughly Kähler structures on X ; so this problem is similar to (but not quite the same) the mirror problem of Farb–Looijenga.

Let $G \subseteq \mathrm{Aut}(\mathcal{D})/[1]$ be a finite subgroup.

Does there exist $\sigma \in \mathrm{Stab}(\mathcal{D})/\mathbb{C}$ such that $\Phi \cdot \sigma = \sigma$ for all $\Phi \in G$?

- There are various examples of \mathcal{D} where there are not many interesting finite order elements in $\mathrm{Aut}(\mathcal{D})$, but there are many interesting finite order elements in $\mathrm{Aut}(\mathcal{D})/[1]$.
- The shift functor $[1]$ (or rather $[2]$) has no classical counterpart in Teichmüller theory, so it is natural to consider this stronger question.

Categorical realization problems

Let \mathcal{D} be a triangulated category, and $G \subseteq \mathrm{Aut}(\mathcal{D})$ be a finite subgroup. Does there exist $\sigma \in \mathrm{Stab}(\mathcal{D})$ such that $\Phi \cdot \sigma = \sigma$ for all $\Phi \in G$?

(Rephrase: any finite subgroup of $\mathrm{Aut}(\mathcal{D})$ can be realized as symmetries with respect to a stability condition on \mathcal{D} .)

- When $\mathcal{D} = D^b(X)$, stability conditions on \mathcal{D} are roughly Kähler structures on X ; so this problem is similar to (but not quite the same) the mirror problem of Farb–Looijenga.

Let $G \subseteq \mathrm{Aut}(\mathcal{D})/[1]$ be a finite subgroup.

Does there exist $\sigma \in \mathrm{Stab}(\mathcal{D})/\mathbb{C}$ such that $\Phi \cdot \sigma = \sigma$ for all $\Phi \in G$?

- There are various examples of \mathcal{D} where there are not many interesting finite order elements in $\mathrm{Aut}(\mathcal{D})$, but there are many interesting finite order elements in $\mathrm{Aut}(\mathcal{D})/[1]$.
- The shift functor $[1]$ (or rather $[2]$) has no classical counterpart in Teichmüller theory, so it is natural to consider this stronger question.

Main theorems (F.–Lai, 2023)

- The answers to both problems are yes, for $D = D^b\mathrm{Coh}(X)$ where X is a curve, a (twisted) abelian surface, a generic twisted K3 surface, or a K3 surface of Picard number $\rho = 1$.

For K3 surfaces of $\rho = 1$, we obtain:

- every finite subgroup of $\mathrm{Aut}(D)$ is of **order 2**, and is generated by an **anti-symplectic involution**;
- classification and counting formula of the conjugacy classes of finite subgroups of $\mathrm{Aut}(D)$ and $\mathrm{Aut}(D)/[1]$;
- one-to-one correspondence between $\{\text{maximal finite subgroups of } \mathrm{Aut}(D)/[1]\}$ and $\{\text{elliptic points of } \mathrm{Stab}_{\mathrm{red}}^{\dagger}(D)/\mathbb{C}\}$
(analogue: one-to-one correspondence between $\{\text{maximal finite subgroups of } \mathrm{PSL}(2, \mathbb{Z})\}$ and $\{\text{elliptic points of } \mathbb{H}\}$)

Here, $\mathrm{Stab}_{\mathrm{red}}^{\dagger}(D) = \{(Z, P) \mid Z^2 = 0 \text{ in } N(D) \otimes \mathbb{C}\} \subseteq \mathrm{Stab}^{\dagger}(D)$.

Main theorems (F.–Lai, 2023)

- The answers to both problems are yes, for $D = D^b\mathrm{Coh}(X)$ where X is a curve, a (twisted) abelian surface, a generic twisted K3 surface, or a K3 surface of Picard number $\rho = 1$.

For K3 surfaces of $\rho = 1$, we obtain:

- every finite subgroup of $\mathrm{Aut}(D)$ is of order 2, and is generated by an anti-symplectic involution;
- classification and counting formula of the conjugacy classes of finite subgroups of $\mathrm{Aut}(D)$ and $\mathrm{Aut}(D)/[1]$;
- one-to-one correspondence between $\{\text{maximal finite subgroups of } \mathrm{Aut}(D)/[1]\}$ and $\{\text{elliptic points of } \mathrm{Stab}_{\mathrm{red}}^{\dagger}(D)/\mathbb{C}\}$
(analogue: one-to-one correspondence between $\{\text{maximal finite subgroups of } \mathrm{PSL}(2, \mathbb{Z})\}$ and $\{\text{elliptic points of } \mathbb{H}\})$

Here, $\mathrm{Stab}_{\mathrm{red}}^{\dagger}(D) = \{(Z, P) \mid Z^2 = 0 \text{ in } N(D) \otimes \mathbb{C}\} \subseteq \mathrm{Stab}^{\dagger}(D)$.

Main theorems (F.–Lai, 2023)

- The answers to both problems are yes, for $D = D^b\mathrm{Coh}(X)$ where X is a curve, a (twisted) abelian surface, a generic twisted K3 surface, or a K3 surface of Picard number $\rho = 1$.

For K3 surfaces of $\rho = 1$, we obtain:

- every finite subgroup of $\mathrm{Aut}(D)$ is of **order 2**, and is generated by an **anti-symplectic involution**;
- classification and counting formula of the conjugacy classes of finite subgroups of $\mathrm{Aut}(D)$ and $\mathrm{Aut}(D)/[1]$;
- one-to-one correspondence between $\{\text{maximal finite subgroups of } \mathrm{Aut}(D)/[1]\}$ and $\{\text{elliptic points of } \mathrm{Stab}_{\mathrm{red}}^{\dagger}(D)/\mathbb{C}\}$
(analogue: one-to-one correspondence between $\{\text{maximal finite subgroups of } \mathrm{PSL}(2, \mathbb{Z})\}$ and $\{\text{elliptic points of } \mathbb{H}\}$)

Here, $\mathrm{Stab}_{\mathrm{red}}^{\dagger}(D) = \{(Z, P) \mid Z^2 = 0 \text{ in } N(D) \otimes \mathbb{C}\} \subseteq \mathrm{Stab}^{\dagger}(D)$.

Main theorems (F.–Lai, 2023)

- The answers to both problems are yes, for $D = D^b\mathrm{Coh}(X)$ where X is a curve, a (twisted) abelian surface, a generic twisted K3 surface, or a K3 surface of Picard number $\rho = 1$.

For K3 surfaces of $\rho = 1$, we obtain:

- every finite subgroup of $\mathrm{Aut}(D)$ is of **order 2**, and is generated by an **anti-symplectic involution**;
- classification and counting formula of the conjugacy classes of finite subgroups of $\mathrm{Aut}(D)$ and $\mathrm{Aut}(D)/[1]$;
- one-to-one correspondence between $\{\text{maximal finite subgroups of } \mathrm{Aut}(D)/[1]\}$ and $\{\text{elliptic points of } \mathrm{Stab}_{\mathrm{red}}^{\dagger}(D)/\mathbb{C}\}$
(analogue: one-to-one correspondence between $\{\text{maximal finite subgroups of } \mathrm{PSL}(2, \mathbb{Z})\}$ and $\{\text{elliptic points of } \mathbb{H}\})$

Here, $\mathrm{Stab}_{\mathrm{red}}^{\dagger}(D) = \{(Z, P) \mid Z^2 = 0 \text{ in } N(D) \otimes \mathbb{C}\} \subseteq \mathrm{Stab}^{\dagger}(D)$.

Main theorems (F.–Lai, 2023)

- The answers to both problems are yes, for $D = D^b\mathrm{Coh}(X)$ where X is a curve, a (twisted) abelian surface, a generic twisted K3 surface, or a K3 surface of Picard number $\rho = 1$.

For K3 surfaces of $\rho = 1$, we obtain:

- every finite subgroup of $\mathrm{Aut}(D)$ is of **order 2**, and is generated by an **anti-symplectic involution**;
- classification and counting formula of the conjugacy classes of finite subgroups of $\mathrm{Aut}(D)$ and $\mathrm{Aut}(D)/[1]$;
- one-to-one correspondence between $\{\text{maximal finite subgroups of } \mathrm{Aut}(D)/[1]\}$ and $\{\text{elliptic points of } \mathrm{Stab}_{\mathrm{red}}^{\dagger}(D)/\mathbb{C}\}$
(analogue: one-to-one correspondence between $\{\text{maximal finite subgroups of } \mathrm{PSL}(2, \mathbb{Z})\}$ and $\{\text{elliptic points of } \mathbb{H}\})$

Here, $\mathrm{Stab}_{\mathrm{red}}^{\dagger}(D) = \{(Z, P) \mid Z^2 = 0 \text{ in } N(D) \otimes \mathbb{C}\} \subseteq \mathrm{Stab}^{\dagger}(D)$.

Main theorems (F.–Lai, 2023), cont'd

It is well-known that the rationality of a cubic fourfold Y is closely related to the existence of an associated K3 surface. For instance, Kuznetsov conjectured that Y is rational if and only if there exists a K3 surface X such that $\mathrm{Ku}(Y) \cong D^b(X)$.

On the other hand, not every K3 surface can be associated with a cubic fourfold: there is a functor on $\mathrm{Ku}(Y)$

$$T_Y: E \mapsto j^*(E \otimes \mathcal{O}(1))$$

which satisfies $T_Y^3 = [2]$. It is asked by Huybrechts whether the existence of such order 3 element of $\mathrm{Aut}(D^b(X))/[2]$ characterizes K3 surfaces with associated cubic fourfolds.

- We prove that when X is a K3 surface of Picard number one, the group $\mathrm{Aut}(D^b(X))/[2]$ contains an order 3 element if and only if it admits an associated cubic fourfold.

Main theorems (F.–Lai, 2023), cont'd

It is well-known that the rationality of a cubic fourfold Y is closely related to the existence of an associated K3 surface. For instance, Kuznetsov conjectured that Y is rational if and only if there exists a K3 surface X such that $\mathrm{Ku}(Y) \cong D^b(X)$.

On the other hand, not every K3 surface can be associated with a cubic fourfold: there is a functor on $\mathrm{Ku}(Y)$

$$T_Y: E \mapsto j^*(E \otimes \mathcal{O}(1))$$

which satisfies $T_Y^3 = [2]$. It is asked by Huybrechts whether the existence of such order 3 element of $\mathrm{Aut}(D^b(X))/[2]$ characterizes K3 surfaces with associated cubic fourfolds.

- We prove that when X is a K3 surface of Picard number one, the group $\mathrm{Aut}(D^b(X))/[2]$ contains an order 3 element if and only if it admits an associated cubic fourfold.

Main theorems (F.–Lai, 2023), cont'd

It is well-known that the rationality of a cubic fourfold Y is closely related to the existence of an associated K3 surface. For instance, Kuznetsov conjectured that Y is rational if and only if there exists a K3 surface X such that $\mathrm{Ku}(Y) \cong D^b(X)$.

On the other hand, not every K3 surface can be associated with a cubic fourfold: there is a functor on $\mathrm{Ku}(Y)$

$$T_Y: E \mapsto j^*(E \otimes \mathcal{O}(1))$$

which satisfies $T_Y^3 = [2]$. It is asked by Huybrechts whether the existence of such order 3 element of $\mathrm{Aut}(D^b(X))/[2]$ characterizes K3 surfaces with associated cubic fourfolds.

- We prove that when X is a K3 surface of Picard number one, the group $\mathrm{Aut}(D^b(X))/[2]$ contains an order 3 element if and only if it admits an associated cubic fourfold.

Main theorems (F.–Lai, 2023), cont'd

It is well-known that the rationality of a cubic fourfold Y is closely related to the existence of an associated K3 surface. For instance, Kuznetsov conjectured that Y is rational if and only if there exists a K3 surface X such that $\mathrm{Ku}(Y) \cong D^b(X)$.

On the other hand, not every K3 surface can be associated with a cubic fourfold: there is a functor on $\mathrm{Ku}(Y)$

$$T_Y: E \mapsto j^*(E \otimes \mathcal{O}(1))$$

which satisfies $T_Y^3 = [2]$. It is asked by Huybrechts whether the existence of such order 3 element of $\mathrm{Aut}(D^b(X))/[2]$ characterizes K3 surfaces with associated cubic fourfolds.

- We prove that when X is a K3 surface of Picard number one, the group $\mathrm{Aut}(D^b(X))/[2]$ contains an order 3 element if and only if it admits an associated cubic fourfold.

Main theorems (F.–Lai, 2023), cont'd

It is well-known that the rationality of a cubic fourfold Y is closely related to the existence of an associated K3 surface. For instance, Kuznetsov conjectured that Y is rational if and only if there exists a K3 surface X such that $\mathrm{Ku}(Y) \cong D^b(X)$.

On the other hand, not every K3 surface can be associated with a cubic fourfold: there is a functor on $\mathrm{Ku}(Y)$

$$T_Y: E \mapsto j^*(E \otimes \mathcal{O}(1))$$

which satisfies $T_Y^3 = [2]$. It is asked by Huybrechts whether the existence of such order 3 element of $\mathrm{Aut}(D^b(X))/[2]$ characterizes K3 surfaces with associated cubic fourfolds.

- We prove that when X is a K3 surface of Picard number one, the group $\mathrm{Aut}(D^b(X))/[2]$ contains an order 3 element if and only if it admits an associated cubic fourfold.

Main theorems (F.–Lai, 2023), cont'd

It is well-known that the rationality of a cubic fourfold Y is closely related to the existence of an associated K3 surface. For instance, Kuznetsov conjectured that Y is rational if and only if there exists a K3 surface X such that $\mathrm{Ku}(Y) \cong D^b(X)$.

On the other hand, not every K3 surface can be associated with a cubic fourfold: there is a functor on $\mathrm{Ku}(Y)$

$$T_Y: E \mapsto j^*(E \otimes \mathcal{O}(1))$$

which satisfies $T_Y^3 = [2]$. It is asked by Huybrechts whether the existence of such order 3 element of $\mathrm{Aut}(D^b(X))/[2]$ characterizes K3 surfaces with associated cubic fourfolds.

- We prove that when X is a K3 surface of Picard number one, the group $\mathrm{Aut}(D^b(X))/[2]$ contains an order 3 element if and only if it admits an associated cubic fourfold.

Main theorems (F.–Lai, 2023), cont'd

$\Phi \in \text{Aut}(D)$ can be classified into (modulo quotienting certain subgroup):

- finite order up to shifts
- reducible, which further classified into:
 - ▶ “ (-2) -reducible”: spherical twists T_S
 - ▶ “0-reducible”: which fixes a class $w \in N(D)$ with $w^2 = 0$ (e.g. $\otimes \mathcal{O}(1)$)
- hyperbolic: $\rho([\Phi]_{N(D)}) > 1$

Modulo certain conjectures regarding the (polynomial) entropy of the reducible autoequivalences, we expect the following trichotomy:

- finite order if and only if $h_{\text{cat}} = h_{\text{poly}} = 0$
- reducible if and only if $h_{\text{cat}} = 0$ and $h_{\text{poly}} > 0$
- hyperbolic if and only if $h_{\text{cat}} > 0$

Main theorems (F.–Lai, 2023), cont'd

$\Phi \in \text{Aut}(D)$ can be classified into (modulo quotienting certain subgroup):

- finite order up to shifts
- reducible, which further classified into:
 - ▶ “ (-2) -reducible”: spherical twists T_S
 - ▶ “0-reducible”: which fixes a class $w \in N(D)$ with $w^2 = 0$ (e.g. $\otimes \mathcal{O}(1)$)
- hyperbolic: $\rho([\Phi]_{N(D)}) > 1$

Modulo certain conjectures regarding the (polynomial) entropy of the reducible autoequivalences, we expect the following trichotomy:

- finite order if and only if $h_{\text{cat}} = h_{\text{poly}} = 0$
- reducible if and only if $h_{\text{cat}} = 0$ and $h_{\text{poly}} > 0$
- hyperbolic if and only if $h_{\text{cat}} > 0$

Main theorems (F.–Lai, 2023), cont'd

$\Phi \in \text{Aut}(D)$ can be classified into (modulo quotienting certain subgroup):

- finite order up to shifts
- reducible, which further classified into:
 - ▶ “ (-2) -reducible”: spherical twists T_S
 - ▶ “0-reducible”: which fixes a class $w \in N(D)$ with $w^2 = 0$ (e.g. $\otimes \mathcal{O}(1)$)
- hyperbolic: $\rho([\Phi]_{N(D)}) > 1$

Modulo certain conjectures regarding the (polynomial) entropy of the reducible autoequivalences, we expect the following trichotomy:

- finite order if and only if $h_{\text{cat}} = h_{\text{poly}} = 0$
- reducible if and only if $h_{\text{cat}} = 0$ and $h_{\text{poly}} > 0$
- hyperbolic if and only if $h_{\text{cat}} > 0$

Main theorems (F.–Lai, 2023), cont'd

$\Phi \in \text{Aut}(D)$ can be classified into (modulo quotienting certain subgroup):

- finite order up to shifts
- reducible, which further classified into:
 - ▶ “ (-2) -reducible”: spherical twists T_S
 - ▶ “0-reducible”: which fixes a class $w \in N(D)$ with $w^2 = 0$ (e.g. $\otimes \mathcal{O}(1)$)
- hyperbolic: $\rho([\Phi]_{N(D)}) > 1$

Modulo certain conjectures regarding the (polynomial) entropy of the reducible autoequivalences, we expect the following trichotomy:

- finite order if and only if $h_{\text{cat}} = h_{\text{poly}} = 0$
- reducible if and only if $h_{\text{cat}} = 0$ and $h_{\text{poly}} > 0$
- hyperbolic if and only if $h_{\text{cat}} > 0$

Main theorems (F.–Lai, 2023), cont'd

$\Phi \in \text{Aut}(D)$ can be classified into (modulo quotienting certain subgroup):

- finite order up to shifts
- reducible, which further classified into:
 - ▶ “ (-2) -reducible”: spherical twists T_S
 - ▶ “0-reducible”: which fixes a class $w \in N(D)$ with $w^2 = 0$ (e.g. $\otimes \mathcal{O}(1)$)
- hyperbolic: $\rho([\Phi]_{N(D)}) > 1$

Modulo certain conjectures regarding the (polynomial) entropy of the reducible autoequivalences, we expect the following trichotomy:

- finite order if and only if $h_{\text{cat}} = h_{\text{poly}} = 0$
- reducible if and only if $h_{\text{cat}} = 0$ and $h_{\text{poly}} > 0$
- hyperbolic if and only if $h_{\text{cat}} > 0$

Main theorems (F.–Lai, 2023), cont'd

$\Phi \in \text{Aut}(D)$ can be classified into (modulo quotienting certain subgroup):

- finite order up to shifts
- reducible, which further classified into:
 - ▶ “ (-2) -reducible”: spherical twists T_S
 - ▶ “0-reducible”: which fixes a class $w \in N(D)$ with $w^2 = 0$ (e.g. $\otimes \mathcal{O}(1)$)
- hyperbolic: $\rho([\Phi]_{N(D)}) > 1$

Modulo certain conjectures regarding the (polynomial) entropy of the reducible autoequivalences, we expect the following trichotomy:

- finite order if and only if $h_{\text{cat}} = h_{\text{poly}} = 0$
- reducible if and only if $h_{\text{cat}} = 0$ and $h_{\text{poly}} > 0$
- hyperbolic if and only if $h_{\text{cat}} > 0$

Strategy and Difficulties

For K3 or abelian surfaces, Bridgeland (2008) showed that there is an $\text{Aut}(D)$ -equivariant covering map

$$\text{Stab}_{\text{red}}^{\dagger}(D)/\mathbb{C} \xrightarrow{\pi} Q_0^+(D)$$

where $Q_0^+(D) = \{v \in \mathbb{P}(N(D) \otimes \mathbb{C}) \mid v^2 = 0, v\bar{v} > 0\} \setminus \bigcup_{\delta^2 = -2} \delta^{\perp}$.

- For abelian surfaces, there is no spherical objects in D , so:
 - ▶ we do not need to remove “ δ^{\perp} ” since there is no (-2) -classes
 - ▶ π is an isomorphism

and it is not hard to show that finite subgroups of $\text{Aut}(D)$ fix a point in $Q_0^+(D)$ using basic Lie theory.

- For K3 surfaces, one needs to resolve the following two issues:
 - ▶ the fixed points of finite subgroups of $\text{Aut}(D)$ need to avoid “ δ^{\perp} ”
 - ▶ $\text{Deck}(\pi)$ is much more complicated

It turns out that both issues can be resolved for $\rho = 1$ –

Strategy and Difficulties

For K3 or abelian surfaces, Bridgeland (2008) showed that there is an $\text{Aut}(D)$ -equivariant covering map

$$\text{Stab}_{\text{red}}^+(D)/\mathbb{C} \xrightarrow{\pi} Q_0^+(D)$$

where $Q_0^+(D) = \{v \in \mathbb{P}(N(D) \otimes \mathbb{C}) \mid v^2 = 0, v\bar{v} > 0\} \setminus \bigcup_{\delta^2 = -2} \delta^\perp$.

- For abelian surfaces, there is no spherical objects in D , so:

- ▶ we do not need to remove " δ^\perp " since there is no (-2) -classes
- ▶ π is an isomorphism

and it is not hard to show that finite subgroups of $\text{Aut}(D)$ fix a point in $Q_0^+(D)$ using basic Lie theory.

- For K3 surfaces, one needs to resolve the following two issues:

- ▶ the fixed points of finite subgroups of $\text{Aut}(D)$ need to avoid " δ^\perp "
- ▶ $\text{Deck}(\pi)$ is much more complicated

It turns out that both issues can be resolved for $\rho = 1$ –

Strategy and Difficulties

For K3 or abelian surfaces, Bridgeland (2008) showed that there is an $\text{Aut}(D)$ -equivariant covering map

$$\text{Stab}_{\text{red}}^{\dagger}(D)/\mathbb{C} \xrightarrow{\pi} Q_0^+(D)$$

where $Q_0^+(D) = \{v \in \mathbb{P}(N(D) \otimes \mathbb{C}) \mid v^2 = 0, v\bar{v} > 0\} \setminus \bigcup_{\delta^2 = -2} \delta^{\perp}$.

- For abelian surfaces, there is no spherical objects in D , so:
 - ▶ we do not need to remove “ δ^{\perp} ” since there is no (-2) -classes
 - ▶ π is an isomorphism

and it is not hard to show that finite subgroups of $\text{Aut}(D)$ fix a point in $Q_0^+(D)$ using basic Lie theory.

- For K3 surfaces, one needs to resolve the following two issues:
 - ▶ the fixed points of finite subgroups of $\text{Aut}(D)$ need to avoid “ δ^{\perp} ”
 - ▶ $\text{Deck}(\pi)$ is much more complicated

It turns out that both issues can be resolved for $\rho = 1$ –

Strategy and Difficulties

For K3 or abelian surfaces, Bridgeland (2008) showed that there is an $\text{Aut}(D)$ -equivariant covering map

$$\text{Stab}_{\text{red}}^{\dagger}(D)/\mathbb{C} \xrightarrow{\pi} Q_0^+(D)$$

where $Q_0^+(D) = \{v \in \mathbb{P}(N(D) \otimes \mathbb{C}) \mid v^2 = 0, v\bar{v} > 0\} \setminus \bigcup_{\delta^2 = -2} \delta^{\perp}$.

- For abelian surfaces, there is no spherical objects in D , so:
 - ▶ we do not need to remove “ δ^{\perp} ” since there is no (-2) -classes
 - ▶ π is an isomorphism

and it is not hard to show that finite subgroups of $\text{Aut}(D)$ fix a point in $Q_0^+(D)$ using basic Lie theory.

- For K3 surfaces, one needs to resolve the following two issues:
 - ▶ the fixed points of finite subgroups of $\text{Aut}(D)$ need to avoid “ δ^{\perp} ”
 - ▶ $\text{Deck}(\pi)$ is much more complicated

It turns out that both issues can be resolved for $\rho = 1 -$

Strategy and Difficulties

For K3 or abelian surfaces, Bridgeland (2008) showed that there is an $\text{Aut}(D)$ -equivariant covering map

$$\text{Stab}_{\text{red}}^{\dagger}(D)/\mathbb{C} \xrightarrow{\pi} Q_0^+(D)$$

where $Q_0^+(D) = \{v \in \mathbb{P}(N(D) \otimes \mathbb{C}) \mid v^2 = 0, v\bar{v} > 0\} \setminus \bigcup_{\delta^2 = -2} \delta^{\perp}$.

- For abelian surfaces, there is no spherical objects in D , so:
 - ▶ we do not need to remove “ δ^{\perp} ” since there is no (-2) -classes
 - ▶ π is an isomorphism

and it is not hard to show that finite subgroups of $\text{Aut}(D)$ fix a point in $Q_0^+(D)$ using basic Lie theory.

- For K3 surfaces, one needs to resolve the following two issues:
 - ▶ the fixed points of finite subgroups of $\text{Aut}(D)$ need to avoid “ δ^{\perp} ”
 - ▶ $\text{Deck}(\pi)$ is much more complicated

It turns out that both issues can be resolved for $\rho = 1$ –

Strategy and Difficulties

For K3 or abelian surfaces, Bridgeland (2008) showed that there is an $\text{Aut}(D)$ -equivariant covering map

$$\text{Stab}_{\text{red}}^{\dagger}(D)/\mathbb{C} \xrightarrow{\pi} Q_0^+(D)$$

where $Q_0^+(D) = \{v \in \mathbb{P}(N(D) \otimes \mathbb{C}) \mid v^2 = 0, v\bar{v} > 0\} \setminus \bigcup_{\delta^2 = -2} \delta^{\perp}$.

- For abelian surfaces, there is no spherical objects in D , so:
 - ▶ we do not need to remove “ δ^{\perp} ” since there is no (-2) -classes
 - ▶ π is an isomorphism

and it is not hard to show that finite subgroups of $\text{Aut}(D)$ fix a point in $Q_0^+(D)$ using basic Lie theory.

- For K3 surfaces, one needs to resolve the following two issues:
 - ▶ the fixed points of finite subgroups of $\text{Aut}(D)$ need to avoid “ δ^{\perp} ”
 - ▶ $\text{Deck}(\pi)$ is much more complicated

It turns out that both issues can be resolved for $\rho = 1$ –

Strategy and Difficulties

For K3 or abelian surfaces, Bridgeland (2008) showed that there is an $\text{Aut}(D)$ -equivariant covering map

$$\text{Stab}_{\text{red}}^{\dagger}(D)/\mathbb{C} \xrightarrow{\pi} Q_0^+(D)$$

where $Q_0^+(D) = \{v \in \mathbb{P}(N(D) \otimes \mathbb{C}) \mid v^2 = 0, v\bar{v} > 0\} \setminus \bigcup_{\delta^2 = -2} \delta^{\perp}$.

- For abelian surfaces, there is no spherical objects in D , so:
 - ▶ we do not need to remove “ δ^{\perp} ” since there is no (-2) -classes
 - ▶ π is an isomorphism

and it is not hard to show that finite subgroups of $\text{Aut}(D)$ fix a point in $Q_0^+(D)$ using basic Lie theory.

- For K3 surfaces, one needs to resolve the following two issues:
 - ▶ the fixed points of finite subgroups of $\text{Aut}(D)$ need to avoid “ δ^{\perp} ”
 - ▶ $\text{Deck}(\pi)$ is much more complicated

It turns out that both issues can be resolved for $\rho = 1$ –

Strategy and Difficulties

For K3 or abelian surfaces, Bridgeland (2008) showed that there is an $\text{Aut}(D)$ -equivariant covering map

$$\text{Stab}_{\text{red}}^{\dagger}(D)/\mathbb{C} \xrightarrow{\pi} Q_0^+(D)$$

where $Q_0^+(D) = \{v \in \mathbb{P}(N(D) \otimes \mathbb{C}) \mid v^2 = 0, v\bar{v} > 0\} \setminus \bigcup_{\delta^2 = -2} \delta^{\perp}$.

- For abelian surfaces, there is no spherical objects in D , so:
 - ▶ we do not need to remove “ δ^{\perp} ” since there is no (-2) -classes
 - ▶ π is an isomorphism

and it is not hard to show that finite subgroups of $\text{Aut}(D)$ fix a point in $Q_0^+(D)$ using basic Lie theory.

- For K3 surfaces, one needs to resolve the following two issues:
 - ▶ the fixed points of finite subgroups of $\text{Aut}(D)$ need to avoid “ δ^{\perp} ”
 - ▶ $\text{Deck}(\pi)$ is much more complicated

It turns out that both issues can be resolved for $\rho = 1$ –

Strategy and Difficulties

For K3 or abelian surfaces, Bridgeland (2008) showed that there is an $\text{Aut}(D)$ -equivariant covering map

$$\text{Stab}_{\text{red}}^{\dagger}(D)/\mathbb{C} \xrightarrow{\pi} Q_0^+(D)$$

where $Q_0^+(D) = \{v \in \mathbb{P}(N(D) \otimes \mathbb{C}) \mid v^2 = 0, v\bar{v} > 0\} \setminus \bigcup_{\delta^2 = -2} \delta^{\perp}$.

- For abelian surfaces, there is no spherical objects in D , so:
 - ▶ we do not need to remove “ δ^{\perp} ” since there is no (-2) -classes
 - ▶ π is an isomorphism

and it is not hard to show that finite subgroups of $\text{Aut}(D)$ fix a point in $Q_0^+(D)$ using basic Lie theory.

- For K3 surfaces, one needs to resolve the following two issues:
 - ▶ the fixed points of finite subgroups of $\text{Aut}(D)$ need to avoid “ δ^{\perp} ”
 - ▶ $\text{Deck}(\pi)$ is much more complicated

It turns out that both issues can be resolved for $\rho = 1$ –

Avoiding δ^\perp

Suppose X is a K3 surface of $\rho = 1$ and degree $2n$.

- We have $Q_0^+(D) \cong \mathbb{H} \setminus \text{"}(-2)\text{"-points"}$.
- By Dolgachev (1996) and Kawatani (2014), the action of $\text{Aut}(D)$ on $Q_0^+(D)$ factors through $\text{Im}(\text{Aut}(D) \xrightarrow{f} \text{PSL}(2, \mathbb{R})) = \Gamma_0^+(n)$ the Fricke modular group, where $\Gamma_0^+(n) = \left\langle \Gamma_0(n), \begin{bmatrix} & -1/\sqrt{n} \\ \sqrt{n} & \end{bmatrix} =: \omega_n \right\rangle$.

We showed that the following statements are equivalent:

- $f(\Phi)$ fixes a (-2) -point in \mathbb{H}
- $f(\Phi)$ is an involution, and $f(\Phi) = g_0 \omega_n$ for some $g_0 \in \Gamma_0(n)$
- $\Phi = T_S \Psi$ for some spherical object S and some $\Psi \in \text{Deck}(\pi)$

Moreover, we showed that autoequivalences of the form $T_S \Psi$ must be of infinite order in $\text{Aut}(D)/[1]$. This resolves the first issue.

Avoiding δ^\perp

Suppose X is a K3 surface of $\rho = 1$ and degree $2n$.

- We have $Q_0^+(D) \cong \mathbb{H} \setminus \text{"}(-2)\text{-points"}$.
- By Dolgachev (1996) and Kawatani (2014), the action of $\text{Aut}(D)$ on $Q_0^+(D)$ factors through $\text{Im}(\text{Aut}(D) \xrightarrow{f} \text{PSL}(2, \mathbb{R})) = \Gamma_0^+(n)$ the Fricke modular group, where $\Gamma_0^+(n) = \left\langle \Gamma_0(n), \begin{bmatrix} & -1/\sqrt{n} \\ \sqrt{n} & \end{bmatrix} =: \omega_n \right\rangle$.

We showed that the following statements are equivalent:

- $f(\Phi)$ fixes a (-2) -point in \mathbb{H}
- $f(\Phi)$ is an involution, and $f(\Phi) = g_0 \omega_n$ for some $g_0 \in \Gamma_0(n)$
- $\Phi = T_S \Psi$ for some spherical object S and some $\Psi \in \text{Deck}(\pi)$

Moreover, we showed that autoequivalences of the form $T_S \Psi$ must be of infinite order in $\text{Aut}(D)/[1]$. This resolves the first issue.

Avoiding δ^\perp

Suppose X is a K3 surface of $\rho = 1$ and degree $2n$.

- We have $Q_0^+(D) \cong \mathbb{H} \setminus \text{"}(-2)\text{-points"}$.
- By Dolgachev (1996) and Kawatani (2014), the action of $\text{Aut}(D)$ on $Q_0^+(D)$ factors through $\text{Im}(\text{Aut}(D) \xrightarrow{f} \text{PSL}(2, \mathbb{R})) = \Gamma_0^+(n)$ the Fricke modular group, where $\Gamma_0^+(n) = \left\langle \Gamma_0(n), \begin{bmatrix} & -1/\sqrt{n} \\ \sqrt{n} & \end{bmatrix} =: \omega_n \right\rangle$.

We showed that the following statements are equivalent:

- $f(\Phi)$ fixes a (-2) -point in \mathbb{H}
- $f(\Phi)$ is an involution, and $f(\Phi) = g_0 \omega_n$ for some $g_0 \in \Gamma_0(n)$
- $\Phi = T_S \Psi$ for some spherical object S and some $\Psi \in \text{Deck}(\pi)$

Moreover, we showed that autoequivalences of the form $T_S \Psi$ must be of infinite order in $\text{Aut}(D)/[1]$. This resolves the first issue.

Avoiding δ^\perp

Suppose X is a K3 surface of $\rho = 1$ and degree $2n$.

- We have $Q_0^+(D) \cong \mathbb{H} \setminus \text{"}(-2)\text{-points"}$.
- By Dolgachev (1996) and Kawatani (2014), the action of $\text{Aut}(D)$ on $Q_0^+(D)$ factors through $\text{Im}(\text{Aut}(D) \xrightarrow{f} \text{PSL}(2, \mathbb{R})) = \Gamma_0^+(n)$ the Fricke modular group, where $\Gamma_0^+(n) = \left\langle \Gamma_0(n), \begin{bmatrix} & -1/\sqrt{n} \\ \sqrt{n} & \end{bmatrix} =: \omega_n \right\rangle$.

We showed that the following statements are equivalent:

- $f(\Phi)$ fixes a (-2) -point in \mathbb{H}
- $f(\Phi)$ is an involution, and $f(\Phi) = g_0 \omega_n$ for some $g_0 \in \Gamma_0(n)$
- $\Phi = T_S \Psi$ for some spherical object S and some $\Psi \in \text{Deck}(\pi)$

Moreover, we showed that autoequivalences of the form $T_S \Psi$ must be of infinite order in $\text{Aut}(D)/[1]$. This resolves the first issue.

Avoiding δ^\perp

Suppose X is a K3 surface of $\rho = 1$ and degree $2n$.

- We have $Q_0^+(D) \cong \mathbb{H} \setminus \text{"}(-2)\text{-points"}$.
- By Dolgachev (1996) and Kawatani (2014), the action of $\text{Aut}(D)$ on $Q_0^+(D)$ factors through $\text{Im}(\text{Aut}(D) \xrightarrow{f} \text{PSL}(2, \mathbb{R})) = \Gamma_0^+(n)$ the Fricke modular group, where $\Gamma_0^+(n) = \left\langle \Gamma_0(n), \begin{bmatrix} & -1/\sqrt{n} \\ \sqrt{n} & \end{bmatrix} =: \omega_n \right\rangle$.

We showed that the following statements are equivalent:

- $f(\Phi)$ fixes a (-2) -point in \mathbb{H}
- $f(\Phi)$ is an involution, and $f(\Phi) = g_0 \omega_n$ for some $g_0 \in \Gamma_0(n)$
- $\Phi = T_S \Psi$ for some spherical object S and some $\Psi \in \text{Deck}(\pi)$

Moreover, we showed that autoequivalences of the form $T_S \Psi$ must be of infinite order in $\text{Aut}(D)/[1]$. This resolves the first issue.

Avoiding δ^\perp

Suppose X is a K3 surface of $\rho = 1$ and degree $2n$.

- We have $Q_0^+(D) \cong \mathbb{H} \setminus \text{"}(-2)\text{-points"}$.
- By Dolgachev (1996) and Kawatani (2014), the action of $\text{Aut}(D)$ on $Q_0^+(D)$ factors through $\text{Im}(\text{Aut}(D) \xrightarrow{f} \text{PSL}(2, \mathbb{R})) = \Gamma_0^+(n)$ the Fricke modular group, where $\Gamma_0^+(n) = \left\langle \Gamma_0(n), \begin{bmatrix} & -1/\sqrt{n} \\ \sqrt{n} & \end{bmatrix} =: \omega_n \right\rangle$.

We showed that the following statements are equivalent:

- $f(\Phi)$ fixes a (-2) -point in \mathbb{H}
- $f(\Phi)$ is an involution, and $f(\Phi) = g_0 \omega_n$ for some $g_0 \in \Gamma_0(n)$
- $\Phi = T_S \Psi$ for some spherical object S and some $\Psi \in \text{Deck}(\pi)$

Moreover, we showed that autoequivalences of the form $T_S \Psi$ must be of infinite order in $\text{Aut}(D)/[1]$. This resolves the first issue.

Avoiding δ^\perp

Suppose X is a K3 surface of $\rho = 1$ and degree $2n$.

- We have $Q_0^+(D) \cong \mathbb{H} \setminus \text{"}(-2)\text{-points"}$.
- By Dolgachev (1996) and Kawatani (2014), the action of $\text{Aut}(D)$ on $Q_0^+(D)$ factors through $\text{Im}(\text{Aut}(D) \xrightarrow{f} \text{PSL}(2, \mathbb{R})) = \Gamma_0^+(n)$ the Fricke modular group, where $\Gamma_0^+(n) = \left\langle \Gamma_0(n), \begin{bmatrix} & -1/\sqrt{n} \\ \sqrt{n} & \end{bmatrix} =: \omega_n \right\rangle$.

We showed that the following statements are equivalent:

- $f(\Phi)$ fixes a (-2) -point in \mathbb{H}
- $f(\Phi)$ is an involution, and $f(\Phi) = g_0 \omega_n$ for some $g_0 \in \Gamma_0(n)$
- $\Phi = T_S \Psi$ for some spherical object S and some $\Psi \in \text{Deck}(\pi)$

Moreover, we showed that autoequivalences of the form $T_S \Psi$ must be of infinite order in $\text{Aut}(D)/[1]$. This resolves the first issue.

Lifting of fixed points

- Let $\Phi \in \text{Aut}(D)/[1]$ be of finite order, then the previous discussion shows that it fixes a point in $Q_0^+(D)$.
- Kawatani (2019): $\pi_1(Q_0^+(D)) \cong \star_{\text{free}} T_S^2$.
- Bayer–Bridgeland (2017): $\text{Stab}_{\text{red}}^\dagger(D)/\mathbb{C}$ is contractible.
- Combining these two results, we have $\text{Deck}(\pi) \cong \star_{\text{free}} T_S^2$.
- We showed that this implies the fixed point of Φ in $Q_0^+(D)$ can be lifted to a fixed point in $\text{Stab}_{\text{red}}^\dagger(D)/\mathbb{C}$, which proves the realization problem for cyclic subgroups of $\text{Aut}(D)/[1]$.
- Finally, we showed that any finite subgroup of $\text{Aut}(D)/[1]$ is cyclic using classification results of finite subgroups of $\Gamma_0^+(n)$ by Fricke. This concludes the proof.

Lifting of fixed points

- Let $\Phi \in \text{Aut}(D)/[1]$ be of finite order, then the previous discussion shows that it fixes a point in $Q_0^+(D)$.
- Kawatani (2019): $\pi_1(Q_0^+(D)) \cong \star_{\text{free}} T_S^2$.
- Bayer–Bridgeland (2017): $\text{Stab}_{\text{red}}^\dagger(D)/\mathbb{C}$ is contractible.
- Combining these two results, we have $\text{Deck}(\pi) \cong \star_{\text{free}} T_S^2$.
- We showed that this implies the fixed point of Φ in $Q_0^+(D)$ can be lifted to a fixed point in $\text{Stab}_{\text{red}}^\dagger(D)/\mathbb{C}$, which proves the realization problem for cyclic subgroups of $\text{Aut}(D)/[1]$.
- Finally, we showed that any finite subgroup of $\text{Aut}(D)/[1]$ is cyclic using classification results of finite subgroups of $\Gamma_0^+(n)$ by Fricke. This concludes the proof.

Lifting of fixed points

- Let $\Phi \in \operatorname{Aut}(D)/[1]$ be of finite order, then the previous discussion shows that it fixes a point in $Q_0^+(D)$.
- Kawatani (2019): $\pi_1(Q_0^+(D)) \cong \star_{\text{free}} T_S^2$.
- Bayer–Bridgeland (2017): $\operatorname{Stab}_{\text{red}}^\dagger(D)/\mathbb{C}$ is contractible.
- Combining these two results, we have $\operatorname{Deck}(\pi) \cong \star_{\text{free}} T_S^2$.
- We showed that this implies the fixed point of Φ in $Q_0^+(D)$ can be lifted to a fixed point in $\operatorname{Stab}_{\text{red}}^\dagger(D)/\mathbb{C}$, which proves the realization problem for cyclic subgroups of $\operatorname{Aut}(D)/[1]$.
- Finally, we showed that any finite subgroup of $\operatorname{Aut}(D)/[1]$ is cyclic using classification results of finite subgroups of $\Gamma_0^+(n)$ by Fricke. This concludes the proof.

Lifting of fixed points

- Let $\Phi \in \operatorname{Aut}(D)/[1]$ be of finite order, then the previous discussion shows that it fixes a point in $Q_0^+(D)$.
- Kawatani (2019): $\pi_1(Q_0^+(D)) \cong \star_{\text{free}} T_S^2$.
- Bayer–Bridgeland (2017): $\operatorname{Stab}_{\text{red}}^\dagger(D)/\mathbb{C}$ is contractible.
- Combining these two results, we have $\operatorname{Deck}(\pi) \cong \star_{\text{free}} T_S^2$.
- We showed that this implies the fixed point of Φ in $Q_0^+(D)$ can be lifted to a fixed point in $\operatorname{Stab}_{\text{red}}^\dagger(D)/\mathbb{C}$, which proves the realization problem for cyclic subgroups of $\operatorname{Aut}(D)/[1]$.
- Finally, we showed that any finite subgroup of $\operatorname{Aut}(D)/[1]$ is cyclic using classification results of finite subgroups of $\Gamma_0^+(n)$ by Fricke. This concludes the proof.

Lifting of fixed points

- Let $\Phi \in \operatorname{Aut}(D)/[1]$ be of finite order, then the previous discussion shows that it fixes a point in $Q_0^+(D)$.
- Kawatani (2019): $\pi_1(Q_0^+(D)) \cong \star_{\text{free}} T_S^2$.
- Bayer–Bridgeland (2017): $\operatorname{Stab}_{\text{red}}^\dagger(D)/\mathbb{C}$ is contractible.
- Combining these two results, we have $\operatorname{Deck}(\pi) \cong \star_{\text{free}} T_S^2$.
- We showed that this implies the fixed point of Φ in $Q_0^+(D)$ can be lifted to a fixed point in $\operatorname{Stab}_{\text{red}}^\dagger(D)/\mathbb{C}$, which proves the realization problem for cyclic subgroups of $\operatorname{Aut}(D)/[1]$.
- Finally, we showed that any finite subgroup of $\operatorname{Aut}(D)/[1]$ is cyclic using classification results of finite subgroups of $\Gamma_0^+(n)$ by Fricke. This concludes the proof.

Lifting of fixed points

- Let $\Phi \in \operatorname{Aut}(D)/[1]$ be of finite order, then the previous discussion shows that it fixes a point in $Q_0^+(D)$.
- Kawatani (2019): $\pi_1(Q_0^+(D)) \cong \star_{\text{free}} T_S^2$.
- Bayer–Bridgeland (2017): $\operatorname{Stab}_{\text{red}}^\dagger(D)/\mathbb{C}$ is contractible.
- Combining these two results, we have $\operatorname{Deck}(\pi) \cong \star_{\text{free}} T_S^2$.
- We showed that this implies the fixed point of Φ in $Q_0^+(D)$ can be lifted to a fixed point in $\operatorname{Stab}_{\text{red}}^\dagger(D)/\mathbb{C}$, which proves the realization problem for cyclic subgroups of $\operatorname{Aut}(D)/[1]$.
- Finally, we showed that any finite subgroup of $\operatorname{Aut}(D)/[1]$ is cyclic using classification results of finite subgroups of $\Gamma_0^+(n)$ by Fricke. This concludes the proof.

A few further problems

- Find good representatives of 0-reducible autoequivalences.
- Do 0-reducible autoequivalences have zero entropy? ($h(\otimes \mathcal{O}(1)) = 0$)
- Generalize the realization results to:
 - ▶ general special cubic fourfolds $\mathrm{Ku}(Y)$
 - ▶ K3 surfaces of Picard number $\rho \geq 2$
 - ▶ ...?

Thank you for your attention!

Reference: F.-Lai, *Nielsen realization problem for Bridgeland stability conditions on generic K3 surfaces*, arXiv:2302.12663