

Dynamical aspects of categories

Yu-Wei Fan (Tsinghua U.)

Outline:

- ▶ Background and examples of $\text{Stab}(\Delta \text{ category})$
(relations with Teichmüller theory and Calabi–Yau geometry)
- ▶ Study $\text{Aut}(D)$ via its action on $\text{Stab}(D)$
(dynamical invariants, classifications)
- ▶ Quasi-convergent paths in $\text{Stab}(D)$ and SOD
(Daniel Halpern-Leistner: relating Stab , QH^* , birational geometry)

Stab and Teichmüller

- ▶ D : triangulated category (e.g. $D^b\mathrm{Coh}(X)$)

Stab and Teichmüller

- ▶ D : triangulated category (e.g. $D^b\mathrm{Coh}(X)$)
- ▶ $\sigma = (Z, P) \in \mathrm{Stab}(D)$ consists of:
 - ▶ $Z: \mathrm{Ob}(D) \rightarrow \mathbb{C}$, where $Z(B) = Z(A) + Z(C) \ \forall A \rightarrow B \rightarrow C \xrightarrow{+1}$
 - ▶ $\{P(\phi)\}_{\phi \in \mathbb{R}}$ semistable objects of phase ϕ

Stab and Teichmüller

- ▶ D : triangulated category (e.g. $D^b\text{Coh}(X)$)
- ▶ $\sigma = (Z, P) \in \text{Stab}(D)$ consists of:
 - ▶ $Z: \text{Ob}(D) \rightarrow \mathbb{C}$, where $Z(B) = Z(A) + Z(C) \forall A \rightarrow B \rightarrow C \xrightarrow{+1}$
 - ▶ $\{P(\phi)\}_{\phi \in \mathbb{R}}$ semistable objects of phase ϕ

Analogy between Teichmüller theory and stability data:

Riemann surface S	Triangulated category D
curve C	object E
$C_1 \cap C_2$	$\text{Hom}(E_1, E_2)$
metric g	stability σ
geodesic	semistable object
$C_1 \# \cdots \# C_n$	HN filtration $\text{gr}_i(E) = E_i$
length $\sum \ell_g(C_i)$	mass $\sum Z_\sigma(E_i) $
$\text{MCG}(S) \curvearrowright \text{Teich}(S)$	$\text{Aut}(D) \curvearrowright \text{Stab}(D)$
Dehn twist	spherical twist

Other things that admit categorical analogues: Topological entropy, pseudo-Anosov maps, systoles and systolic inequalities, $\text{SL}(2, \mathbb{R})$ -action, counting of saddle connections, etc.

Stab and mirror symmetry

- ▶ $\sigma \in \text{Stab}(D^b(X))$ is analogous to a Kähler/symplectic structure on X .
- ▶ It is conjectured that for a Calabi–Yau manifold X with Kähler class ω , there is a stability condition on $D^b(X)$ with central charge

$$Z_\omega(E) = - \int_X e^{-i\omega} \text{ch}(E) \hat{\Gamma}_X + \cdots .$$

Stab and mirror symmetry

- ▶ $\sigma \in \text{Stab}(D^b(X))$ is analogous to a Kähler/symplectic structure on X .
- ▶ It is conjectured that for a Calabi–Yau manifold X with Kähler class ω , there is a stability condition on $D^b(X)$ with central charge

$$Z_\omega(E) = - \int_X e^{-i\omega} \text{ch}(E) \hat{f}_X + \dots.$$

- ▶ $\sigma \in \text{Stab}(\text{Fuk}(\check{X}))$ is analogous to a complex structure on \check{X} .
- ▶ It is conjectured that for a Calabi–Yau manifold \check{X} with holomorphic volume form Ω , there is a stability condition on $\text{Fuk}(\check{X})$ with central charge

$$Z_\Omega(L) = \int_L \Omega,$$

and semistable objects given by special Lagrangians.

Stab and mirror symmetry

- ▶ $\sigma \in \text{Stab}(D^b(X))$ is analogous to a Kähler/symplectic structure on X .
- ▶ It is conjectured that for a Calabi–Yau manifold X with Kähler class ω , there is a stability condition on $D^b(X)$ with central charge

$$Z_\omega(E) = - \int_X e^{-i\omega} \text{ch}(E) \hat{\Gamma}_X + \dots.$$

- ▶ $\sigma \in \text{Stab}(\text{Fuk}(\check{X}))$ is analogous to a complex structure on \check{X} .
- ▶ It is conjectured that for a Calabi–Yau manifold \check{X} with holomorphic volume form Ω , there is a stability condition on $\text{Fuk}(\check{X})$ with central charge

$$Z_\Omega(L) = \int_L \Omega,$$

and semistable objects given by special Lagrangians.

- ▶ There are deep connections among $\text{Stab}(D^b(X))$, quantum cohomology of X , birational geometry of X , which we'll discuss later.

Examples of stability conditions

Example: Let X be an elliptic curve. Then any $\sigma \in \text{Stab}(D^b(X))$ is (up to a natural group action) equivalent to the slope stability:

- ▶ $Z(E) = -\deg(E) + i \cdot \text{rank}(E)$.
- ▶ E is σ -semistable if and only if it is slope semistable.

Examples of stability conditions

Example: Let X be an elliptic curve. Then any $\sigma \in \text{Stab}(D^b(X))$ is (up to a natural group action) equivalent to the slope stability:

- ▶ $Z(E) = -\deg(E) + i \cdot \text{rank}(E)$.
- ▶ E is σ -semistable if and only if it is slope semistable.

Its mirror \check{X} is also an elliptic curve. Then $\check{\sigma} \in \text{Stab}(\text{Fuk}(\check{X}))$ is given by:

- ▶ $Z(L) = \int_L dz$.
- ▶ L is $\check{\sigma}$ -semistable if and only if it is a special Lagrangian (i.e. straight line).

Examples of stability conditions

Example: Let X be an elliptic curve. Then any $\sigma \in \text{Stab}(D^b(X))$ is (up to a natural group action) equivalent to the slope stability:

- ▶ $Z(E) = -\deg(E) + i \cdot \text{rank}(E)$.
- ▶ E is σ -semistable if and only if it is slope semistable.

Its mirror \check{X} is also an elliptic curve. Then $\check{\sigma} \in \text{Stab}(\text{Fuk}(\check{X}))$ is given by:

- ▶ $Z(L) = \int_L dz$.
- ▶ L is $\check{\sigma}$ -semistable if and only if it is a special Lagrangian (i.e. straight line).

Example: $\text{Stab}(D^b(\mathbb{P}^1)) \cong \mathbb{C}^2$ has wall-crossings and consists of geometric and algebraic stability conditions. (We'll see a picture later.)

Examples of stability conditions

Example: Let X be an elliptic curve. Then any $\sigma \in \text{Stab}(D^b(X))$ is (up to a natural group action) equivalent to the slope stability:

- ▶ $Z(E) = -\deg(E) + i \cdot \text{rank}(E)$.
- ▶ E is σ -semistable if and only if it is slope semistable.

Its mirror \check{X} is also an elliptic curve. Then $\check{\sigma} \in \text{Stab}(\text{Fuk}(\check{X}))$ is given by:

- ▶ $Z(L) = \int_L dz$.
- ▶ L is $\check{\sigma}$ -semistable if and only if it is a special Lagrangian (i.e. straight line).

Example: $\text{Stab}(D^b(\mathbb{P}^1)) \cong \mathbb{C}^2$ has wall-crossings and consists of geometric and algebraic stability conditions. (We'll see a picture later.)

Its mirror is FS $\left(W: \mathbb{C}^* \xrightarrow{z+1/z} \mathbb{C}\right)$. By Haiden–Katzarkov–Kontsevich, its

stability space can be parametrized by 1-forms $\phi_{a,b} = \exp\left(z + a + \frac{b}{z}\right) \frac{dz}{z}$:

- ▶ $Z_{a,b} = \int_L \phi_{a,b}$.
- ▶ L is $\sigma_{a,b}$ -semistable if and only if it is a finite length sLag w.r.t. $\phi_{a,b}$.

Examples of Aut acting on Stab

There are natural actions $\mathrm{Aut}(D) \curvearrowright \mathrm{Stab}(D)$ and $\mathrm{Aut}(D)/[1] \curvearrowright \mathrm{Stab}(D)/\mathbb{C}$.

Examples of Aut acting on Stab

There are natural actions $\text{Aut}(D) \curvearrowright \text{Stab}(D)$ and $\text{Aut}(D)/[1] \curvearrowright \text{Stab}(D)/\mathbb{C}$.

Example: Let X be an elliptic curve. Then the action of $\text{Aut}(D)/[1]$ on $\text{Stab}(D)/\mathbb{C} \cong \mathbb{H}$ factors through $\text{PSL}(2, \mathbb{Z}) \curvearrowright \mathbb{H}$. In this case, we have

$$\mathcal{M}_{\text{K\"ah}}(X) \cong \mathbb{H}/\text{PSL}(2, \mathbb{Z}).$$



Examples of Aut acting on Stab

There are natural actions $\text{Aut}(D) \curvearrowright \text{Stab}(D)$ and $\text{Aut}(D)/[1] \curvearrowright \text{Stab}(D)/\mathbb{C}$.

Example: Let X be an elliptic curve. Then the action of $\text{Aut}(D)/[1]$ on $\text{Stab}(D)/\mathbb{C} \cong \mathbb{H}$ factors through $\text{PSL}(2, \mathbb{Z}) \curvearrowright \mathbb{H}$. In this case, we have

$$\mathcal{M}_{\text{K\"ah}}(X) \cong \mathbb{H}/\text{PSL}(2, \mathbb{Z}).$$

Example: The action of $\text{Aut}(D^b(\mathbb{P}^1))/[1]$ on $\text{Stab}(D^b(\mathbb{P}^1))/\mathbb{C} \cong \mathbb{C}$ factors through an \mathbb{Z} -action (given by $\mathcal{O}(1) \otimes -$):



Consider the orbit of $\Phi \in \text{Aut}(D)$ acting on $\sigma \in \text{Stab}(D)$

$$\{\dots, \Phi^{-1}\sigma, \sigma, \Phi\sigma, \dots\}.$$

We can use their mass functions $m_\sigma(\bullet)$ and phase functions $\phi_\sigma^\pm(\bullet)$ to extract dynamical invariants of Φ .

- ▶ The growth rate of $m_{\Phi^n\sigma}(\bullet)$ gives the categorical generalization of topological entropy.
- ▶ The growth rate of $\phi_{\Phi^n\sigma}^\pm(\bullet)$ gives the categorical generalization of Poincaré translation number.

Consider the orbit of $\Phi \in \text{Aut}(D)$ acting on $\sigma \in \text{Stab}(D)$

$$\{\dots, \Phi^{-1}\sigma, \sigma, \Phi\sigma, \dots\}.$$

We can use their mass functions $m_\sigma(\bullet)$ and phase functions $\phi_\sigma^\pm(\bullet)$ to extract dynamical invariants of Φ .

- ▶ The growth rate of $m_{\Phi^n\sigma}(\bullet)$ gives the categorical generalization of topological entropy.
- ▶ The growth rate of $\phi_{\Phi^n\sigma}^\pm(\bullet)$ gives the categorical generalization of Poincaré translation number.

Using the analogy between $\text{MCG}(S) \curvearrowright \text{Teich}(S)$ and $\text{Aut}(D) \curvearrowright \text{Stab}(D)$, we can also study:

- ▶ trichotomy like finite order/reducible/pseudo-Anosov;
- ▶ properties of finite order autoequivalences;
- ▶ various categorical generalizations of pseudo-Anosov maps.

Topological entropy

Let (X, d) be a compact metric space and $f: X \rightarrow X$ continuous. Consider

$$N(n, \epsilon) := \max \left\{ \ell: \exists x_1, \dots, x_\ell \text{ s.t. } \max_{0 \leq k \leq n} \{d(f^k(x_i), f^k(x_j))\} \geq \epsilon \forall x_i, x_j \right\}$$

Topological entropy

Let (X, d) be a compact metric space and $f: X \rightarrow X$ continuous. Consider

$$N(n, \epsilon) := \max \left\{ \ell: \exists x_1, \dots, x_\ell \text{ s.t. } \max_{0 \leq k \leq n} \{d(f^k(x_i), f^k(x_j))\} \geq \epsilon \forall x_i, x_j \right\}$$

The topological entropy of f is defined to be

$$h_{\text{top}}(f) := \lim_{\epsilon \rightarrow 0} \left(\limsup_{n \rightarrow \infty} \frac{1}{n} \log N(n, \epsilon) \right) \in [0, \infty].$$

Topological entropy

Let (X, d) be a compact metric space and $f: X \rightarrow X$ continuous. Consider

$$N(n, \epsilon) := \max \left\{ \ell: \exists x_1, \dots, x_\ell \text{ s.t. } \max_{0 \leq k \leq n} \{d(f^k(x_i), f^k(x_j))\} \geq \epsilon \forall x_i, x_j \right\}$$

The topological entropy of f is defined to be

$$h_{\text{top}}(f) := \lim_{\epsilon \rightarrow 0} \left(\limsup_{n \rightarrow \infty} \frac{1}{n} \log N(n, \epsilon) \right) \in [0, \infty].$$

- It's a topological invariant measuring the “complexity” of f .

Topological entropy

Let (X, d) be a compact metric space and $f: X \rightarrow X$ continuous. Consider

$$N(n, \epsilon) := \max \left\{ \ell: \exists x_1, \dots, x_\ell \text{ s.t. } \max_{0 \leq k \leq n} \{d(f^k(x_i), f^k(x_j))\} \geq \epsilon \forall x_i, x_j \right\}$$

The topological entropy of f is defined to be

$$h_{\text{top}}(f) := \lim_{\epsilon \rightarrow 0} \left(\limsup_{n \rightarrow \infty} \frac{1}{n} \log N(n, \epsilon) \right) \in [0, \infty].$$

- ▶ It's a topological invariant measuring the “complexity” of f .
- ▶ $f^n = \text{id}_X \implies h_{\text{top}}(f) = 0$.

Topological entropy

Let (X, d) be a compact metric space and $f: X \rightarrow X$ continuous. Consider

$$N(n, \epsilon) := \max \left\{ \ell: \exists x_1, \dots, x_\ell \text{ s.t. } \max_{0 \leq k \leq n} \{d(f^k(x_i), f^k(x_j))\} \geq \epsilon \forall x_i, x_j \right\}$$

The topological entropy of f is defined to be

$$h_{\text{top}}(f) := \lim_{\epsilon \rightarrow 0} \left(\limsup_{n \rightarrow \infty} \frac{1}{n} \log N(n, \epsilon) \right) \in [0, \infty].$$

- ▶ It's a topological invariant measuring the “complexity” of f .
- ▶ $f^n = \text{id}_X \implies h_{\text{top}}(f) = 0$.
- ▶ (Yomdin) If X is a compact differentiable manifold and f is smooth, then $h_{\text{top}}(f) \geq \log \rho(f_{H^*(X, \mathbb{C})}^*)$.

Topological entropy

Let (X, d) be a compact metric space and $f: X \rightarrow X$ continuous. Consider

$$N(n, \epsilon) := \max \left\{ \ell: \exists x_1, \dots, x_\ell \text{ s.t. } \max_{0 \leq k \leq n} \{d(f^k(x_i), f^k(x_j))\} \geq \epsilon \forall x_i, x_j \right\}$$

The topological entropy of f is defined to be

$$h_{\text{top}}(f) := \lim_{\epsilon \rightarrow 0} \left(\limsup_{n \rightarrow \infty} \frac{1}{n} \log N(n, \epsilon) \right) \in [0, \infty].$$

- ▶ It's a topological invariant measuring the “complexity” of f .
- ▶ $f^n = \text{id}_X \implies h_{\text{top}}(f) = 0$.
- ▶ (Yomdin) If X is a compact differentiable manifold and f is smooth, then $h_{\text{top}}(f) \geq \log \rho(f_{H^*(X, \mathbb{C})}^*)$.
- ▶ (Gromov) Moreover, if X is Kähler and f is holomorphic, then $h_{\text{top}}(f) = \log \rho(f_{H^*(X, \mathbb{C})}^*)$.

Topological entropy

Let (X, d) be a compact metric space and $f: X \rightarrow X$ continuous. Consider

$$N(n, \epsilon) := \max \left\{ \ell: \exists x_1, \dots, x_\ell \text{ s.t. } \max_{0 \leq k \leq n} \{d(f^k(x_i), f^k(x_j))\} \geq \epsilon \forall x_i, x_j \right\}$$

The topological entropy of f is defined to be

$$h_{\text{top}}(f) := \lim_{\epsilon \rightarrow 0} \left(\limsup_{n \rightarrow \infty} \frac{1}{n} \log N(n, \epsilon) \right) \in [0, \infty].$$

- ▶ It's a topological invariant measuring the “complexity” of f .
- ▶ $f^n = \text{id}_X \implies h_{\text{top}}(f) = 0$.
- ▶ (Yomdin) If X is a compact differentiable manifold and f is smooth, then $h_{\text{top}}(f) \geq \log \rho(f_{H^*(X, \mathbb{C})}^*)$.
- ▶ (Gromov) Moreover, if X is Kähler and f is holomorphic, then $h_{\text{top}}(f) = \log \rho(f_{H^*(X, \mathbb{C})}^*)$.
- ▶ (Cantat) If a compact complex surface X admits $h_{\text{top}}(f) > 0$, then X is either a torus, a K3 surface, an Enriques surface, or a rational surface.

Categorical entropy

Let $\Phi \in \text{Aut}(D)$, $\sigma \in \text{Stab}(D)$, and $G \in D$ a split generator.

$$h_{\sigma}(\Phi) := \limsup_{n \rightarrow \infty} \frac{1}{n} \log m_{\sigma}(\Phi^n G).$$

Categorical entropy

Let $\Phi \in \text{Aut}(D)$, $\sigma \in \text{Stab}(D)$, and $G \in D$ a split generator.

$$h_\sigma(\Phi) := \limsup_{n \rightarrow \infty} \frac{1}{n} \log m_\sigma(\Phi^n G).$$

$h_\sigma(\Phi)$: Not always computable; not known to be independent of $\pi_0(\text{Stab}(D))$.

Categorical entropy

Let $\Phi \in \text{Aut}(D)$, $\sigma \in \text{Stab}(D)$, and $G \in D$ a split generator.

$$h_{\sigma}(\Phi) := \limsup_{n \rightarrow \infty} \frac{1}{n} \log m_{\sigma}(\Phi^n G).$$

$h_{\sigma}(\Phi)$: Not always computable; not known to be independent of $\pi_0(\text{Stab}(D))$.

$$h_{\text{cat}}(\Phi) := \lim_{n \rightarrow \infty} \frac{1}{n} \log \left(\sum_k \dim \text{Hom}(G, \Phi^n G[k]) \right).$$

Categorical entropy

Let $\Phi \in \text{Aut}(D)$, $\sigma \in \text{Stab}(D)$, and $G \in D$ a split generator.

$$h_\sigma(\Phi) := \limsup_{n \rightarrow \infty} \frac{1}{n} \log m_\sigma(\Phi^n G).$$

$h_\sigma(\Phi)$: Not always computable; not known to be independent of $\pi_0(\text{Stab}(D))$.

$$h_{\text{cat}}(\Phi) := \lim_{n \rightarrow \infty} \frac{1}{n} \log \left(\sum_k \dim \text{Hom}(G, \Phi^n G[k]) \right).$$

► (Ikeda) $h_{\text{cat}}(\Phi) \geq h_\sigma(\Phi) \geq \log \rho([\Phi])$ for all $\sigma \in \text{Stab}(D)$.

Categorical entropy

Let $\Phi \in \operatorname{Aut}(D)$, $\sigma \in \operatorname{Stab}(D)$, and $G \in D$ a split generator.

$$h_{\sigma}(\Phi) := \limsup_{n \rightarrow \infty} \frac{1}{n} \log m_{\sigma}(\Phi^n G).$$

$h_{\sigma}(\Phi)$: Not always computable; not known to be independent of $\pi_0(\operatorname{Stab}(D))$.

$$h_{\text{cat}}(\Phi) := \lim_{n \rightarrow \infty} \frac{1}{n} \log \left(\sum_k \dim \operatorname{Hom}(G, \Phi^n G[k]) \right).$$

- ▶ (Ikeda) $h_{\text{cat}}(\Phi) \geq h_{\sigma}(\Phi) \geq \log \rho([\Phi])$ for all $\sigma \in \operatorname{Stab}(D)$.
- ▶ (Kikuta–Takahashi) For $f \in \operatorname{Aut}(X)$, $h_{\text{cat}}(\mathbb{L}f^*) = h_{\text{top}}(f) = \log \rho([\mathbb{L}f^*])$.

Categorical entropy

Let $\Phi \in \text{Aut}(D)$, $\sigma \in \text{Stab}(D)$, and $G \in D$ a split generator.

$$h_\sigma(\Phi) := \limsup_{n \rightarrow \infty} \frac{1}{n} \log m_\sigma(\Phi^n G).$$

$h_\sigma(\Phi)$: Not always computable; not known to be independent of $\pi_0(\text{Stab}(D))$.

$$h_{\text{cat}}(\Phi) := \lim_{n \rightarrow \infty} \frac{1}{n} \log \left(\sum_k \dim \text{Hom}(G, \Phi^n G[k]) \right).$$

- ▶ (Ikeda) $h_{\text{cat}}(\Phi) \geq h_\sigma(\Phi) \geq \log \rho([\Phi])$ for all $\sigma \in \text{Stab}(D)$.
- ▶ (Kikuta–Takahashi) For $f \in \text{Aut}(X)$, $h_{\text{cat}}(\mathbb{L}f^*) = h_{\text{top}}(f) = \log \rho([\mathbb{L}f^*])$.
- ▶ (F.) $\exists \Phi \in \text{Aut}(D^b(X))$ with $h_{\text{cat}}(\Phi) > \log \rho([\Phi])$ for some CY X .

Categorical entropy

Let $\Phi \in \text{Aut}(D)$, $\sigma \in \text{Stab}(D)$, and $G \in D$ a split generator.

$$h_\sigma(\Phi) := \limsup_{n \rightarrow \infty} \frac{1}{n} \log m_\sigma(\Phi^n G).$$

$h_\sigma(\Phi)$: Not always computable; not known to be independent of $\pi_0(\text{Stab}(D))$.

$$h_{\text{cat}}(\Phi) := \lim_{n \rightarrow \infty} \frac{1}{n} \log \left(\sum_k \dim \text{Hom}(G, \Phi^n G[k]) \right).$$

- ▶ (Ikeda) $h_{\text{cat}}(\Phi) \geq h_\sigma(\Phi) \geq \log \rho([\Phi])$ for all $\sigma \in \text{Stab}(D)$.
- ▶ (Kikuta–Takahashi) For $f \in \text{Aut}(X)$, $h_{\text{cat}}(\mathbb{L}f^*) = h_{\text{top}}(f) = \log \rho([\mathbb{L}f^*])$.
- ▶ (F.) $\exists \Phi \in \text{Aut}(D^b(X))$ with $h_{\text{cat}}(\Phi) > \log \rho([\Phi])$ for some CY X .

Remark: Existence of manifold with corner structure of the compactification of Stab in D. H.-L.'s program $\implies \text{hom}(G, E) \leq C \cdot m_\sigma(E)$ for all $E \implies$

- ▶ $h_{\text{cat}}(\Phi) = h_\sigma(\Phi)$.
- ▶ $\mathcal{M}_\sigma^{\text{ss}}(\nu)$ admits a proper good moduli space.

Categorical polynomial entropy

For $h_{\text{cat}}(\Phi) = 0$, one can consider a more refined invariant, its polynomial entropy

$$h_{\text{poly}}(\Phi) := \limsup_{n \rightarrow \infty} \frac{\log(\sum_k \dim \text{Hom}(G, \Phi^n G[k]))}{\log n}.$$

Categorical polynomial entropy

For $h_{\text{cat}}(\Phi) = 0$, one can consider a more refined invariant, its polynomial entropy

$$h_{\text{poly}}(\Phi) := \limsup_{n \rightarrow \infty} \frac{\log(\sum_k \dim \text{Hom}(G, \Phi^n G[k]))}{\log n}.$$

F.–Fu–Ouchi:

- Yomdin-type lower bound still holds.

Categorical polynomial entropy

For $h_{\text{cat}}(\Phi) = 0$, one can consider a more refined invariant, its polynomial entropy

$$h_{\text{poly}}(\Phi) := \limsup_{n \rightarrow \infty} \frac{\log(\sum_k \dim \text{Hom}(G, \Phi^n G[k]))}{\log n}.$$

F.-Fu-Ouchi:

- ▶ Yomdin-type lower bound still holds.
- ▶ Let L be a line bundle on X . Then $h_{\text{cat}}(- \otimes L) = 0$, and

$$\nu(L) \leq h_{\text{poly}}(- \otimes L) \leq \dim(X)$$

where $\nu(L) = \max\{m \mid c_1(L)^m \neq 0\}$ the numerical dimension of L .

Categorical polynomial entropy

For $h_{\text{cat}}(\Phi) = 0$, one can consider a more refined invariant, its polynomial entropy

$$h_{\text{poly}}(\Phi) := \limsup_{n \rightarrow \infty} \frac{\log(\sum_k \dim \text{Hom}(G, \Phi^n G[k]))}{\log n}.$$

F.-Fu-Ouchi:

- ▶ Yomdin-type lower bound still holds.
- ▶ Let L be a line bundle on X . Then $h_{\text{cat}}(- \otimes L) = 0$, and

$$\nu(L) \leq h_{\text{poly}}(- \otimes L) \leq \dim(X)$$

where $\nu(L) = \max\{m \mid c_1(L)^m \neq 0\}$ the numerical dimension of L .

- ▶ Let X be an elliptic curve, and $\Phi \in \text{Aut}(D^b(X))$. Then
 - ▶ $h_{\text{cat}}(\Phi) = h_{\text{poly}}(\Phi) = 0$ iff $[\Phi] \in \text{SL}(2, \mathbb{Z})$ is elliptic.
 - ▶ $h_{\text{cat}}(\Phi) = 0$ and $h_{\text{poly}}(\Phi) = 1$ iff $[\Phi] \in \text{SL}(2, \mathbb{Z})$ is parabolic.
 - ▶ $h_{\text{cat}}(\Phi) > 0$ iff $[\Phi] \in \text{SL}(2, \mathbb{Z})$ is hyperbolic.

Shifting numbers

(F.-Filip) The limits $\tau^\pm(\Phi) := \lim_{n \rightarrow \infty} \frac{1}{n} \phi_\sigma^\pm(\Phi^n G)$ always exist, and are independent of the choices of G and σ .

Shifting numbers

(F.–Filip) The limits $\tau^\pm(\Phi) := \lim_{n \rightarrow \infty} \frac{1}{n} \phi_\sigma^\pm(\Phi^n G)$ always exist, and are independent of the choices of G and σ .

Poincaré translation numbers ρ	Shifting numbers
$\mathbb{Z} \hookrightarrow \text{Homeo}_{\mathbb{Z}}^+(\mathbb{R}) \twoheadrightarrow \text{Homeo}^+(S^1)$	$\mathbb{Z} \hookrightarrow \text{Aut}(\mathcal{D}) \twoheadrightarrow \text{Aut}(\mathcal{D})/[1]$
$f \in \text{Homeo}_{\mathbb{Z}}^+(\mathbb{R})$	$\Phi \in \text{Aut}(\mathcal{D})$
$x_0 \in \mathbb{R}$	$G \in \mathcal{D}$
amount of translation	phases $\phi_\sigma^\pm: \text{Ob}(\mathcal{D}) \rightarrow \mathbb{R}$
$f^{(n)}(x_0) - x_0$	$\phi_\sigma^\pm(\Phi^n G) - \phi_\sigma^\pm(G)$
translation number	upper/lower shifting numbers

Shifting numbers

(F.–Filip) The limits $\tau^\pm(\Phi) := \lim_{n \rightarrow \infty} \frac{1}{n} \phi_\sigma^\pm(\Phi^n G)$ always exist, and are independent of the choices of G and σ .

Poincaré translation numbers ρ	Shifting numbers
$\mathbb{Z} \hookrightarrow \text{Homeo}_{\mathbb{Z}}^+(\mathbb{R}) \twoheadrightarrow \text{Homeo}^+(S^1)$	$\mathbb{Z} \hookrightarrow \text{Aut}(\mathcal{D}) \twoheadrightarrow \text{Aut}(\mathcal{D})/[1]$
$f \in \text{Homeo}_{\mathbb{Z}}^+(\mathbb{R})$	$\Phi \in \text{Aut}(\mathcal{D})$
$x_0 \in \mathbb{R}$	$G \in \mathcal{D}$
amount of translation	phases $\phi_\sigma^\pm: \text{Ob}(\mathcal{D}) \rightarrow \mathbb{R}$
$f^{(n)}(x_0) - x_0$	$\phi_\sigma^\pm(\Phi^n G) - \phi_\sigma^\pm(G)$
translation number	upper/lower shifting numbers

Note: $\tau^\pm(\text{Serre}_D)$ is the upper/lower Serre dimension of D .

Shifting numbers

(F.–Filip) The limits $\tau^\pm(\Phi) := \lim_{n \rightarrow \infty} \frac{1}{n} \phi_\sigma^\pm(\Phi^n G)$ always exist, and are independent of the choices of G and σ .

Poincaré translation numbers ρ	Shifting numbers
$\mathbb{Z} \hookrightarrow \text{Homeo}_{\mathbb{Z}}^+(\mathbb{R}) \twoheadrightarrow \text{Homeo}^+(S^1)$	$\mathbb{Z} \hookrightarrow \text{Aut}(\mathcal{D}) \twoheadrightarrow \text{Aut}(\mathcal{D})/[1]$
$f \in \text{Homeo}_{\mathbb{Z}}^+(\mathbb{R})$	$\Phi \in \text{Aut}(\mathcal{D})$
$x_0 \in \mathbb{R}$	$G \in \mathcal{D}$
amount of translation	phases $\phi_\sigma^\pm: \text{Ob}(\mathcal{D}) \rightarrow \mathbb{R}$
$f^{(n)}(x_0) - x_0$	$\phi_\sigma^\pm(\Phi^n G) - \phi_\sigma^\pm(G)$
translation number	upper/lower shifting numbers

Note: $\tau^\pm(\text{Serre}_D)$ is the upper/lower Serre dimension of D .

► When X is an elliptic curve, $\tau = \tau^\pm$ can be decomposed as

$$\text{Aut}(D^b(X)) \rightarrow \text{Homeo}_{\mathbb{Z}}^+(\mathbb{R}) \xrightarrow{\rho} \mathbb{R}.$$

Shifting numbers

(F.–Filip) The limits $\tau^\pm(\Phi) := \lim_{n \rightarrow \infty} \frac{1}{n} \phi_\sigma^\pm(\Phi^n G)$ always exist, and are independent of the choices of G and σ .

Poincaré translation numbers ρ	Shifting numbers
$\mathbb{Z} \hookrightarrow \text{Homeo}_{\mathbb{Z}}^+(\mathbb{R}) \twoheadrightarrow \text{Homeo}^+(S^1)$	$\mathbb{Z} \hookrightarrow \text{Aut}(\mathcal{D}) \twoheadrightarrow \text{Aut}(\mathcal{D})/[1]$
$f \in \text{Homeo}_{\mathbb{Z}}^+(\mathbb{R})$	$\Phi \in \text{Aut}(\mathcal{D})$
$x_0 \in \mathbb{R}$	$G \in \mathcal{D}$
amount of translation	phases $\phi_\sigma^\pm: \text{Ob}(\mathcal{D}) \rightarrow \mathbb{R}$
$f^{(n)}(x_0) - x_0$	$\phi_\sigma^\pm(\Phi^n G) - \phi_\sigma^\pm(G)$
translation number	upper/lower shifting numbers

Note: $\tau^\pm(\text{Serre}_D)$ is the upper/lower Serre dimension of D .

- ▶ When X is an elliptic curve, $\tau = \tau^\pm$ can be decomposed as

$$\text{Aut}(D^b(X)) \rightarrow \text{Homeo}_{\mathbb{Z}}^+(\mathbb{R}) \xrightarrow{\rho} \mathbb{R}.$$

- ▶ When X is an abelian surface, τ coincides with a standard quasimorphism on certain Lie group of Hermitian type.

Shifting numbers

(F.–Filip) The limits $\tau^\pm(\Phi) := \lim_{n \rightarrow \infty} \frac{1}{n} \phi_\sigma^\pm(\Phi^n G)$ always exist, and are independent of the choices of G and σ .

Poincaré translation numbers ρ	Shifting numbers
$\mathbb{Z} \hookrightarrow \text{Homeo}_{\mathbb{Z}}^+(\mathbb{R}) \twoheadrightarrow \text{Homeo}^+(S^1)$	$\mathbb{Z} \hookrightarrow \text{Aut}(\mathcal{D}) \twoheadrightarrow \text{Aut}(\mathcal{D})/[1]$
$f \in \text{Homeo}_{\mathbb{Z}}^+(\mathbb{R})$	$\Phi \in \text{Aut}(\mathcal{D})$
$x_0 \in \mathbb{R}$	$G \in \mathcal{D}$
amount of translation	phases $\phi_\sigma^\pm: \text{Ob}(\mathcal{D}) \rightarrow \mathbb{R}$
$f^{(n)}(x_0) - x_0$	$\phi_\sigma^\pm(\Phi^n G) - \phi_\sigma^\pm(G)$
translation number	upper/lower shifting numbers

Note: $\tau^\pm(\text{Serre}_D)$ is the upper/lower Serre dimension of D .

- ▶ When X is an elliptic curve, $\tau = \tau^\pm$ can be decomposed as

$$\text{Aut}(D^b(X)) \rightarrow \text{Homeo}_{\mathbb{Z}}^+(\mathbb{R}) \xrightarrow{\rho} \mathbb{R}.$$

- ▶ When X is an abelian surface, τ coincides with a standard quasimorphism on certain Lie group of Hermitian type.
- ▶ (F.) When X is an abelian variety $\tau = \tau^\pm: \text{Aut}(D^b(X)) \rightarrow \mathbb{R}$ is a quasimorphism.

Finite order autoequivalences

Finite order autoequivalences

- ▶ Nielsen asked: Let $G \subseteq \text{MCG}(S) = \text{Diff}(S)/\{\text{isotopy}\}$ be a finite subgroup. Does there always exist a lifting $G \subseteq \text{Diff}(S)$?

Finite order autoequivalences

- ▶ Nielsen asked: Let $G \subseteq \text{MCG}(S) = \text{Diff}(S)/\{\text{isotopy}\}$ be a finite subgroup. Does there always exist a lifting $G \subseteq \text{Diff}(S)$?
- ▶ Kerckhoff: Yes! Moreover, there exists a metric g such that $G \subseteq \text{Isom}(S, g)$. Or equivalently, G fixes a point in $\text{Teich}(S)$.

Finite order autoequivalences

- ▶ Nielsen asked: Let $G \subseteq \text{MCG}(S) = \text{Diff}(S)/\{\text{isotopy}\}$ be a finite subgroup. Does there always exist a lifting $G \subseteq \text{Diff}(S)$?
- ▶ Kerckhoff: Yes! Moreover, there exists a metric g such that $G \subseteq \text{Isom}(S, g)$. Or equivalently, G fixes a point in $\text{Teich}(S)$.
- ▶ Farb–Looijenga also proved similar statements for K3 surfaces (under certain conditions), where g is replaced by complex structure or Ricci-flat metric.

Finite order autoequivalences

- ▶ Nielsen asked: Let $G \subseteq \text{MCG}(S) = \text{Diff}(S)/\{\text{isotopy}\}$ be a finite subgroup. Does there always exist a lifting $G \subseteq \text{Diff}(S)$?
- ▶ Kerckhoff: Yes! Moreover, there exists a metric g such that $G \subseteq \text{Isom}(S, g)$. Or equivalently, G fixes a point in $\text{Teich}(S)$.
- ▶ Farb–Looijenga also proved similar statements for K3 surfaces (under certain conditions), where g is replaced by complex structure or Ricci-flat metric.

Categorical realization problem: Let $G \subseteq \text{Aut}(D)/[1]$ be a finite subgroup. Does there exist $\sigma \in \text{Stab}(D)/\mathbb{C}$ such that $\Phi \cdot \sigma = \sigma$ for all $\Phi \in G$?

Finite order autoequivalences

- ▶ Nielsen asked: Let $G \subseteq \text{MCG}(S) = \text{Diff}(S)/\{\text{isotopy}\}$ be a finite subgroup. Does there always exist a lifting $G \subseteq \text{Diff}(S)$?
- ▶ Kerckhoff: Yes! Moreover, there exists a metric g such that $G \subseteq \text{Isom}(S, g)$. Or equivalently, G fixes a point in $\text{Teich}(S)$.
- ▶ Farb–Looijenga also proved similar statements for K3 surfaces (under certain conditions), where g is replaced by complex structure or Ricci-flat metric.

Categorical realization problem: Let $G \subseteq \text{Aut}(D)/[1]$ be a finite subgroup. Does there exist $\sigma \in \text{Stab}(D)/\mathbb{C}$ such that $\Phi \cdot \sigma = \sigma$ for all $\Phi \in G$?

- ▶ When $D = D^b(X)$, stability conditions on D are roughly Kähler structures, so this is similar to (but not exactly) the mirror problem of Farb–Looijenga.

Finite order autoequivalences

- ▶ Nielsen asked: Let $G \subseteq \text{MCG}(S) = \text{Diff}(S)/\{\text{isotopy}\}$ be a finite subgroup. Does there always exist a lifting $G \subseteq \text{Diff}(S)$?
- ▶ Kerckhoff: Yes! Moreover, there exists a metric g such that $G \subseteq \text{Isom}(S, g)$. Or equivalently, G fixes a point in $\text{Teich}(S)$.
- ▶ Farb–Looijenga also proved similar statements for K3 surfaces (under certain conditions), where g is replaced by complex structure or Ricci-flat metric.

Categorical realization problem: Let $G \subseteq \text{Aut}(D)/[1]$ be a finite subgroup. Does there exist $\sigma \in \text{Stab}(D)/\mathbb{C}$ such that $\Phi \cdot \sigma = \sigma$ for all $\Phi \in G$?

- ▶ When $D = D^b(X)$, stability conditions on D are roughly Kähler structures, so this is similar to (but not exactly) the mirror problem of Farb–Looijenga.
- ▶ (F.–Lai) Yes, for $D = D^b(X)$ where X is a curve, an abelian surface, a generic twisted K3 surface, or a K3 surface of Picard number $\rho = 1$.

Finite order autoequivalences

- ▶ Nielsen asked: Let $G \subseteq \mathrm{MCG}(S) = \mathrm{Diff}(S)/\{\text{isotopy}\}$ be a finite subgroup. Does there always exist a lifting $G \subseteq \mathrm{Diff}(S)$?
- ▶ Kerckhoff: Yes! Moreover, there exists a metric g such that $G \subseteq \mathrm{Isom}(S, g)$. Or equivalently, G fixes a point in $\mathrm{Teich}(S)$.
- ▶ Farb–Looijenga also proved similar statements for K3 surfaces (under certain conditions), where g is replaced by complex structure or Ricci-flat metric.

Categorical realization problem: Let $G \subseteq \mathrm{Aut}(D)/[1]$ be a finite subgroup. Does there exist $\sigma \in \mathrm{Stab}(D)/\mathbb{C}$ such that $\Phi \cdot \sigma = \sigma$ for all $\Phi \in G$?

- ▶ When $D = D^b(X)$, stability conditions on D are roughly Kähler structures, so this is similar to (but not exactly) the mirror problem of Farb–Looijenga.
- ▶ (F.–Lai) Yes, for $D = D^b(X)$ where X is a curve, an abelian surface, a generic twisted K3 surface, or a K3 surface of Picard number $\rho = 1$.
- ▶ Moreover, this is the main theorem that we use to fully classify finite subgroups of $\mathrm{Aut}(D)/[1]$ for K3 surfaces of $\rho = 1$.

Finite order autoequivalences

- ▶ Nielsen asked: Let $G \subseteq \text{MCG}(S) = \text{Diff}(S)/\{\text{isotopy}\}$ be a finite subgroup. Does there always exist a lifting $G \subseteq \text{Diff}(S)$?
- ▶ Kerckhoff: Yes! Moreover, there exists a metric g such that $G \subseteq \text{Isom}(S, g)$. Or equivalently, G fixes a point in $\text{Teich}(S)$.
- ▶ Farb–Looijenga also proved similar statements for K3 surfaces (under certain conditions), where g is replaced by complex structure or Ricci-flat metric.

Categorical realization problem: Let $G \subseteq \text{Aut}(D)/[1]$ be a finite subgroup. Does there exist $\sigma \in \text{Stab}(D)/\mathbb{C}$ such that $\Phi \cdot \sigma = \sigma$ for all $\Phi \in G$?

- ▶ When $D = D^b(X)$, stability conditions on D are roughly Kähler structures, so this is similar to (but not exactly) the mirror problem of Farb–Looijenga.
- ▶ (F.–Lai) Yes, for $D = D^b(X)$ where X is a curve, an abelian surface, a generic twisted K3 surface, or a K3 surface of Picard number $\rho = 1$.
- ▶ Moreover, this is the main theorem that we use to fully classify finite subgroups of $\text{Aut}(D)/[1]$ for K3 surfaces of $\rho = 1$.
- ▶ Corollary: $\text{Aut}(D^b(X))/[1]$ contains an order 3 element if and only if X admits an associated cubic fourfold.

X : K3 surface, $g=1$, $H^2=2n$,

$$\begin{aligned} & \mathbb{H} / \Gamma_0^+(n) \\ & \parallel \\ & \langle \Gamma_0(n), \begin{bmatrix} 0 & \sqrt{n} \\ \sqrt{n} & 0 \end{bmatrix} \rangle \end{aligned}$$

$$\begin{aligned} & \Gamma_0^+(n) \backslash \Gamma_0(n) \\ & \Downarrow \\ & \text{Stab} = \langle g \rangle \\ & g^2 = 1. \end{aligned}$$

- Spherical twist,
- ∞ order autoeq.
- $h_{\text{cat}}=0$, $h_{\text{poly}}=1$

- $\text{Stab} = \langle g \rangle$, $g \in \Gamma_0(n)$
 $\text{ord}(g) = 2 \text{ or } 3$. \rightarrow finite order autoeq.
- $h_{\text{cat}} = h_{\text{poly}} = 0$.

cusps $\longleftrightarrow \otimes \mathbb{Q}_X(1)$ for some $D(X) = D(X')$

- infinite order autoeq.
- $h_{\text{cat}}=0$, $h_{\text{poly}}=2$.

Daniel Halpern-Leistner's NMMP proposal




Daniel Halpern-Leistner's NMMP proposal



- ▶ Every quasi-convergent paths in Stab (in the above example, satisfies $\phi_{\sigma_t}(\mathcal{O}) - \phi_{\sigma_t}(\mathcal{O}(-1)) \rightarrow +\infty$) give rise to a corresponding semiorthogonal decomposition (SOD), with stability on each component.

Daniel Halpern-Leistner's NMMP proposal




- ▶ Every quasi-convergent paths in Stab (in the above example, satisfies $\phi_{\sigma_t}(\mathcal{O}) - \phi_{\sigma_t}(\mathcal{O}(-1)) \rightarrow +\infty$) give rise to a corresponding semiorthogonal decomposition (SOD), with stability on each component.
- ▶ Every SOD (s.t. each component admits stability) can be realized by quasi-convergent paths.

Daniel Halpern-Leistner's NMMP proposal




- ▶ Every quasi-convergent paths in Stab (in the above example, satisfies $\phi_{\sigma_t}(\mathcal{O}) - \phi_{\sigma_t}(\mathcal{O}(-1)) \rightarrow +\infty$) give rise to a corresponding semiorthogonal decomposition (SOD), with stability on each component.
- ▶ Every SOD (s.t. each component admits stability) can be realized by quasi-convergent paths.
- ▶ Solution of the quantum differential equation gives a quasi-convergent path, therefore a canonical SOD of $D^b(X)$.

Daniel Halpern-Leistner's NMMP proposal



- ▶ Every quasi-convergent paths in Stab (in the above example, satisfies $\phi_{\sigma_t}(\mathcal{O}) - \phi_{\sigma_t}(\mathcal{O}(-1)) \rightarrow +\infty$) give rise to a corresponding semiorthogonal decomposition (SOD), with stability on each component.
- ▶ Every SOD (s.t. each component admits stability) can be realized by quasi-convergent paths.
- ▶ Solution of the quantum differential equation gives a quasi-convergent path, therefore a canonical SOD of $D^b(X)$.
- ▶ For every $X' \rightarrow X$ birational morphism of smooth projective varieties, the canonical SOD of X' refines the canonical SOD of X .

Daniel Halpern-Leistner's NMMP proposal



- ▶ Every quasi-convergent paths in Stab (in the above example, satisfies $\phi_{\sigma_t}(\mathcal{O}) - \phi_{\sigma_t}(\mathcal{O}(-1)) \rightarrow +\infty$) give rise to a corresponding semiorthogonal decomposition (SOD), with stability on each component.
- ▶ Every SOD (s.t. each component admits stability) can be realized by quasi-convergent paths.
- ▶ Solution of the quantum differential equation gives a quasi-convergent path, therefore a canonical SOD of $D^b(X)$.
- ▶ For every $X' \rightarrow X$ birational morphism of smooth projective varieties, the canonical SOD of X' refines the canonical SOD of X .
- ▶ Corollary: Existence of noncommutative minimal model, D-equivalence conjecture, (one side of) Dubrovin conjecture.

Thank you for your attention!