

# Nielsen realization for Bridgeland stability conditions on K3 surfaces

Yu-Wei Fan (Tsinghua)  
Joint work with Kuan-Wen Lai (Bonn)

May 2023, Nanjing

(arXiv:2302.12663)

## A general problem

Let  $\mathcal{D}$  be a triangulated category (e.g.  $D^b\mathrm{Coh}(X)$ ). Study  
the group of autoequivalences  $\mathrm{Aut}(\mathcal{D})$ .

(Why? It contains  $\mathrm{Aut}(X)$ , and “hidden symmetries” like spherical twists)

- Complexity: categorical entropy, categorical polynomial entropy
- Group structures?
- Spaces that it acts on, ideally: hyperbolic space,  $\mathrm{CAT}(0)$  space, etc.
- Classifications (e.g. finite order, “reducible”, “pseudo-Anosov”, etc.): via entropy, or via its action on certain spaces, etc.

Next, we explain “Nielsen realization” in the title –

## A general problem

Let  $\mathcal{D}$  be a triangulated category (e.g.  $D^b\mathrm{Coh}(X)$ ). Study  
the group of autoequivalences  $\mathrm{Aut}(\mathcal{D})$ .

(Why? It contains  $\mathrm{Aut}(X)$ , and “hidden symmetries” like spherical twists)

- Complexity: categorical entropy, categorical polynomial entropy
- Group structures?
- Spaces that it acts on, ideally: hyperbolic space,  $\mathrm{CAT}(0)$  space, etc.
- Classifications (e.g. finite order, “reducible”, “pseudo-Anosov”, etc.):  
via entropy, or via its action on certain spaces, etc.

Next, we explain “Nielsen realization” in the title –

## A general problem

Let  $\mathcal{D}$  be a triangulated category (e.g.  $D^b\mathrm{Coh}(X)$ ). Study  
the group of autoequivalences  $\mathrm{Aut}(\mathcal{D})$ .

(Why? It contains  $\mathrm{Aut}(X)$ , and “hidden symmetries” like spherical twists)

- Complexity: categorical entropy, categorical polynomial entropy
- Group structures?
- Spaces that it acts on, ideally: hyperbolic space,  $\mathrm{CAT}(0)$  space, etc.
- Classifications (e.g. finite order, “reducible”, “pseudo-Anosov”, etc.):  
via entropy, or via its action on certain spaces, etc.

Next, we explain “Nielsen realization” in the title –

## A general problem

Let  $\mathcal{D}$  be a triangulated category (e.g.  $D^b\mathrm{Coh}(X)$ ). Study  
the group of autoequivalences  $\mathrm{Aut}(\mathcal{D})$ .

(Why? It contains  $\mathrm{Aut}(X)$ , and “hidden symmetries” like spherical twists)

- Complexity: categorical entropy, categorical polynomial entropy
- Group structures?
- Spaces that it acts on, ideally: hyperbolic space,  $\mathrm{CAT}(0)$  space, etc.
- Classifications (e.g. finite order, “reducible”, “pseudo-Anosov”, etc.):  
via entropy, or via its action on certain spaces, etc.

Next, we explain “Nielsen realization” in the title –

## A general problem

Let  $\mathcal{D}$  be a triangulated category (e.g.  $D^b\mathrm{Coh}(X)$ ). Study  
the group of autoequivalences  $\mathrm{Aut}(\mathcal{D})$ .

(Why? It contains  $\mathrm{Aut}(X)$ , and “hidden symmetries” like spherical twists)

- Complexity: categorical entropy, categorical polynomial entropy
- Group structures?
- Spaces that it acts on, ideally: hyperbolic space,  $\mathrm{CAT}(0)$  space, etc.
- Classifications (e.g. finite order, “reducible”, “pseudo-Anosov”, etc.):  
via entropy, or via its action on certain spaces, etc.

Next, we explain “Nielsen realization” in the title –

# Nielsen–Thurston classification

- $\Sigma$ : Riemann surface
- $\text{MCG}(\Sigma) = \text{Diff}(\Sigma)/\text{isotopy}$ : mapping class group
- each mapping class is either:
  - ▶ finite order
  - ▶ reducible
  - ▶ pseudo-Anosov

For instance –

- elements of  $\text{MCG}(T^2) = \text{SL}(2, \mathbb{Z})$  are either:
  - ▶ elliptic (finite order)
  - ▶ parabolic (Dehn twist)
  - ▶ hyperbolic (two transverse foliations: stretch one by  $\lambda > 1$ , compress the other by  $1/\lambda$ )

Nielsen realization problem concerns mapping classes of finite order –

# Nielsen–Thurston classification

- $\Sigma$ : Riemann surface
- $\text{MCG}(\Sigma) = \text{Diff}(\Sigma)/\text{isotopy}$ : mapping class group
- each mapping class is either:
  - ▶ finite order
  - ▶ reducible
  - ▶ pseudo-Anosov

For instance –

- elements of  $\text{MCG}(T^2) = \text{SL}(2, \mathbb{Z})$  are either:
  - ▶ elliptic (finite order)
  - ▶ parabolic (Dehn twist)
  - ▶ hyperbolic (two transverse foliations: stretch one by  $\lambda > 1$ , compress the other by  $1/\lambda$ )

Nielsen realization problem concerns mapping classes of finite order –



# Nielsen–Thurston classification

- $\Sigma$ : Riemann surface
- $\text{MCG}(\Sigma) = \text{Diff}(\Sigma)/\text{isotopy}$ : mapping class group
- each mapping class is either:
  - ▶ finite order
  - ▶ reducible
  - ▶ pseudo-Anosov

For instance –

- elements of  $\text{MCG}(T^2) = \text{SL}(2, \mathbb{Z})$  are either:
  - ▶ elliptic (finite order)
  - ▶ parabolic (Dehn twist)
  - ▶ hyperbolic (two transverse foliations: stretch one by  $\lambda > 1$ , compress the other by  $1/\lambda$ )

Nielsen realization problem concerns mapping classes of finite order –

# Nielsen–Thurston classification

- $\Sigma$ : Riemann surface
- $\text{MCG}(\Sigma) = \text{Diff}(\Sigma)/\text{isotopy}$ : mapping class group
- each mapping class is either:
  - ▶ finite order
  - ▶ reducible
  - ▶ pseudo-Anosov

For instance –

- elements of  $\text{MCG}(T^2) = \text{SL}(2, \mathbb{Z})$  are either:
  - ▶ elliptic (finite order)
  - ▶ parabolic (Dehn twist)
  - ▶ hyperbolic (two transverse foliations: stretch one by  $\lambda > 1$ , compress the other by  $1/\lambda$ )

Nielsen realization problem concerns mapping classes of finite order –

# Nielsen–Thurston classification

- $\Sigma$ : Riemann surface
- $\text{MCG}(\Sigma) = \text{Diff}(\Sigma)/\text{isotopy}$ : mapping class group
- each mapping class is either:
  - ▶ finite order
  - ▶ reducible
  - ▶ pseudo-Anosov

For instance –

- elements of  $\text{MCG}(T^2) = \text{SL}(2, \mathbb{Z})$  are either:
  - ▶ elliptic (finite order)
  - ▶ parabolic (Dehn twist)
  - ▶ hyperbolic (two transverse foliations: stretch one by  $\lambda > 1$ , compress the other by  $1/\lambda$ )

Nielsen realization problem concerns mapping classes of finite order –

# Realization problem

- Nielsen asked (1923): Let  $G \subseteq \text{MCG}(\Sigma)$  be a finite subgroup. Does there always exist a lifting  $G \subseteq \text{Diff}(\Sigma)$ ? (Recall that  $\text{MCG}(\Sigma) = \text{Diff}(\Sigma)/\text{isotopy}$ ).
- Kerckhoff (1983): Yes! Moreover, there exists a metric  $g$  such that  $G \subseteq \text{Isom}(\Sigma, g)$ . Or equivalently,  $G$  fixes a point in  $\text{Teich}(\Sigma)$ . (There is a natural action of  $\text{MCG}(\Sigma)$  on  $\text{Teich}(\Sigma)$ , e.g.  $\text{MCG}(T^2) = \text{SL}(2, \mathbb{Z})$  acts on  $\text{Teich}(T^2) = \mathbb{H}$ .) (Rephrase: any finite subgroup of  $\text{MCG}(\Sigma)$  can be realized as symmetries with respect to a metric on  $\Sigma$ .)
- Farb–Looijenga (2021) also proved similar statements for K3 surfaces (under certain conditions), where  $g$  is replaced by complex structure or Ricci-flat metric on the K3 surface.

# Realization problem

- Nielsen asked (1923): Let  $G \subseteq \text{MCG}(\Sigma)$  be a finite subgroup. Does there always exist a lifting  $G \subseteq \text{Diff}(\Sigma)$ ? (Recall that  $\text{MCG}(\Sigma) = \text{Diff}(\Sigma)/\text{isotopy}$ ).
- Kerckhoff (1983): Yes! Moreover, there exists a metric  $g$  such that  $G \subseteq \text{Isom}(\Sigma, g)$ . Or equivalently,  $G$  fixes a point in  $\text{Teich}(\Sigma)$ . (There is a natural action of  $\text{MCG}(\Sigma)$  on  $\text{Teich}(\Sigma)$ , e.g.  $\text{MCG}(T^2) = \text{SL}(2, \mathbb{Z})$  acts on  $\text{Teich}(T^2) = \mathbb{H}$ .) (Rephrase: any finite subgroup of  $\text{MCG}(\Sigma)$  can be realized as symmetries with respect to a metric on  $\Sigma$ .)
- Farb–Looijenga (2021) also proved similar statements for K3 surfaces (under certain conditions), where  $g$  is replaced by complex structure or Ricci-flat metric on the K3 surface.

# Realization problem

- Nielsen asked (1923): Let  $G \subseteq \text{MCG}(\Sigma)$  be a finite subgroup. Does there always exist a lifting  $G \subseteq \text{Diff}(\Sigma)$ ?  
(Recall that  $\text{MCG}(\Sigma) = \text{Diff}(\Sigma)/\text{isotopy}$ ).
- Kerckhoff (1983): Yes! Moreover, there exists a metric  $g$  such that  $G \subseteq \text{Isom}(\Sigma, g)$ . Or equivalently,  $G$  fixes a point in  $\text{Teich}(\Sigma)$ .  
(There is a natural action of  $\text{MCG}(\Sigma)$  on  $\text{Teich}(\Sigma)$ , e.g.  $\text{MCG}(T^2) = \text{SL}(2, \mathbb{Z})$  acts on  $\text{Teich}(T^2) = \mathbb{H}$ .)  
(Rephrase: any finite subgroup of  $\text{MCG}(\Sigma)$  can be realized as symmetries with respect to a metric on  $\Sigma$ .)
- Farb–Looijenga (2021) also proved similar statements for K3 surfaces (under certain conditions), where  $g$  is replaced by complex structure or Ricci-flat metric on the K3 surface.

# Realization problem

- Nielsen asked (1923): Let  $G \subseteq \text{MCG}(\Sigma)$  be a finite subgroup. Does there always exist a lifting  $G \subseteq \text{Diff}(\Sigma)$ ?  
(Recall that  $\text{MCG}(\Sigma) = \text{Diff}(\Sigma)/\text{isotopy}$ ).
- Kerckhoff (1983): Yes! Moreover, there exists a metric  $g$  such that  $G \subseteq \text{Isom}(\Sigma, g)$ . Or equivalently,  $G$  fixes a point in  $\text{Teich}(\Sigma)$ .  
(There is a natural action of  $\text{MCG}(\Sigma)$  on  $\text{Teich}(\Sigma)$ ,  
e.g.  $\text{MCG}(T^2) = \text{SL}(2, \mathbb{Z})$  acts on  $\text{Teich}(T^2) = \mathbb{H}$ .)  
(Rephrase: any finite subgroup of  $\text{MCG}(\Sigma)$  can be realized as symmetries with respect to a metric on  $\Sigma$ .)
- Farb–Looijenga (2021) also proved similar statements for K3 surfaces (under certain conditions), where  $g$  is replaced by complex structure or Ricci-flat metric on the K3 surface.

## A categorical realization problem

Let  $\mathcal{D}$  be a triangulated category, and  $G \subseteq \operatorname{Aut}(\mathcal{D})$  be a finite subgroup. Does there exist  $\sigma \in \operatorname{Stab}(\mathcal{D})$  such that  $\Phi \cdot \sigma = \sigma$  for all  $\Phi \in G$ ?

(Rephrase: any finite subgroup of  $\operatorname{Aut}(\mathcal{D})$  can be realized as symmetries with respect to a stability condition on  $\mathcal{D}$ .)

- There is a whole dictionary of analogy between Teichmüller theory and stability conditions on triangulated categories (Haiden, Katzarkov, Kontsevich, Bridgeland, Smith, etc.). This problem is the categorical version of the Nielsen realization problem.
- When  $\mathcal{D} = D^b\operatorname{Coh}(X)$ , stability conditions on  $\mathcal{D}$  are roughly Kähler structures on  $X$ ; so this problem is similar to (but not quite the same) the mirror problem of Farb–Looijenga.



## A categorical realization problem

Let  $\mathcal{D}$  be a triangulated category, and  $G \subseteq \operatorname{Aut}(\mathcal{D})$  be a finite subgroup. Does there exist  $\sigma \in \operatorname{Stab}(\mathcal{D})$  such that  $\Phi \cdot \sigma = \sigma$  for all  $\Phi \in G$ ?

(Rephrase: any finite subgroup of  $\operatorname{Aut}(\mathcal{D})$  can be realized as symmetries with respect to a stability condition on  $\mathcal{D}$ .)

- There is a whole dictionary of analogy between Teichmüller theory and stability conditions on triangulated categories (Haiden, Katzarkov, Kontsevich, Bridgeland, Smith, etc.). This problem is the categorical version of the Nielsen realization problem.
- When  $\mathcal{D} = D^b\operatorname{Coh}(X)$ , stability conditions on  $\mathcal{D}$  are roughly Kähler structures on  $X$ ; so this problem is similar to (but not quite the same) the mirror problem of Farb–Looijenga.

# A categorical realization problem

Let  $\mathcal{D}$  be a triangulated category, and  $G \subseteq \operatorname{Aut}(\mathcal{D})$  be a finite subgroup. Does there exist  $\sigma \in \operatorname{Stab}(\mathcal{D})$  such that  $\Phi \cdot \sigma = \sigma$  for all  $\Phi \in G$ ?

(Rephrase: any finite subgroup of  $\operatorname{Aut}(\mathcal{D})$  can be realized as symmetries with respect to a stability condition on  $\mathcal{D}$ .)

- There is a whole dictionary of analogy between Teichmüller theory and stability conditions on triangulated categories (Haiden, Katzarkov, Kontsevich, Bridgeland, Smith, etc.). This problem is the categorical version of the Nielsen realization problem.
- When  $\mathcal{D} = D^b\operatorname{Coh}(X)$ , stability conditions on  $\mathcal{D}$  are roughly Kähler structures on  $X$ ; so this problem is similar to (but not quite the same) the mirror problem of Farb–Looijenga.

# A categorical realization problem

Let  $\mathcal{D}$  be a triangulated category, and  $G \subseteq \operatorname{Aut}(\mathcal{D})$  be a finite subgroup. Does there exist  $\sigma \in \operatorname{Stab}(\mathcal{D})$  such that  $\Phi \cdot \sigma = \sigma$  for all  $\Phi \in G$ ?

(Rephrase: any finite subgroup of  $\operatorname{Aut}(\mathcal{D})$  can be realized as symmetries with respect to a stability condition on  $\mathcal{D}$ .)

- There is a whole dictionary of analogy between Teichmüller theory and stability conditions on triangulated categories (Haiden, Katzarkov, Kontsevich, Bridgeland, Smith, etc.). This problem is the categorical version of the Nielsen realization problem.
- When  $\mathcal{D} = D^b\operatorname{Coh}(X)$ , stability conditions on  $\mathcal{D}$  are roughly Kähler structures on  $X$ ; so this problem is similar to (but not quite the same) the mirror problem of Farb–Looijenga.

# A dictionary of analogy

(after Gaiotto, Moore, Neitzke; Bridgeland, Smith; Dimitrov, Haiden, Katzarkov, Kontsevich, etc.)

Riemann surface $\Sigma$	Triangulated category $\mathcal{D}$
curve $C$	object $E$
$C_1 \cap C_2$	$\mathrm{Hom}(E_1, E_2)$
metric $g$	Bridgeland stability condition $\sigma$
geodesics	semistable objects
length $\ell_g(C)$	mass $m_\sigma(E)$
$\mathrm{MCG}(\Sigma)$	$\mathrm{Aut}(\mathcal{D})$
$\mathrm{Teich}(\Sigma)$	$\mathrm{Stab}(\mathcal{D})$
...	...

Many analogous statements in Teichmüller theory can be proved in the categorical setting for  $\mathcal{D} = D^b\mathrm{Coh}(\text{elliptic curve})$ . An interesting general question is whether some of these can be generalized to  $\dim \geq 2$ .

## A dictionary of analogy

(after Gaiotto, Moore, Neitzke; Bridgeland, Smith; Dimitrov, Haiden, Katzarkov, Kontsevich, etc.)

Riemann surface $\Sigma$	Triangulated category $\mathcal{D}$
curve $C$	object $E$
$C_1 \cap C_2$	$\mathrm{Hom}(E_1, E_2)$
metric $g$	Bridgeland stability condition $\sigma$
geodesics	semistable objects
length $\ell_g(C)$	mass $m_\sigma(E)$
$\mathrm{MCG}(\Sigma)$	$\mathrm{Aut}(\mathcal{D})$
$\mathrm{Teich}(\Sigma)$	$\mathrm{Stab}(\mathcal{D})$
...	...

Many analogous statements in Teichmüller theory can be proved in the categorical setting for  $\mathcal{D} = D^b\mathrm{Coh}(\text{elliptic curve})$ . An interesting general question is whether some of these can be generalized to  $\dim \geq 2$ .

# A categorical realization problem II

Let  $G \subseteq \mathrm{Aut}(\mathcal{D})/[1]$  be a finite subgroup.

Does there exist  $\sigma \in \mathrm{Stab}(\mathcal{D})/\mathbb{C}$  such that  $\Phi \cdot \sigma = \sigma$  for all  $\Phi \in G$ ?

- This statement is stronger than the previous one.
- There are many examples of  $\mathcal{D}$  where there are not many interesting finite order elements in  $\mathrm{Aut}(\mathcal{D})$ , but there are many interesting finite order elements in  $\mathrm{Aut}(\mathcal{D})/[1]$ .
- The shift functor  $[1]$  (or rather  $[2]$ ) has no classical counterpart in Teichmüller theory, so it is natural to consider this stronger question.

# A categorical realization problem II

Let  $G \subseteq \mathrm{Aut}(\mathcal{D})/[1]$  be a finite subgroup.

Does there exist  $\sigma \in \mathrm{Stab}(\mathcal{D})/\mathbb{C}$  such that  $\Phi \cdot \sigma = \sigma$  for all  $\Phi \in G$ ?

- This statement is stronger than the previous one.
- There are many examples of  $\mathcal{D}$  where there are not many interesting finite order elements in  $\mathrm{Aut}(\mathcal{D})$ , but there are many interesting finite order elements in  $\mathrm{Aut}(\mathcal{D})/[1]$ .
- The shift functor  $[1]$  (or rather  $[2]$ ) has no classical counterpart in Teichmüller theory, so it is natural to consider this stronger question.

## A categorical realization problem II

Let  $G \subseteq \operatorname{Aut}(\mathcal{D})/[1]$  be a finite subgroup.

Does there exist  $\sigma \in \operatorname{Stab}(\mathcal{D})/\mathbb{C}$  such that  $\Phi \cdot \sigma = \sigma$  for all  $\Phi \in G$ ?

- This statement is stronger than the previous one.
- There are many examples of  $\mathcal{D}$  where there are not many interesting finite order elements in  $\operatorname{Aut}(\mathcal{D})$ , but there are many interesting finite order elements in  $\operatorname{Aut}(\mathcal{D})/[1]$ .
- The shift functor  $[1]$  (or rather  $[2]$ ) has no classical counterpart in Teichmüller theory, so it is natural to consider this stronger question.



## A categorical realization problem II

Let  $G \subseteq \operatorname{Aut}(\mathcal{D})/[1]$  be a finite subgroup.

Does there exist  $\sigma \in \operatorname{Stab}(\mathcal{D})/\mathbb{C}$  such that  $\Phi \cdot \sigma = \sigma$  for all  $\Phi \in G$ ?

- This statement is stronger than the previous one.
- There are many examples of  $\mathcal{D}$  where there are not many interesting finite order elements in  $\operatorname{Aut}(\mathcal{D})$ , but there are many interesting finite order elements in  $\operatorname{Aut}(\mathcal{D})/[1]$ .
- The shift functor  $[1]$  (or rather  $[2]$ ) has no classical counterpart in Teichmüller theory, so it is natural to consider this stronger question.

# Main theorems (F.–Lai, 2023)

- The answer is yes, for  $\mathcal{D} = D^b\mathrm{Coh}(X)$  where  $X$  is a curve, a (twisted) abelian surface, a generic twisted K3 surface, or a K3 surface of Picard number  $\rho = 1$ .
- For K3 surfaces of  $\rho = 1$ , we obtain:
  - ▶ classification and counting formula of the conjugacy classes of finite subgroups of  $\mathrm{Aut}(\mathcal{D})$  and  $\mathrm{Aut}(\mathcal{D})/[1]$ ;
  - ▶ one-to-one correspondence between  $\{\text{maximal finite subgroups of } \mathrm{Aut}(\mathcal{D})/[1]\}$  and  $\{\text{elliptic points of } \mathrm{Stab}_{\mathrm{red}}^{\dagger}(\mathcal{D})/\mathbb{C}\}$   
(analogue: one-to-one correspondence between  $\{\text{maximal finite subgroups of } \mathrm{PSL}(2, \mathbb{Z})\}$  and  $\{\text{elliptic points of } \mathbb{H}\}$ )

Here,  $\mathrm{Stab}_{\mathrm{red}}^{\dagger}(\mathcal{D}) = \{(Z, P) \mid Z^2 = 0 \text{ in } N(\mathcal{D}) \otimes \mathbb{C}\} \subseteq \mathrm{Stab}^{\dagger}(\mathcal{D})$ .

## Main theorems (F.–Lai, 2023)

- The answer is yes, for  $\mathcal{D} = D^b\mathrm{Coh}(X)$  where  $X$  is a curve, a (twisted) abelian surface, a generic twisted K3 surface, or a K3 surface of Picard number  $\rho = 1$ .
- For K3 surfaces of  $\rho = 1$ , we obtain:
  - ▶ classification and counting formula of the conjugacy classes of finite subgroups of  $\mathrm{Aut}(\mathcal{D})$  and  $\mathrm{Aut}(\mathcal{D})/[1]$ ;
  - ▶ one-to-one correspondence between  $\{\text{maximal finite subgroups of } \mathrm{Aut}(\mathcal{D})/[1]\}$  and  $\{\text{elliptic points of } \mathrm{Stab}_{\mathrm{red}}^{\dagger}(\mathcal{D})/\mathbb{C}\}$   
(analogue: one-to-one correspondence between  $\{\text{maximal finite subgroups of } \mathrm{PSL}(2, \mathbb{Z})\}$  and  $\{\text{elliptic points of } \mathbb{H}\}$ )

Here,  $\mathrm{Stab}_{\mathrm{red}}^{\dagger}(\mathcal{D}) = \{(Z, P) \mid Z^2 = 0 \text{ in } N(\mathcal{D}) \otimes \mathbb{C}\} \subseteq \mathrm{Stab}^{\dagger}(\mathcal{D})$ .

## Main theorems (F.–Lai, 2023)

- The answer is yes, for  $\mathcal{D} = D^b\mathrm{Coh}(X)$  where  $X$  is a curve, a (twisted) abelian surface, a generic twisted K3 surface, or a K3 surface of Picard number  $\rho = 1$ .
- For K3 surfaces of  $\rho = 1$ , we obtain:
  - ▶ classification and counting formula of the conjugacy classes of finite subgroups of  $\mathrm{Aut}(\mathcal{D})$  and  $\mathrm{Aut}(\mathcal{D})/[1]$ ;
  - ▶ one-to-one correspondence between  $\{\text{maximal finite subgroups of } \mathrm{Aut}(\mathcal{D})/[1]\}$  and  $\{\text{elliptic points of } \mathrm{Stab}_{\mathrm{red}}^{\dagger}(\mathcal{D})/\mathbb{C}\}$   
(analogue: one-to-one correspondence between  $\{\text{maximal finite subgroups of } \mathrm{PSL}(2, \mathbb{Z})\}$  and  $\{\text{elliptic points of } \mathbb{H}\}$ )

Here,  $\mathrm{Stab}_{\mathrm{red}}^{\dagger}(\mathcal{D}) = \{(Z, P) \mid Z^2 = 0 \text{ in } N(\mathcal{D}) \otimes \mathbb{C}\} \subseteq \mathrm{Stab}^{\dagger}(\mathcal{D})$ .

# Main theorems (F.–Lai, 2023), cont'd

(Still in the case of K3 surfaces of  $\rho = 1$ )

$\Phi \in \text{Aut}(\mathcal{D})$  can be classified into (modulo quotienting certain subgroup):

- finite order up to shifts
- reducible, which further classified into:
  - ▶ “ $(-2)$ -reducible”: spherical twists  $T_S$
  - ▶ “0-reducible”: which fixes a class  $w \in N(\mathcal{D})$  with  $w^2 = 0$  (e.g.  $\otimes \mathcal{O}(1)$ )
- hyperbolic:  $\rho([\Phi]_{N(\mathcal{D})}) > 1$

Modulo certain conjectures regarding the (polynomial) entropy of the reducible autoequivalences, we expect the following trichotomy:

- finite order if and only if  $h_{\text{cat}} = h_{\text{poly}} = 0$
- reducible if and only if  $h_{\text{cat}} = 0$  and  $h_{\text{poly}} > 0$
- hyperbolic if and only if  $h_{\text{cat}} > 0$

# Main theorems (F.–Lai, 2023), cont'd

(Still in the case of K3 surfaces of  $\rho = 1$ )

$\Phi \in \text{Aut}(\mathcal{D})$  can be classified into (modulo quotienting certain subgroup):

- finite order up to shifts
- reducible, which further classified into:
  - ▶ “ $(-2)$ -reducible”: spherical twists  $T_S$
  - ▶ “0-reducible”: which fixes a class  $w \in N(\mathcal{D})$  with  $w^2 = 0$  (e.g.  $\otimes \mathcal{O}(1)$ )
- hyperbolic:  $\rho([\Phi]_{N(\mathcal{D})}) > 1$

Modulo certain conjectures regarding the (polynomial) entropy of the reducible autoequivalences, we expect the following trichotomy:

- finite order if and only if  $h_{\text{cat}} = h_{\text{poly}} = 0$
- reducible if and only if  $h_{\text{cat}} = 0$  and  $h_{\text{poly}} > 0$
- hyperbolic if and only if  $h_{\text{cat}} > 0$

## Main theorems (F.–Lai, 2023), cont'd

(Still in the case of K3 surfaces of  $\rho = 1$ )

$\Phi \in \text{Aut}(\mathcal{D})$  can be classified into (modulo quotienting certain subgroup):

- finite order up to shifts
- reducible, which further classified into:
  - ▶ “ $(-2)$ -reducible”: spherical twists  $T_S$
  - ▶ “0-reducible”: which fixes a class  $w \in N(\mathcal{D})$  with  $w^2 = 0$  (e.g.  $\otimes \mathcal{O}(1)$ )
- hyperbolic:  $\rho([\Phi]_{N(\mathcal{D})}) > 1$

Modulo certain conjectures regarding the (polynomial) entropy of the reducible autoequivalences, we expect the following trichotomy:

- finite order if and only if  $h_{\text{cat}} = h_{\text{poly}} = 0$
- reducible if and only if  $h_{\text{cat}} = 0$  and  $h_{\text{poly}} > 0$
- hyperbolic if and only if  $h_{\text{cat}} > 0$

## Main theorems (F.–Lai, 2023), cont'd

(Still in the case of K3 surfaces of  $\rho = 1$ )

$\Phi \in \text{Aut}(\mathcal{D})$  can be classified into (modulo quotienting certain subgroup):

- finite order up to shifts
- reducible, which further classified into:
  - ▶ “ $(-2)$ -reducible”: spherical twists  $T_S$
  - ▶ “0-reducible”: which fixes a class  $w \in N(\mathcal{D})$  with  $w^2 = 0$  (e.g.  $\otimes \mathcal{O}(1)$ )
- hyperbolic:  $\rho([\Phi]_{N(\mathcal{D})}) > 1$

Modulo certain conjectures regarding the (polynomial) entropy of the reducible autoequivalences, we expect the following trichotomy:

- finite order if and only if  $h_{\text{cat}} = h_{\text{poly}} = 0$
- reducible if and only if  $h_{\text{cat}} = 0$  and  $h_{\text{poly}} > 0$
- hyperbolic if and only if  $h_{\text{cat}} > 0$



# Main theorems (F.–Lai, 2023), cont'd

(Still in the case of K3 surfaces of  $\rho = 1$ )

$\Phi \in \text{Aut}(\mathcal{D})$  can be classified into (modulo quotienting certain subgroup):

- finite order up to shifts
- reducible, which further classified into:
  - ▶ “ $(-2)$ -reducible”: spherical twists  $T_S$
  - ▶ “0-reducible”: which fixes a class  $w \in N(\mathcal{D})$  with  $w^2 = 0$  (e.g.  $\otimes \mathcal{O}(1)$ )
- hyperbolic:  $\rho([\Phi]_{N(\mathcal{D})}) > 1$

Modulo certain conjectures regarding the (polynomial) entropy of the reducible autoequivalences, we expect the following trichotomy:

- finite order if and only if  $h_{\text{cat}} = h_{\text{poly}} = 0$
- reducible if and only if  $h_{\text{cat}} = 0$  and  $h_{\text{poly}} > 0$
- hyperbolic if and only if  $h_{\text{cat}} > 0$

## Main theorems (F.–Lai, 2023), cont'd

(Still in the case of K3 surfaces of  $\rho = 1$ )

$\Phi \in \text{Aut}(\mathcal{D})$  can be classified into (modulo quotienting certain subgroup):

- finite order up to shifts
- reducible, which further classified into:
  - ▶ “ $(-2)$ -reducible”: spherical twists  $T_S$
  - ▶ “0-reducible”: which fixes a class  $w \in N(\mathcal{D})$  with  $w^2 = 0$  (e.g.  $\otimes \mathcal{O}(1)$ )
- hyperbolic:  $\rho([\Phi]_{N(\mathcal{D})}) > 1$

Modulo certain conjectures regarding the (polynomial) entropy of the reducible autoequivalences, we expect the following trichotomy:

- finite order if and only if  $h_{\text{cat}} = h_{\text{poly}} = 0$
- reducible if and only if  $h_{\text{cat}} = 0$  and  $h_{\text{poly}} > 0$
- hyperbolic if and only if  $h_{\text{cat}} > 0$

# Strategy and Difficulties

For K3 or abelian surfaces, Bridgeland (2008) showed that there is an  $\text{Aut}(\mathcal{D})$ -equivariant covering map

$$\text{Stab}_{\text{red}}^+(\mathcal{D})/\mathbb{C} \xrightarrow{\pi} Q_0^+(\mathcal{D})$$

where  $Q_0^+(\mathcal{D}) = \{v \in \mathbb{P}(N(\mathcal{D}) \otimes \mathbb{C}) \mid v^2 = 0, v\bar{v} > 0\} \setminus \bigcup_{\delta^2 = -2} \delta^\perp$ .

- For abelian surfaces, there is no spherical objects in  $\mathcal{D}$ , so:

- ▶ we do not need to remove " $\delta^\perp$ " since there is no  $(-2)$ -classes
- ▶  $\pi$  is an isomorphism

and it is not hard to show that finite subgroups of  $\text{Aut}(\mathcal{D})$  fix a point in  $Q_0^+(\mathcal{D})$  using basic Lie theory.

- For K3 surfaces, one needs to resolve the following two issues:

- ▶ the fixed points of finite subgroups of  $\text{Aut}(\mathcal{D})$  need to avoid " $\delta^\perp$ "
- ▶  $\text{Deck}(\pi)$  is much more complicated

It turns out that both issues can be resolved for  $\rho = 1$  –

# Strategy and Difficulties

For K3 or abelian surfaces, Bridgeland (2008) showed that there is an  $\text{Aut}(\mathcal{D})$ -equivariant covering map

$$\text{Stab}_{\text{red}}^+(\mathcal{D})/\mathbb{C} \xrightarrow{\pi} Q_0^+(\mathcal{D})$$

where  $Q_0^+(\mathcal{D}) = \{v \in \mathbb{P}(N(\mathcal{D}) \otimes \mathbb{C}) \mid v^2 = 0, v\bar{v} > 0\} \setminus \bigcup_{\delta^2 = -2} \delta^\perp$ .

- For abelian surfaces, there is no spherical objects in  $\mathcal{D}$ , so:

- ▶ we do not need to remove " $\delta^\perp$ " since there is no  $(-2)$ -classes
- ▶  $\pi$  is an isomorphism

and it is not hard to show that finite subgroups of  $\text{Aut}(\mathcal{D})$  fix a point in  $Q_0^+(\mathcal{D})$  using basic Lie theory.

- For K3 surfaces, one needs to resolve the following two issues:

- ▶ the fixed points of finite subgroups of  $\text{Aut}(\mathcal{D})$  need to avoid " $\delta^\perp$ "
- ▶  $\text{Deck}(\pi)$  is much more complicated

It turns out that both issues can be resolved for  $\rho = 1$  –

# Strategy and Difficulties

For K3 or abelian surfaces, Bridgeland (2008) showed that there is an  $\text{Aut}(\mathcal{D})$ -equivariant covering map

$$\text{Stab}_{\text{red}}^+(\mathcal{D})/\mathbb{C} \xrightarrow{\pi} Q_0^+(\mathcal{D})$$

where  $Q_0^+(\mathcal{D}) = \{v \in \mathbb{P}(N(\mathcal{D}) \otimes \mathbb{C}) \mid v^2 = 0, v\bar{v} > 0\} \setminus \bigcup_{\delta^2 = -2} \delta^\perp$ .

- For abelian surfaces, there is no spherical objects in  $\mathcal{D}$ , so:
  - ▶ we do not need to remove “ $\delta^\perp$ ” since there is no  $(-2)$ -classes
  - ▶  $\pi$  is an isomorphism

and it is not hard to show that finite subgroups of  $\text{Aut}(\mathcal{D})$  fix a point in  $Q_0^+(\mathcal{D})$  using basic Lie theory.

- For K3 surfaces, one needs to resolve the following two issues:
  - ▶ the fixed points of finite subgroups of  $\text{Aut}(\mathcal{D})$  need to avoid “ $\delta^\perp$ ”
  - ▶  $\text{Deck}(\pi)$  is much more complicated

It turns out that both issues can be resolved for  $\rho = 1$  –

# Strategy and Difficulties

For K3 or abelian surfaces, Bridgeland (2008) showed that there is an  $\text{Aut}(\mathcal{D})$ -equivariant covering map

$$\text{Stab}_{\text{red}}^+(\mathcal{D})/\mathbb{C} \xrightarrow{\pi} Q_0^+(\mathcal{D})$$

where  $Q_0^+(\mathcal{D}) = \{v \in \mathbb{P}(N(\mathcal{D}) \otimes \mathbb{C}) \mid v^2 = 0, v\bar{v} > 0\} \setminus \bigcup_{\delta^2 = -2} \delta^\perp$ .

- For abelian surfaces, there is no spherical objects in  $\mathcal{D}$ , so:
  - ▶ we do not need to remove “ $\delta^\perp$ ” since there is no  $(-2)$ -classes
  - ▶  $\pi$  is an isomorphism

and it is not hard to show that finite subgroups of  $\text{Aut}(\mathcal{D})$  fix a point in  $Q_0^+(\mathcal{D})$  using basic Lie theory.

- For K3 surfaces, one needs to resolve the following two issues:
  - ▶ the fixed points of finite subgroups of  $\text{Aut}(\mathcal{D})$  need to avoid “ $\delta^\perp$ ”
  - ▶  $\text{Deck}(\pi)$  is much more complicated

It turns out that both issues can be resolved for  $\rho = 1 -$

## Strategy and Difficulties

For K3 or abelian surfaces, Bridgeland (2008) showed that there is an  $\text{Aut}(\mathcal{D})$ -equivariant covering map

$$\text{Stab}_{\text{red}}^+(\mathcal{D})/\mathbb{C} \xrightarrow{\pi} Q_0^+(\mathcal{D})$$

where  $Q_0^+(\mathcal{D}) = \{v \in \mathbb{P}(N(\mathcal{D}) \otimes \mathbb{C}) \mid v^2 = 0, v\bar{v} > 0\} \setminus \bigcup_{\delta^2 = -2} \delta^\perp$ .

- For abelian surfaces, there is no spherical objects in  $\mathcal{D}$ , so:
  - ▶ we do not need to remove “ $\delta^\perp$ ” since there is no  $(-2)$ -classes
  - ▶  $\pi$  is an isomorphism

and it is not hard to show that finite subgroups of  $\text{Aut}(\mathcal{D})$  fix a point in  $Q_0^+(\mathcal{D})$  using basic Lie theory.

- For K3 surfaces, one needs to resolve the following two issues:
  - ▶ the fixed points of finite subgroups of  $\text{Aut}(\mathcal{D})$  need to avoid “ $\delta^\perp$ ”
  - ▶  $\text{Deck}(\pi)$  is much more complicated

It turns out that both issues can be resolved for  $\rho = 1$  –

# Strategy and Difficulties

For K3 or abelian surfaces, Bridgeland (2008) showed that there is an  $\text{Aut}(\mathcal{D})$ -equivariant covering map

$$\text{Stab}_{\text{red}}^+(\mathcal{D})/\mathbb{C} \xrightarrow{\pi} Q_0^+(\mathcal{D})$$

where  $Q_0^+(\mathcal{D}) = \{v \in \mathbb{P}(N(\mathcal{D}) \otimes \mathbb{C}) \mid v^2 = 0, v\bar{v} > 0\} \setminus \bigcup_{\delta^2 = -2} \delta^\perp$ .

- For abelian surfaces, there is no spherical objects in  $\mathcal{D}$ , so:
  - ▶ we do not need to remove “ $\delta^\perp$ ” since there is no  $(-2)$ -classes
  - ▶  $\pi$  is an isomorphism

and it is not hard to show that finite subgroups of  $\text{Aut}(\mathcal{D})$  fix a point in  $Q_0^+(\mathcal{D})$  using basic Lie theory.

- For K3 surfaces, one needs to resolve the following two issues:
  - ▶ the fixed points of finite subgroups of  $\text{Aut}(\mathcal{D})$  need to avoid “ $\delta^\perp$ ”
  - ▶  $\text{Deck}(\pi)$  is much more complicated

It turns out that both issues can be resolved for  $\rho = 1$  –



# Strategy and Difficulties

For K3 or abelian surfaces, Bridgeland (2008) showed that there is an  $\text{Aut}(\mathcal{D})$ -equivariant covering map

$$\text{Stab}_{\text{red}}^+(\mathcal{D})/\mathbb{C} \xrightarrow{\pi} Q_0^+(\mathcal{D})$$

where  $Q_0^+(\mathcal{D}) = \{v \in \mathbb{P}(N(\mathcal{D}) \otimes \mathbb{C}) \mid v^2 = 0, v\bar{v} > 0\} \setminus \bigcup_{\delta^2 = -2} \delta^\perp$ .

- For abelian surfaces, there is no spherical objects in  $\mathcal{D}$ , so:
  - ▶ we do not need to remove “ $\delta^\perp$ ” since there is no  $(-2)$ -classes
  - ▶  $\pi$  is an isomorphism

and it is not hard to show that finite subgroups of  $\text{Aut}(\mathcal{D})$  fix a point in  $Q_0^+(\mathcal{D})$  using basic Lie theory.

- For K3 surfaces, one needs to resolve the following two issues:
  - ▶ the fixed points of finite subgroups of  $\text{Aut}(\mathcal{D})$  need to avoid “ $\delta^\perp$ ”
  - ▶  $\text{Deck}(\pi)$  is much more complicated

It turns out that both issues can be resolved for  $\rho = 1$  –

# Strategy and Difficulties

For K3 or abelian surfaces, Bridgeland (2008) showed that there is an  $\text{Aut}(\mathcal{D})$ -equivariant covering map

$$\text{Stab}_{\text{red}}^+(\mathcal{D})/\mathbb{C} \xrightarrow{\pi} Q_0^+(\mathcal{D})$$

where  $Q_0^+(\mathcal{D}) = \{v \in \mathbb{P}(N(\mathcal{D}) \otimes \mathbb{C}) \mid v^2 = 0, v\bar{v} > 0\} \setminus \bigcup_{\delta^2 = -2} \delta^\perp$ .

- For abelian surfaces, there is no spherical objects in  $\mathcal{D}$ , so:
  - ▶ we do not need to remove “ $\delta^\perp$ ” since there is no  $(-2)$ -classes
  - ▶  $\pi$  is an isomorphism

and it is not hard to show that finite subgroups of  $\text{Aut}(\mathcal{D})$  fix a point in  $Q_0^+(\mathcal{D})$  using basic Lie theory.

- For K3 surfaces, one needs to resolve the following two issues:
  - ▶ the fixed points of finite subgroups of  $\text{Aut}(\mathcal{D})$  need to avoid “ $\delta^\perp$ ”
  - ▶  $\text{Deck}(\pi)$  is much more complicated

It turns out that both issues can be resolved for  $\rho = 1$  –

# Strategy and Difficulties

For K3 or abelian surfaces, Bridgeland (2008) showed that there is an  $\text{Aut}(\mathcal{D})$ -equivariant covering map

$$\text{Stab}_{\text{red}}^+(\mathcal{D})/\mathbb{C} \xrightarrow{\pi} Q_0^+(\mathcal{D})$$

where  $Q_0^+(\mathcal{D}) = \{v \in \mathbb{P}(N(\mathcal{D}) \otimes \mathbb{C}) \mid v^2 = 0, v\bar{v} > 0\} \setminus \bigcup_{\delta^2 = -2} \delta^\perp$ .

- For abelian surfaces, there is no spherical objects in  $\mathcal{D}$ , so:
  - ▶ we do not need to remove “ $\delta^\perp$ ” since there is no  $(-2)$ -classes
  - ▶  $\pi$  is an isomorphism

and it is not hard to show that finite subgroups of  $\text{Aut}(\mathcal{D})$  fix a point in  $Q_0^+(\mathcal{D})$  using basic Lie theory.

- For K3 surfaces, one needs to resolve the following two issues:
  - ▶ the fixed points of finite subgroups of  $\text{Aut}(\mathcal{D})$  need to avoid “ $\delta^\perp$ ”
  - ▶  $\text{Deck}(\pi)$  is much more complicated

It turns out that both issues can be resolved for  $\rho = 1$  –

## Avoiding $\delta^\perp$

Suppose  $X$  is a K3 surface of  $\rho = 1$  and degree  $2n$ .

- We have  $Q_0^+(\mathcal{D}) \cong \mathbb{H} \setminus \text{"}(-2)\text{-points"}$ .
- By Dolgachev (1996) and Kawatani (2014), the action of  $\text{Aut}(\mathcal{D})$  on  $Q_0^+(\mathcal{D})$  factors through  $\text{Im}(\text{Aut}(\mathcal{D}) \xrightarrow{f} \text{PSL}(2, \mathbb{R})) = \Gamma_0^+(n)$  the Fricke modular group, where  $\Gamma_0^+(n) = \left\langle \Gamma_0(n), \begin{bmatrix} & -1/\sqrt{n} \\ \sqrt{n} & \end{bmatrix} =: \omega_n \right\rangle$ .

We showed that the following statements are equivalent:

- $f(\Phi)$  fixes a  $(-2)$ -point in  $\mathbb{H}$
- $f(\Phi)$  is an involution, and  $f(\Phi) = g_0 \omega_n$  for some  $g_0 \in \Gamma_0(n)$
- $\Phi = T_S \Psi$  for some spherical object  $S$  and some  $\Psi \in \text{Deck}(\pi)$

Moreover, we showed that autoequivalences of the form  $T_S \Psi$  must be of infinite order in  $\text{Aut}(\mathcal{D})/[1]$ . This resolves the first issue.

## Avoiding $\delta^\perp$

Suppose  $X$  is a K3 surface of  $\rho = 1$  and degree  $2n$ .

- We have  $Q_0^+(\mathcal{D}) \cong \mathbb{H} \setminus \text{"}(-2)\text{-points"}$ .
- By Dolgachev (1996) and Kawatani (2014), the action of  $\text{Aut}(\mathcal{D})$  on  $Q_0^+(\mathcal{D})$  factors through  $\text{Im}(\text{Aut}(\mathcal{D}) \xrightarrow{f} \text{PSL}(2, \mathbb{R})) = \Gamma_0^+(n)$  the Fricke modular group, where  $\Gamma_0^+(n) = \left\langle \Gamma_0(n), \begin{bmatrix} & -1/\sqrt{n} \\ \sqrt{n} & \end{bmatrix} =: \omega_n \right\rangle$ .

We showed that the following statements are equivalent:

- $f(\Phi)$  fixes a  $(-2)$ -point in  $\mathbb{H}$
- $f(\Phi)$  is an involution, and  $f(\Phi) = g_0 \omega_n$  for some  $g_0 \in \Gamma_0(n)$
- $\Phi = T_S \Psi$  for some spherical object  $S$  and some  $\Psi \in \text{Deck}(\pi)$

Moreover, we showed that autoequivalences of the form  $T_S \Psi$  must be of infinite order in  $\text{Aut}(\mathcal{D})/[1]$ . This resolves the first issue.

## Avoiding $\delta^\perp$

Suppose  $X$  is a K3 surface of  $\rho = 1$  and degree  $2n$ .

- We have  $Q_0^+(\mathcal{D}) \cong \mathbb{H} \setminus \text{"}(-2)\text{-points"}$ .
- By Dolgachev (1996) and Kawatani (2014), the action of  $\text{Aut}(\mathcal{D})$  on  $Q_0^+(\mathcal{D})$  factors through  $\text{Im}(\text{Aut}(\mathcal{D}) \xrightarrow{f} \text{PSL}(2, \mathbb{R})) = \Gamma_0^+(n)$  the Fricke modular group, where  $\Gamma_0^+(n) = \left\langle \Gamma_0(n), \begin{bmatrix} & -1/\sqrt{n} \\ \sqrt{n} & \end{bmatrix} =: \omega_n \right\rangle$ .

We showed that the following statements are equivalent:

- $f(\Phi)$  fixes a  $(-2)$ -point in  $\mathbb{H}$
- $f(\Phi)$  is an involution, and  $f(\Phi) = g_0 \omega_n$  for some  $g_0 \in \Gamma_0(n)$
- $\Phi = T_S \Psi$  for some spherical object  $S$  and some  $\Psi \in \text{Deck}(\pi)$

Moreover, we showed that autoequivalences of the form  $T_S \Psi$  must be of infinite order in  $\text{Aut}(\mathcal{D})/[1]$ . This resolves the first issue.

## Avoiding $\delta^\perp$

Suppose  $X$  is a K3 surface of  $\rho = 1$  and degree  $2n$ .

- We have  $Q_0^+(\mathcal{D}) \cong \mathbb{H} \setminus \text{"}(-2)\text{-points"}$ .
- By Dolgachev (1996) and Kawatani (2014), the action of  $\text{Aut}(\mathcal{D})$  on  $Q_0^+(\mathcal{D})$  factors through  $\text{Im}(\text{Aut}(\mathcal{D}) \xrightarrow{f} \text{PSL}(2, \mathbb{R})) = \Gamma_0^+(n)$  the Fricke modular group, where  $\Gamma_0^+(n) = \left\langle \Gamma_0(n), \begin{bmatrix} & -1/\sqrt{n} \\ \sqrt{n} & \end{bmatrix} =: \omega_n \right\rangle$ .

We showed that the following statements are equivalent:

- $f(\Phi)$  fixes a  $(-2)$ -point in  $\mathbb{H}$
- $f(\Phi)$  is an involution, and  $f(\Phi) = g_0 \omega_n$  for some  $g_0 \in \Gamma_0(n)$
- $\Phi = T_S \Psi$  for some spherical object  $S$  and some  $\Psi \in \text{Deck}(\pi)$

Moreover, we showed that autoequivalences of the form  $T_S \Psi$  must be of infinite order in  $\text{Aut}(\mathcal{D})/[1]$ . This resolves the first issue.

## Avoiding $\delta^\perp$

Suppose  $X$  is a K3 surface of  $\rho = 1$  and degree  $2n$ .

- We have  $Q_0^+(\mathcal{D}) \cong \mathbb{H} \setminus \text{"}(-2)\text{-points"}$ .
- By Dolgachev (1996) and Kawatani (2014), the action of  $\text{Aut}(\mathcal{D})$  on  $Q_0^+(\mathcal{D})$  factors through  $\text{Im}(\text{Aut}(\mathcal{D}) \xrightarrow{f} \text{PSL}(2, \mathbb{R})) = \Gamma_0^+(n)$  the Fricke modular group, where  $\Gamma_0^+(n) = \left\langle \Gamma_0(n), \begin{bmatrix} & -1/\sqrt{n} \\ \sqrt{n} & \end{bmatrix} =: \omega_n \right\rangle$ .

We showed that the following statements are equivalent:

- $f(\Phi)$  fixes a  $(-2)$ -point in  $\mathbb{H}$
- $f(\Phi)$  is an involution, and  $f(\Phi) = g_0 \omega_n$  for some  $g_0 \in \Gamma_0(n)$
- $\Phi = T_S \Psi$  for some spherical object  $S$  and some  $\Psi \in \text{Deck}(\pi)$

Moreover, we showed that autoequivalences of the form  $T_S \Psi$  must be of infinite order in  $\text{Aut}(\mathcal{D})/[1]$ . This resolves the first issue.



## Avoiding $\delta^\perp$

Suppose  $X$  is a K3 surface of  $\rho = 1$  and degree  $2n$ .

- We have  $Q_0^+(\mathcal{D}) \cong \mathbb{H} \setminus \text{"}(-2)\text{-points"}$ .
- By Dolgachev (1996) and Kawatani (2014), the action of  $\text{Aut}(\mathcal{D})$  on  $Q_0^+(\mathcal{D})$  factors through  $\text{Im}(\text{Aut}(\mathcal{D}) \xrightarrow{f} \text{PSL}(2, \mathbb{R})) = \Gamma_0^+(n)$  the Fricke modular group, where  $\Gamma_0^+(n) = \left\langle \Gamma_0(n), \begin{bmatrix} & -1/\sqrt{n} \\ \sqrt{n} & \end{bmatrix} =: \omega_n \right\rangle$ .

We showed that the following statements are equivalent:

- $f(\Phi)$  fixes a  $(-2)$ -point in  $\mathbb{H}$
- $f(\Phi)$  is an involution, and  $f(\Phi) = g_0 \omega_n$  for some  $g_0 \in \Gamma_0(n)$
- $\Phi = T_S \Psi$  for some spherical object  $S$  and some  $\Psi \in \text{Deck}(\pi)$

Moreover, we showed that autoequivalences of the form  $T_S \Psi$  must be of infinite order in  $\text{Aut}(\mathcal{D})/[1]$ . This resolves the first issue.

## Avoiding $\delta^\perp$

Suppose  $X$  is a K3 surface of  $\rho = 1$  and degree  $2n$ .

- We have  $Q_0^+(\mathcal{D}) \cong \mathbb{H} \setminus \text{"}(-2)\text{-points"}$ .
- By Dolgachev (1996) and Kawatani (2014), the action of  $\text{Aut}(\mathcal{D})$  on  $Q_0^+(\mathcal{D})$  factors through  $\text{Im}(\text{Aut}(\mathcal{D}) \xrightarrow{f} \text{PSL}(2, \mathbb{R})) = \Gamma_0^+(n)$  the Fricke modular group, where  $\Gamma_0^+(n) = \left\langle \Gamma_0(n), \begin{bmatrix} & -1/\sqrt{n} \\ \sqrt{n} & \end{bmatrix} =: \omega_n \right\rangle$ .

We showed that the following statements are equivalent:

- $f(\Phi)$  fixes a  $(-2)$ -point in  $\mathbb{H}$
- $f(\Phi)$  is an involution, and  $f(\Phi) = g_0 \omega_n$  for some  $g_0 \in \Gamma_0(n)$
- $\Phi = T_S \Psi$  for some spherical object  $S$  and some  $\Psi \in \text{Deck}(\pi)$

Moreover, we showed that autoequivalences of the form  $T_S \Psi$  must be of infinite order in  $\text{Aut}(\mathcal{D})/[1]$ . This resolves the first issue.

# Lifting of fixed points

- Let  $\Phi \in \text{Aut}(\mathcal{D})/[1]$  be of finite order, then the previous discussion shows that it fixes a point in  $Q_0^+(\mathcal{D})$ .
- Kawatani (2019):  $\pi_1(Q_0^+(\mathcal{D})) \cong \star_{\text{free}} T_S^2$ .
- Bayer–Bridgeland (2017):  $\text{Stab}_{\text{red}}^+(\mathcal{D})/\mathbb{C}$  is contractible.
- Combining these two results, we have  $\text{Deck}(\pi) \cong \star_{\text{free}} T_S^2$ .
- We showed that this implies the fixed point of  $\Phi$  in  $Q_0^+(D)$  can be lifted to a fixed point in  $\text{Stab}_{\text{red}}^+(\mathcal{D})/\mathbb{C}$ , which proves the realization problem for cyclic subgroups of  $\text{Aut}(\mathcal{D})/[1]$ .
- Finally, we showed that any finite subgroup of  $\text{Aut}(\mathcal{D})/[1]$  is cyclic using classification results of finite subgroups of  $\Gamma_0^+(n)$  by Fricke. This concludes the proof.

# Lifting of fixed points

- Let  $\Phi \in \text{Aut}(\mathcal{D})/[1]$  be of finite order, then the previous discussion shows that it fixes a point in  $Q_0^+(\mathcal{D})$ .
- Kawatani (2019):  $\pi_1(Q_0^+(\mathcal{D})) \cong \star_{\text{free}} T_S^2$ .
- Bayer–Bridgeland (2017):  $\text{Stab}_{\text{red}}^+(\mathcal{D})/\mathbb{C}$  is contractible.
- Combining these two results, we have  $\text{Deck}(\pi) \cong \star_{\text{free}} T_S^2$ .
- We showed that this implies the fixed point of  $\Phi$  in  $Q_0^+(\mathcal{D})$  can be lifted to a fixed point in  $\text{Stab}_{\text{red}}^+(\mathcal{D})/\mathbb{C}$ , which proves the realization problem for cyclic subgroups of  $\text{Aut}(\mathcal{D})/[1]$ .
- Finally, we showed that any finite subgroup of  $\text{Aut}(\mathcal{D})/[1]$  is cyclic using classification results of finite subgroups of  $\Gamma_0^+(n)$  by Fricke. This concludes the proof.

## Lifting of fixed points

- Let  $\Phi \in \operatorname{Aut}(\mathcal{D})/[1]$  be of finite order, then the previous discussion shows that it fixes a point in  $Q_0^+(\mathcal{D})$ .
- Kawatani (2019):  $\pi_1(Q_0^+(\mathcal{D})) \cong \star_{\text{free}} T_S^2$ .
- Bayer–Bridgeland (2017):  $\operatorname{Stab}_{\text{red}}^\dagger(\mathcal{D})/\mathbb{C}$  is contractible.
- Combining these two results, we have  $\operatorname{Deck}(\pi) \cong \star_{\text{free}} T_S^2$ .
- We showed that this implies the fixed point of  $\Phi$  in  $Q_0^+(\mathcal{D})$  can be lifted to a fixed point in  $\operatorname{Stab}_{\text{red}}^\dagger(\mathcal{D})/\mathbb{C}$ , which proves the realization problem for cyclic subgroups of  $\operatorname{Aut}(\mathcal{D})/[1]$ .
- Finally, we showed that any finite subgroup of  $\operatorname{Aut}(\mathcal{D})/[1]$  is cyclic using classification results of finite subgroups of  $\Gamma_0^+(n)$  by Fricke. This concludes the proof.

## Lifting of fixed points

- Let  $\Phi \in \text{Aut}(\mathcal{D})/[1]$  be of finite order, then the previous discussion shows that it fixes a point in  $Q_0^+(\mathcal{D})$ .
- Kawatani (2019):  $\pi_1(Q_0^+(\mathcal{D})) \cong \star_{\text{free}} T_S^2$ .
- Bayer–Bridgeland (2017):  $\text{Stab}_{\text{red}}^+(\mathcal{D})/\mathbb{C}$  is contractible.
- Combining these two results, we have  $\text{Deck}(\pi) \cong \star_{\text{free}} T_S^2$ .
- We showed that this implies the fixed point of  $\Phi$  in  $Q_0^+(\mathcal{D})$  can be lifted to a fixed point in  $\text{Stab}_{\text{red}}^+(\mathcal{D})/\mathbb{C}$ , which proves the realization problem for cyclic subgroups of  $\text{Aut}(\mathcal{D})/[1]$ .
- Finally, we showed that any finite subgroup of  $\text{Aut}(\mathcal{D})/[1]$  is cyclic using classification results of finite subgroups of  $\Gamma_0^+(n)$  by Fricke. This concludes the proof.

## Lifting of fixed points

- Let  $\Phi \in \operatorname{Aut}(\mathcal{D})/[1]$  be of finite order, then the previous discussion shows that it fixes a point in  $Q_0^+(\mathcal{D})$ .
- Kawatani (2019):  $\pi_1(Q_0^+(\mathcal{D})) \cong \star_{\text{free}} T_S^2$ .
- Bayer–Bridgeland (2017):  $\operatorname{Stab}_{\text{red}}^\dagger(\mathcal{D})/\mathbb{C}$  is contractible.
- Combining these two results, we have  $\operatorname{Deck}(\pi) \cong \star_{\text{free}} T_S^2$ .
- We showed that this implies the fixed point of  $\Phi$  in  $Q_0^+(\mathcal{D})$  can be lifted to a fixed point in  $\operatorname{Stab}_{\text{red}}^\dagger(\mathcal{D})/\mathbb{C}$ , which proves the realization problem for cyclic subgroups of  $\operatorname{Aut}(\mathcal{D})/[1]$ .
- Finally, we showed that any finite subgroup of  $\operatorname{Aut}(\mathcal{D})/[1]$  is cyclic using classification results of finite subgroups of  $\Gamma_0^+(n)$  by Fricke. This concludes the proof.

## Lifting of fixed points

- Let  $\Phi \in \operatorname{Aut}(\mathcal{D})/[1]$  be of finite order, then the previous discussion shows that it fixes a point in  $Q_0^+(\mathcal{D})$ .
- Kawatani (2019):  $\pi_1(Q_0^+(\mathcal{D})) \cong \star_{\text{free}} T_S^2$ .
- Bayer–Bridgeland (2017):  $\operatorname{Stab}_{\text{red}}^\dagger(\mathcal{D})/\mathbb{C}$  is contractible.
- Combining these two results, we have  $\operatorname{Deck}(\pi) \cong \star_{\text{free}} T_S^2$ .
- We showed that this implies the fixed point of  $\Phi$  in  $Q_0^+(\mathcal{D})$  can be lifted to a fixed point in  $\operatorname{Stab}_{\text{red}}^\dagger(\mathcal{D})/\mathbb{C}$ , which proves the realization problem for cyclic subgroups of  $\operatorname{Aut}(\mathcal{D})/[1]$ .
- Finally, we showed that any finite subgroup of  $\operatorname{Aut}(\mathcal{D})/[1]$  is cyclic using classification results of finite subgroups of  $\Gamma_0^+(n)$  by Fricke. This concludes the proof.



## A few further problems

- Do 0-reducible autoequivalences have zero entropy? ( $h(\otimes \mathcal{O}(1)) = 0$ )
- Generalize realization results to:
  - ▶ general special cubic fourfolds  $\mathrm{Ku}(X)$
  - ▶ K3 surfaces of Picard number  $\rho \geq 2$
  - ▶ ...?

**Thank you for your attention!**