

# Invariants of categorical dynamical systems

Yu-Wei Fan (YMSC, Tsinghua)

at Shanghai, ICCM  
January 3, 2024

**Goal of this talk:** Explain reasons that “categorical dynamical systems” could be interesting.

A (discrete) **dynamical system** is a pair  $(X, \phi)$  where

- $\phi: X \rightarrow X$  preserves certain mathematical structures on  $X$ .

We would like to study the long-term behavior of  $\phi^n$  under large iterations.

Examples:

- A linear self-map  $T: V \rightarrow V$  of a vector space  $V$ .
- A continuous self-map  $f: X \rightarrow X$  of a compact metric space  $X$ .
- A holomorphic self-map  $f: X \rightarrow X$  of a compact Kähler manifold  $X$ .
- An endofunctor  $F: \mathcal{D} \rightarrow \mathcal{D}$  of a triangulated category  $\mathcal{D}$ .

**Goal of this talk:** Explain reasons that “categorical dynamical systems” could be interesting.

A (discrete) **dynamical system** is a pair  $(X, \phi)$  where

- $\phi: X \rightarrow X$  preserves certain mathematical structures on  $X$ .

We would like to study the long-term behavior of  $\phi^n$  under large iterations.

Examples:

- A linear self-map  $T: V \rightarrow V$  of a vector space  $V$ .
- A continuous self-map  $f: X \rightarrow X$  of a compact metric space  $X$ .
- A holomorphic self-map  $f: X \rightarrow X$  of a compact Kähler manifold  $X$ .
- An endofunctor  $F: \mathcal{D} \rightarrow \mathcal{D}$  of a triangulated category  $\mathcal{D}$ .

**Goal of this talk:** Explain reasons that “categorical dynamical systems” could be interesting.

A (discrete) **dynamical system** is a pair  $(X, \phi)$  where

- $\phi: X \rightarrow X$  preserves certain mathematical structures on  $X$ .

We would like to study the long-term behavior of  $\phi^n$  under large iterations.

Examples:

- A linear self-map  $T: V \rightarrow V$  of a vector space  $V$ .
- A continuous self-map  $f: X \rightarrow X$  of a compact metric space  $X$ .
- A holomorphic self-map  $f: X \rightarrow X$  of a compact Kähler manifold  $X$ .
- An endofunctor  $F: \mathcal{D} \rightarrow \mathcal{D}$  of a triangulated category  $\mathcal{D}$ .

**Goal of this talk:** Explain reasons that “categorical dynamical systems” could be interesting.

A (discrete) **dynamical system** is a pair  $(X, \phi)$  where

- $\phi: X \rightarrow X$  preserves certain mathematical structures on  $X$ .

We would like to study the long-term behavior of  $\phi^n$  under large iterations.

Examples:

- A linear self-map  $T: V \rightarrow V$  of a vector space  $V$ .
- A continuous self-map  $f: X \rightarrow X$  of a compact metric space  $X$ .
- A holomorphic self-map  $f: X \rightarrow X$  of a compact Kähler manifold  $X$ .
- An endofunctor  $F: \mathcal{D} \rightarrow \mathcal{D}$  of a triangulated category  $\mathcal{D}$ .

**Goal of this talk:** Explain reasons that “categorical dynamical systems” could be interesting.

A (discrete) **dynamical system** is a pair  $(X, \phi)$  where

- $\phi: X \rightarrow X$  preserves certain mathematical structures on  $X$ .

We would like to study the long-term behavior of  $\phi^n$  under large iterations.

Examples:

- A linear self-map  $T: V \rightarrow V$  of a vector space  $V$ .
- A continuous self-map  $f: X \rightarrow X$  of a compact metric space  $X$ .
- A holomorphic self-map  $f: X \rightarrow X$  of a compact Kähler manifold  $X$ .
- An endofunctor  $F: \mathcal{D} \rightarrow \mathcal{D}$  of a triangulated category  $\mathcal{D}$ .

**Goal of this talk:** Explain reasons that “categorical dynamical systems” could be interesting.

A (discrete) **dynamical system** is a pair  $(X, \phi)$  where

- $\phi: X \rightarrow X$  preserves certain mathematical structures on  $X$ .

We would like to study the long-term behavior of  $\phi^n$  under large iterations.

Examples:

- A linear self-map  $T: V \rightarrow V$  of a vector space  $V$ .
- A continuous self-map  $f: X \rightarrow X$  of a compact metric space  $X$ .
- A holomorphic self-map  $f: X \rightarrow X$  of a compact Kähler manifold  $X$ .
- An endofunctor  $F: \mathcal{D} \rightarrow \mathcal{D}$  of a triangulated category  $\mathcal{D}$ .

**Goal of this talk:** Explain reasons that “categorical dynamical systems” could be interesting.

A (discrete) **dynamical system** is a pair  $(X, \phi)$  where

- $\phi: X \rightarrow X$  preserves certain mathematical structures on  $X$ .

We would like to study the long-term behavior of  $\phi^n$  under large iterations.

Examples:

- A linear self-map  $T: V \rightarrow V$  of a vector space  $V$ .
- A continuous self-map  $f: X \rightarrow X$  of a compact metric space  $X$ .
- A holomorphic self-map  $f: X \rightarrow X$  of a compact Kähler manifold  $X$ .
- An endofunctor  $F: \mathcal{D} \rightarrow \mathcal{D}$  of a triangulated category  $\mathcal{D}$ .



**Goal of this talk:** Explain reasons that “categorical dynamical systems” could be interesting.

A (discrete) **dynamical system** is a pair  $(X, \phi)$  where

- $\phi: X \rightarrow X$  preserves certain mathematical structures on  $X$ .

We would like to study the long-term behavior of  $\phi^n$  under large iterations.

Examples:

- A linear self-map  $T: V \rightarrow V$  of a vector space  $V$ .
- A continuous self-map  $f: X \rightarrow X$  of a compact metric space  $X$ .
- A holomorphic self-map  $f: X \rightarrow X$  of a compact Kähler manifold  $X$ .
- An endofunctor  $F: \mathcal{D} \rightarrow \mathcal{D}$  of a triangulated category  $\mathcal{D}$ .

# Examples of triangulated categories

Recall that a triangulated category is an additive category with a shift functor  $[1]$  and a collection of exact triangles

$$\cdots \rightarrow A \rightarrow B \rightarrow C \rightarrow A[1] \rightarrow \cdots$$

that satisfy a set of axioms.

(Analogy: Exact sequences  $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$  in abelian categories.)  
( $C$  is the “mapping cone” of  $A \rightarrow B$ .)

Examples:

- $\mathcal{D}^b\mathrm{Coh}(X)$ , where  $X$  is a smooth complex projective variety  
(objects: (complex of) holomorphic vector bundles on  $X$ )
- $\mathcal{D}^\pi\mathrm{Fuk}(Y)$ , where  $Y$  is a symplectic manifold  
(objects: Lagrangian submanifolds in  $Y$ , morphisms:  $L_1 \cap L_2$ )

# Examples of triangulated categories

Recall that a triangulated category is an additive category with a shift functor  $[1]$  and a collection of exact triangles

$$\cdots \rightarrow A \rightarrow B \rightarrow C \rightarrow A[1] \rightarrow \cdots$$

that satisfy a set of axioms.

(Analogy: Exact sequences  $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$  in abelian categories.)  
( $C$  is the “mapping cone” of  $A \rightarrow B$ .)

Examples:

- $\mathcal{D}^b\mathrm{Coh}(X)$ , where  $X$  is a smooth complex projective variety  
(objects: (complex of) holomorphic vector bundles on  $X$ )
- $\mathcal{D}^\pi\mathrm{Fuk}(Y)$ , where  $Y$  is a symplectic manifold  
(objects: Lagrangian submanifolds in  $Y$ , morphisms:  $L_1 \cap L_2$ )

# Examples of triangulated categories

Recall that a triangulated category is an additive category with a shift functor  $[1]$  and a collection of exact triangles

$$\cdots \rightarrow A \rightarrow B \rightarrow C \rightarrow A[1] \rightarrow \cdots$$

that satisfy a set of axioms.

(Analogy: Exact sequences  $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$  in abelian categories.)  
( $C$  is the “mapping cone” of  $A \rightarrow B$ .)

Examples:

- $\mathcal{D}^b\mathrm{Coh}(X)$ , where  $X$  is a smooth complex projective variety  
(objects: (complex of) holomorphic vector bundles on  $X$ )
- $\mathcal{D}^\pi\mathrm{Fuk}(Y)$ , where  $Y$  is a symplectic manifold  
(objects: Lagrangian submanifolds in  $Y$ , morphisms:  $L_1 \cap L_2$ )

# Examples of triangulated categories

Recall that a triangulated category is an additive category with a shift functor  $[1]$  and a collection of exact triangles

$$\cdots \rightarrow A \rightarrow B \rightarrow C \rightarrow A[1] \rightarrow \cdots$$

that satisfy a set of axioms.

(Analogy: Exact sequences  $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$  in abelian categories.)  
( $C$  is the “mapping cone” of  $A \rightarrow B$ .)

Examples:

- $\mathcal{D}^b\mathrm{Coh}(X)$ , where  $X$  is a smooth complex projective variety  
(objects: (complex of) holomorphic vector bundles on  $X$ )
- $\mathcal{D}^\pi\mathrm{Fuk}(Y)$ , where  $Y$  is a symplectic manifold  
(objects: Lagrangian submanifolds in  $Y$ , morphisms:  $L_1 \cap L_2$ )

# Why categorical dynamical systems? (I)

Both **holomorphic dynamics** and **symplectic dynamics** can be discussed in the categorical settings.

- A holomorphic self-map  $f: X \rightarrow X$  induces an endofunctor

$$\mathbb{L}f^*: \mathcal{D}^b\mathrm{Coh}(X) \rightarrow \mathcal{D}^b\mathrm{Coh}(X).$$

- A symplectomorphism  $f: Y \rightarrow Y$  induces an autoequivalence

$$f_*: \mathcal{D}^\pi\mathrm{Fuk}(Y) \rightarrow \mathcal{D}^\pi\mathrm{Fuk}(Y).$$

Moreover, with homological mirror symmetry conjecture, one can consider “mixings” of holomorphic and symplectic dynamical systems on

$$\mathcal{D}^b\mathrm{Coh}(X) \cong \mathcal{D}^\pi\mathrm{Fuk}(Y).$$

# Why categorical dynamical systems? (I)

Both **holomorphic dynamics** and **symplectic dynamics** can be discussed in the categorical settings.

- A holomorphic self-map  $f: X \rightarrow X$  induces an endofunctor

$$\mathbb{L}f^*: \mathcal{D}^b\mathrm{Coh}(X) \rightarrow \mathcal{D}^b\mathrm{Coh}(X).$$

- A symplectomorphism  $f: Y \rightarrow Y$  induces an autoequivalence

$$f_*: \mathcal{D}^\pi\mathrm{Fuk}(Y) \rightarrow \mathcal{D}^\pi\mathrm{Fuk}(Y).$$

Moreover, with homological mirror symmetry conjecture, one can consider “mixings” of holomorphic and symplectic dynamical systems on

$$\mathcal{D}^b\mathrm{Coh}(X) \cong \mathcal{D}^\pi\mathrm{Fuk}(Y).$$

# Why categorical dynamical systems? (I)

Both **holomorphic dynamics** and **symplectic dynamics** can be discussed in the categorical settings.

- A holomorphic self-map  $f: X \rightarrow X$  induces an endofunctor

$$\mathbb{L}f^*: \mathcal{D}^b\mathrm{Coh}(X) \rightarrow \mathcal{D}^b\mathrm{Coh}(X).$$

- A symplectomorphism  $f: Y \rightarrow Y$  induces an autoequivalence

$$f_*: \mathcal{D}^\pi\mathrm{Fuk}(Y) \rightarrow \mathcal{D}^\pi\mathrm{Fuk}(Y).$$

Moreover, with homological mirror symmetry conjecture, one can consider “mixings” of holomorphic and symplectic dynamical systems on

$$\mathcal{D}^b\mathrm{Coh}(X) \cong \mathcal{D}^\pi\mathrm{Fuk}(Y).$$



# Why categorical dynamical systems? (I)

Both **holomorphic dynamics** and **symplectic dynamics** can be discussed in the categorical settings.

- A holomorphic self-map  $f: X \rightarrow X$  induces an endofunctor

$$\mathbb{L}f^*: \mathcal{D}^b\mathrm{Coh}(X) \rightarrow \mathcal{D}^b\mathrm{Coh}(X).$$

- A symplectomorphism  $f: Y \rightarrow Y$  induces an autoequivalence

$$f_*: \mathcal{D}^\pi\mathrm{Fuk}(Y) \rightarrow \mathcal{D}^\pi\mathrm{Fuk}(Y).$$

Moreover, with homological mirror symmetry conjecture, one can consider “mixings” of holomorphic and symplectic dynamical systems on

$$\mathcal{D}^b\mathrm{Coh}(X) \cong \mathcal{D}^\pi\mathrm{Fuk}(Y).$$

## Why categorical dynamical systems? (II)

There is a parallel between **Teichmüller theory** and the theory of **stability conditions on triangulated categories**, developed by Bridgeland, Smith, Dimitrov, Haiden, Katzarkov, Kontsevich, etc.

Riemann surfaces	Triangulated categories
curve $C$	object $E$
$C_1 \cap C_2$	$\mathrm{Hom}(E_1, E_2)$
metric $g$	stability condition $\sigma$
geodesic	stable objects
length $\ell_g(C)$	mass $m_\sigma(E)$
slope of $C$	phase $\phi_\sigma(E)$
diffeomorphisms	autoequivalences
...	...

One can therefore consider categorical analogues of results in **Teichmüller dynamics**.

# Why categorical dynamical systems? (II)

There is a parallel between **Teichmüller theory** and the theory of **stability conditions on triangulated categories**, developed by Bridgeland, Smith, Dimitrov, Haiden, Katzarkov, Kontsevich, etc.

Riemann surfaces	Triangulated categories
curve $C$	object $E$
$C_1 \cap C_2$	$\mathrm{Hom}(E_1, E_2)$
metric $g$	stability condition $\sigma$
geodesic	stable objects
length $\ell_g(C)$	mass $m_\sigma(E)$
slope of $C$	phase $\phi_\sigma(E)$
diffeomorphisms	autoequivalences
...	...

One can therefore consider categorical analogues of results in **Teichmüller dynamics**.

## Why categorical dynamical systems? (II)

There is a parallel between **Teichmüller theory** and the theory of **stability conditions on triangulated categories**, developed by Bridgeland, Smith, Dimitrov, Haiden, Katzarkov, Kontsevich, etc.

Riemann surfaces	Triangulated categories
curve $C$	object $E$
$C_1 \cap C_2$	$\mathrm{Hom}(E_1, E_2)$
metric $g$	stability condition $\sigma$
geodesic	stable objects
length $\ell_g(C)$	mass $m_\sigma(E)$
slope of $C$	phase $\phi_\sigma(E)$
diffeomorphisms	autoequivalences
...	...

One can therefore consider categorical analogues of results in **Teichmüller dynamics**.

# Why categorical dynamical systems? (III)

Analogy with **symbolic dynamics**:

Subshifts	Triangulated categories
finite alphabet $\mathcal{A}$	bounded $t$ -structure
shift-invariant subset $X \subseteq \mathcal{A}^{\mathbb{Z}}$	triangulated categories
shift map $\sigma: X \rightarrow X$	shift functor $[1]$
automorphism $\text{Aut}(X, \sigma)$	autoequivalences

One can also consider categorical analogues of results in symbolic dynamics.

# Why categorical dynamical systems? (III)

Analogy with **symbolic dynamics**:

Subshifts	Triangulated categories
finite alphabets $\mathcal{A}$	bounded $t$ -structure
shift-invariant subset $X \subseteq \mathcal{A}^{\mathbb{Z}}$	triangulated categories
shift map $\sigma: X \rightarrow X$	shift functor $[1]$
automorphism $\text{Aut}(X, \sigma)$	autoequivalences

One can also consider categorical analogues of results in symbolic dynamics.

# Why categorical dynamical systems? (III)

Analogy with [symbolic dynamics](#):

Subshifts	Triangulated categories
finite alphabet $\mathcal{A}$	bounded $t$ -structure
shift-invariant subset $X \subseteq \mathcal{A}^{\mathbb{Z}}$	triangulated categories
shift map $\sigma: X \rightarrow X$	shift functor $[1]$
automorphism $\text{Aut}(X, \sigma)$	autoequivalences

One can also consider categorical analogues of results in symbolic dynamics.

# Outline

- (I). Entropy of holomorphic and symplectic dynamical systems, and mixings of them.
- (II). Finite subgroups of  $\text{Aut}(\mathcal{D})$  acting on  $\text{Stab}(\mathcal{D})$ .
- (III). Shifting numbers and quasimorphisms on  $\text{Aut}(\mathcal{D})$ .



# Topological entropy...

... is hard to compute in general.

Let  $(X, d)$  compact metric space and  $f: X \rightarrow X$  continuous. Consider

$$N(n, \epsilon) := \max \left\{ \ell: \exists x_1, \dots, x_\ell \text{ s.t. } \max_{0 \leq k \leq n} \{d(f^k(x_i), f^k(x_j))\} \geq \epsilon \ \forall x_i, x_j \right\}$$

The **topological entropy** of  $f$  is defined to be

$$h_{\text{top}}(f) := \lim_{\epsilon \rightarrow 0} \left( \limsup_{n \rightarrow \infty} \frac{1}{n} \log N(n, \epsilon) \right) \in [0, \infty].$$

Basic properties:

- It's a topological invariant measuring the “complexity” of  $f$ .
- $f^n = \text{id}_X \implies h_{\text{top}}(f) = 0$ .

# Topological entropy...

... is hard to compute in general.

Let  $(X, d)$  compact metric space and  $f: X \rightarrow X$  continuous. Consider

$$N(n, \epsilon) := \max \left\{ \ell: \exists x_1, \dots, x_\ell \text{ s.t. } \max_{0 \leq k \leq n} \{d(f^k(x_i), f^k(x_j))\} \geq \epsilon \ \forall x_i, x_j \right\}$$

The **topological entropy** of  $f$  is defined to be

$$h_{\text{top}}(f) := \lim_{\epsilon \rightarrow 0} \left( \limsup_{n \rightarrow \infty} \frac{1}{n} \log N(n, \epsilon) \right) \in [0, \infty].$$

Basic properties:

- It's a topological invariant measuring the “complexity” of  $f$ .
- $f^n = \text{id}_X \implies h_{\text{top}}(f) = 0$ .

# Topological entropy...

... is hard to compute in general.

Let  $(X, d)$  compact metric space and  $f: X \rightarrow X$  continuous. Consider

$$N(n, \epsilon) := \max \left\{ \ell: \exists x_1, \dots, x_\ell \text{ s.t. } \max_{0 \leq k \leq n} \{d(f^k(x_i), f^k(x_j))\} \geq \epsilon \ \forall x_i, x_j \right\}$$

The **topological entropy** of  $f$  is defined to be

$$h_{\text{top}}(f) := \lim_{\epsilon \rightarrow 0} \left( \limsup_{n \rightarrow \infty} \frac{1}{n} \log N(n, \epsilon) \right) \in [0, \infty].$$

Basic properties:

- It's a topological invariant measuring the “complexity” of  $f$ .
- $f^n = \text{id}_X \implies h_{\text{top}}(f) = 0$ .

# Topological entropy...

... is hard to compute in general.

Let  $(X, d)$  compact metric space and  $f: X \rightarrow X$  continuous. Consider

$$N(n, \epsilon) := \max \left\{ \ell: \exists x_1, \dots, x_\ell \text{ s.t. } \max_{0 \leq k \leq n} \{d(f^k(x_i), f^k(x_j))\} \geq \epsilon \ \forall x_i, x_j \right\}$$

The **topological entropy** of  $f$  is defined to be

$$h_{\text{top}}(f) := \lim_{\epsilon \rightarrow 0} \left( \limsup_{n \rightarrow \infty} \frac{1}{n} \log N(n, \epsilon) \right) \in [0, \infty].$$

Basic properties:

- It's a topological invariant measuring the “complexity” of  $f$ .
- $f^n = \text{id}_X \implies h_{\text{top}}(f) = 0$ .

## Topological entropy...

... is hard to compute in general.

Let  $(X, d)$  compact metric space and  $f: X \rightarrow X$  continuous. Consider

$$N(n, \epsilon) := \max \left\{ \ell: \exists x_1, \dots, x_\ell \text{ s.t. } \max_{0 \leq k \leq n} \{d(f^k(x_i), f^k(x_j))\} \geq \epsilon \ \forall x_i, x_j \right\}$$

The **topological entropy** of  $f$  is defined to be

$$h_{\text{top}}(f) := \lim_{\epsilon \rightarrow 0} \left( \limsup_{n \rightarrow \infty} \frac{1}{n} \log N(n, \epsilon) \right) \in [0, \infty].$$

Basic properties:

- It's a topological invariant measuring the “complexity” of  $f$ .
- $f^n = \text{id}_X \implies h_{\text{top}}(f) = 0$ .

## Example: Holomorphic maps on compact Kähler manifolds

One of the most fundamental results in (higher dimensional) complex dynamics is the following result of Gromov and Yomdin.

### Theorem (Gromov, Yomdin)

*If  $f: X \rightarrow X$  is a surjective holomorphic map of a compact Kähler manifold, then*

$$h_{\text{top}}(f) = \log \rho(f^*)$$

*where  $\rho$  is the spectral radius of  $f^*: H^*(X, \mathbb{C}) \rightarrow H^*(X, \mathbb{C})$ .*

Here is a geometric application of the topological entropy.

### Theorem (Cantat)

*If a compact complex surface  $X$  admits an automorphism of positive topological entropy, then  $X$  is either a torus, a K3 surface, an Enriques surface, or a rational surface.*

## Example: Holomorphic maps on compact Kähler manifolds

One of the most fundamental results in (higher dimensional) complex dynamics is the following result of Gromov and Yomdin.

### Theorem (Gromov, Yomdin)

*If  $f: X \rightarrow X$  is a surjective holomorphic map of a compact Kähler manifold, then*

$$h_{\text{top}}(f) = \log \rho(f^*)$$

*where  $\rho$  is the spectral radius of  $f^*: H^*(X, \mathbb{C}) \rightarrow H^*(X, \mathbb{C})$ .*

Here is a geometric application of the topological entropy.

### Theorem (Cantat)

*If a compact complex surface  $X$  admits an automorphism of positive topological entropy, then  $X$  is either a torus, a K3 surface, an Enriques surface, or a rational surface.*

## Example: Holomorphic maps on compact Kähler manifolds

One of the most fundamental results in (higher dimensional) complex dynamics is the following result of Gromov and Yomdin.

### Theorem (Gromov, Yomdin)

*If  $f: X \rightarrow X$  is a surjective holomorphic map of a compact Kähler manifold, then*

$$h_{\text{top}}(f) = \log \rho(f^*)$$

*where  $\rho$  is the spectral radius of  $f^*: H^*(X, \mathbb{C}) \rightarrow H^*(X, \mathbb{C})$ .*

Here is a geometric application of the topological entropy.

### Theorem (Cantat)

*If a compact complex surface  $X$  admits an automorphism of positive topological entropy, then  $X$  is either a torus, a K3 surface, an Enriques surface, or a rational surface.*



# Categorical entropy

Let  $F: \mathcal{D} \rightarrow \mathcal{D}$  be as before, and  $G, G' \in \mathcal{D}$  be split generators.

Dimitrov, Haiden, Katzarkov, and Kontsevich defined:

$$h_{\text{cat}}(F) := \lim_{n \rightarrow \infty} \frac{1}{n} \log \left( \sum_{k \in \mathbb{Z}} \dim \operatorname{Hom}(G, F^n G'[k]) \right).$$

Basic properties:

- The limit exists, and is independent of the choice of  $G, G'$ .
- $F^n = [m] \implies h_{\text{cat}}(F) = 0$ .

Example: When  $\mathcal{D} = \mathcal{D}^b \operatorname{Coh}(X)$ :

- Kikuta–Takahashi:  $h_{\text{cat}}(\mathbb{L}f^*) = h_{\text{top}}(f) = \log \rho(f^*) = \log \rho([\mathbb{L}f^*]_{H^*})$ .
- $h_{\text{cat}}(- \otimes L) = 0$ .

However, there exist a Calabi–Yau manifold  $X$  and  $F = T \circ (- \otimes L)$  with  $h_{\text{cat}}(F) > \log \rho([F])$ , where  $T$  is mirror to certain Dehn twist.

# Categorical entropy

Let  $F: \mathcal{D} \rightarrow \mathcal{D}$  be as before, and  $G, G' \in \mathcal{D}$  be split generators. Dimitrov, Haiden, Katzarkov, and Kontsevich defined:

$$h_{\text{cat}}(F) := \lim_{n \rightarrow \infty} \frac{1}{n} \log \left( \sum_{k \in \mathbb{Z}} \dim \operatorname{Hom}(G, F^n G'[k]) \right).$$

Basic properties:

- The limit exists, and is independent of the choice of  $G, G'$ .
- $F^n = [m] \implies h_{\text{cat}}(F) = 0$ .

Example: When  $\mathcal{D} = \mathcal{D}^b \operatorname{Coh}(X)$ :

- Kikuta–Takahashi:  $h_{\text{cat}}(\mathbb{L}f^*) = h_{\text{top}}(f) = \log \rho(f^*) = \log \rho([\mathbb{L}f^*]_{H^*})$ .
- $h_{\text{cat}}(- \otimes L) = 0$ .

However, there exist a Calabi–Yau manifold  $X$  and  $F = T \circ (- \otimes L)$  with  $h_{\text{cat}}(F) > \log \rho([F])$ , where  $T$  is mirror to certain Dehn twist.

# Categorical entropy

Let  $F: \mathcal{D} \rightarrow \mathcal{D}$  be as before, and  $G, G' \in \mathcal{D}$  be split generators. Dimitrov, Haiden, Katzarkov, and Kontsevich defined:

$$h_{\text{cat}}(F) := \lim_{n \rightarrow \infty} \frac{1}{n} \log \left( \sum_{k \in \mathbb{Z}} \dim \operatorname{Hom}(G, F^n G'[k]) \right).$$

Basic properties:

- The limit exists, and is independent of the choice of  $G, G'$ .
- $F^n = [m] \implies h_{\text{cat}}(F) = 0$ .

Example: When  $\mathcal{D} = \mathcal{D}^b \operatorname{Coh}(X)$ :

- Kikuta–Takahashi:  $h_{\text{cat}}(\mathbb{L}f^*) = h_{\text{top}}(f) = \log \rho(f^*) = \log \rho([\mathbb{L}f^*]_{H^*})$ .
- $h_{\text{cat}}(- \otimes L) = 0$ .

However, there exist a Calabi–Yau manifold  $X$  and  $F = T \circ (- \otimes L)$  with  $h_{\text{cat}}(F) > \log \rho([F])$ , where  $T$  is mirror to certain Dehn twist.

## Categorical entropy

Let  $F: \mathcal{D} \rightarrow \mathcal{D}$  be as before, and  $G, G' \in \mathcal{D}$  be split generators. Dimitrov, Haiden, Katzarkov, and Kontsevich defined:

$$h_{\text{cat}}(F) := \lim_{n \rightarrow \infty} \frac{1}{n} \log \left( \sum_{k \in \mathbb{Z}} \dim \operatorname{Hom}(G, F^n G'[k]) \right).$$

Basic properties:

- The limit exists, and is independent of the choice of  $G, G'$ .
- $F^n = [m] \implies h_{\text{cat}}(F) = 0$ .

Example: When  $\mathcal{D} = \mathcal{D}^b \operatorname{Coh}(X)$ :

- Kikuta–Takahashi:  $h_{\text{cat}}(\mathbb{L}f^*) = h_{\text{top}}(f) = \log \rho(f^*) = \log \rho([\mathbb{L}f^*]_{H^*})$ .
- $h_{\text{cat}}(- \otimes L) = 0$ .

However, there exist a Calabi–Yau manifold  $X$  and  $F = T \circ (- \otimes L)$  with  $h_{\text{cat}}(F) > \log \rho([F])$ , where  $T$  is mirror to certain Dehn twist.

## Categorical entropy

Let  $F: \mathcal{D} \rightarrow \mathcal{D}$  be as before, and  $G, G' \in \mathcal{D}$  be split generators. Dimitrov, Haiden, Katzarkov, and Kontsevich defined:

$$h_{\text{cat}}(F) := \lim_{n \rightarrow \infty} \frac{1}{n} \log \left( \sum_{k \in \mathbb{Z}} \dim \operatorname{Hom}(G, F^n G'[k]) \right).$$

Basic properties:

- The limit exists, and is independent of the choice of  $G, G'$ .
- $F^n = [m] \implies h_{\text{cat}}(F) = 0$ .

Example: When  $\mathcal{D} = \mathcal{D}^b \operatorname{Coh}(X)$ :

- Kikuta–Takahashi:  $h_{\text{cat}}(\mathbb{L}f^*) = h_{\text{top}}(f) = \log \rho(f^*) = \log \rho([\mathbb{L}f^*]_{H^*})$ .
- $h_{\text{cat}}(- \otimes L) = 0$ .

However, there exist a Calabi–Yau manifold  $X$  and  $F = T \circ (- \otimes L)$  with  $h_{\text{cat}}(F) > \log \rho([F])$ , where  $T$  is mirror to certain Dehn twist.

## Categorical entropy

Let  $F: \mathcal{D} \rightarrow \mathcal{D}$  be as before, and  $G, G' \in \mathcal{D}$  be split generators. Dimitrov, Haiden, Katzarkov, and Kontsevich defined:

$$h_{\text{cat}}(F) := \lim_{n \rightarrow \infty} \frac{1}{n} \log \left( \sum_{k \in \mathbb{Z}} \dim \operatorname{Hom}(G, F^n G'[k]) \right).$$

Basic properties:

- The limit exists, and is independent of the choice of  $G, G'$ .
- $F^n = [m] \implies h_{\text{cat}}(F) = 0$ .

Example: When  $\mathcal{D} = \mathcal{D}^b \operatorname{Coh}(X)$ :

- Kikuta–Takahashi:  $h_{\text{cat}}(\mathbb{L}f^*) = h_{\text{top}}(f) = \log \rho(f^*) = \log \rho([\mathbb{L}f^*]_{H^*})$ .
- $h_{\text{cat}}(- \otimes L) = 0$ .

However, there exist a Calabi–Yau manifold  $X$  and  $F = T \circ (- \otimes L)$  with  $h_{\text{cat}}(F) > \log \rho([F])$ , where  $T$  is mirror to certain Dehn twist.

# Nielsen–Thurston classification

- $\Sigma$ : Riemann surface
- $\text{MCG}(\Sigma) = \text{Diff}(\Sigma)/\text{isotopy}$ : mapping class group
- each mapping class is either:
  - ▶ finite order
  - ▶ reducible
  - ▶ pseudo-Anosov

For instance –

- elements of  $\text{MCG}(T^2) = \text{SL}(2, \mathbb{Z})$  are either:
  - ▶ elliptic (finite order)
  - ▶ parabolic (Dehn twist)
  - ▶ hyperbolic (two transverse foliations: stretch one by  $\lambda > 1$ , compress the other by  $1/\lambda$ )

# Nielsen–Thurston classification

- $\Sigma$ : Riemann surface
- $\text{MCG}(\Sigma) = \text{Diff}(\Sigma)/\text{isotopy}$ : mapping class group
- each mapping class is either:
  - ▶ finite order
  - ▶ reducible
  - ▶ pseudo-Anosov

For instance –

- elements of  $\text{MCG}(T^2) = \text{SL}(2, \mathbb{Z})$  are either:
  - ▶ elliptic (finite order)
  - ▶ parabolic (Dehn twist)
  - ▶ hyperbolic (two transverse foliations: stretch one by  $\lambda > 1$ , compress the other by  $1/\lambda$ )



# Nielsen–Thurston classification

- $\Sigma$ : Riemann surface
- $\text{MCG}(\Sigma) = \text{Diff}(\Sigma)/\text{isotopy}$ : mapping class group
- each mapping class is either:
  - ▶ finite order
  - ▶ reducible
  - ▶ pseudo-Anosov

For instance –

- elements of  $\text{MCG}(T^2) = \text{SL}(2, \mathbb{Z})$  are either:
  - ▶ elliptic (finite order)
  - ▶ parabolic (Dehn twist)
  - ▶ hyperbolic (two transverse foliations: stretch one by  $\lambda > 1$ , compress the other by  $1/\lambda$ )

# Nielsen–Thurston classification

- $\Sigma$ : Riemann surface
- $\text{MCG}(\Sigma) = \text{Diff}(\Sigma)/\text{isotopy}$ : mapping class group
- each mapping class is either:
  - ▶ finite order
  - ▶ reducible
  - ▶ pseudo-Anosov

For instance –

- elements of  $\text{MCG}(T^2) = \text{SL}(2, \mathbb{Z})$  are either:
  - ▶ elliptic (finite order)
  - ▶ parabolic (Dehn twist)
  - ▶ hyperbolic (two transverse foliations: stretch one by  $\lambda > 1$ , compress the other by  $1/\lambda$ )

# Nielsen–Thurston classification

- $\Sigma$ : Riemann surface
- $\text{MCG}(\Sigma) = \text{Diff}(\Sigma)/\text{isotopy}$ : mapping class group
- each mapping class is either:
  - ▶ finite order
  - ▶ reducible
  - ▶ pseudo-Anosov

For instance –

- elements of  $\text{MCG}(T^2) = \text{SL}(2, \mathbb{Z})$  are either:
  - ▶ elliptic (finite order)
  - ▶ parabolic (Dehn twist)
  - ▶ hyperbolic (two transverse foliations: stretch one by  $\lambda > 1$ , compress the other by  $1/\lambda$ )

# Realization problem of finite subgroups

- Nielsen asked (1923): Let  $G \subseteq \mathrm{MCG}(\Sigma)$  be a finite subgroup. Does there always exist a lifting  $G \subseteq \mathrm{Diff}(\Sigma)$ ?
- Kerckhoff (1983): Yes! Moreover, there exists a metric  $g$  such that  $G \subseteq \mathrm{Isom}(\Sigma, g)$ . Or equivalently,  $G$  fixes a point in  $\mathrm{Teich}(\Sigma)$ . (e.g.  $\mathrm{MCG}(\mathcal{T}^2) = \mathrm{SL}(2, \mathbb{Z})$  acts on  $\mathrm{Teich}(\mathcal{T}^2) = \mathbb{H}$ .)
- Farb–Looijenga (2021) also proved similar statements for K3 surfaces (under certain conditions), where  $g$  is replaced by complex structure or Ricci-flat metric on the K3 surface.
- F.–Lai (2023): Let  $\mathcal{D} = \mathcal{D}^b(X)$  be the derived category of a general K3 surface  $X$ . Then any finite subgroup  $G \subseteq \mathrm{Aut}(\mathcal{D})$  fixes a point in  $\mathrm{Stab}(\mathcal{D})$ . Using this, we provide a full classification of finite subgroups of  $\mathrm{Aut}(\mathcal{D})$ .

# Realization problem of finite subgroups

- Nielsen asked (1923): Let  $G \subseteq \mathrm{MCG}(\Sigma)$  be a finite subgroup. Does there always exist a lifting  $G \subseteq \mathrm{Diff}(\Sigma)$ ?
- Kerckhoff (1983): Yes! Moreover, there exists a metric  $g$  such that  $G \subseteq \mathrm{Isom}(\Sigma, g)$ . Or equivalently,  $G$  fixes a point in  $\mathrm{Teich}(\Sigma)$ . (e.g.  $\mathrm{MCG}(\mathcal{T}^2) = \mathrm{SL}(2, \mathbb{Z})$  acts on  $\mathrm{Teich}(\mathcal{T}^2) = \mathbb{H}$ .)
- Farb–Looijenga (2021) also proved similar statements for K3 surfaces (under certain conditions), where  $g$  is replaced by complex structure or Ricci-flat metric on the K3 surface.
- F.–Lai (2023): Let  $\mathcal{D} = \mathcal{D}^b(X)$  be the derived category of a general K3 surface  $X$ . Then any finite subgroup  $G \subseteq \mathrm{Aut}(\mathcal{D})$  fixes a point in  $\mathrm{Stab}(\mathcal{D})$ . Using this, we provide a full classification of finite subgroups of  $\mathrm{Aut}(\mathcal{D})$ .

# Realization problem of finite subgroups

- Nielsen asked (1923): Let  $G \subseteq \mathrm{MCG}(\Sigma)$  be a finite subgroup. Does there always exist a lifting  $G \subseteq \mathrm{Diff}(\Sigma)$ ?
- Kerckhoff (1983): Yes! Moreover, there exists a metric  $g$  such that  $G \subseteq \mathrm{Isom}(\Sigma, g)$ . Or equivalently,  $G$  fixes a point in  $\mathrm{Teich}(\Sigma)$ .  
(e.g.  $\mathrm{MCG}(\mathcal{T}^2) = \mathrm{SL}(2, \mathbb{Z})$  acts on  $\mathrm{Teich}(\mathcal{T}^2) = \mathbb{H}$ .)
- Farb–Looijenga (2021) also proved similar statements for K3 surfaces (under certain conditions), where  $g$  is replaced by complex structure or Ricci-flat metric on the K3 surface.
- F.–Lai (2023): Let  $\mathcal{D} = \mathcal{D}^b(X)$  be the derived category of a general K3 surface  $X$ . Then any finite subgroup  $G \subseteq \mathrm{Aut}(\mathcal{D})$  fixes a point in  $\mathrm{Stab}(\mathcal{D})$ . Using this, we provide a full classification of finite subgroups of  $\mathrm{Aut}(\mathcal{D})$ .

# Realization problem of finite subgroups

- Nielsen asked (1923): Let  $G \subseteq \mathrm{MCG}(\Sigma)$  be a finite subgroup. Does there always exist a lifting  $G \subseteq \mathrm{Diff}(\Sigma)$ ?
- Kerckhoff (1983): Yes! Moreover, there exists a metric  $g$  such that  $G \subseteq \mathrm{Isom}(\Sigma, g)$ . Or equivalently,  $G$  fixes a point in  $\mathrm{Teich}(\Sigma)$ . (e.g.  $\mathrm{MCG}(T^2) = \mathrm{SL}(2, \mathbb{Z})$  acts on  $\mathrm{Teich}(T^2) = \mathbb{H}$ .)
- Farb–Looijenga (2021) also proved similar statements for K3 surfaces (under certain conditions), where  $g$  is replaced by complex structure or Ricci-flat metric on the K3 surface.
- F.–Lai (2023): Let  $\mathcal{D} = \mathcal{D}^b(X)$  be the derived category of a general K3 surface  $X$ . Then any finite subgroup  $G \subseteq \mathrm{Aut}(\mathcal{D})$  fixes a point in  $\mathrm{Stab}(\mathcal{D})$ . Using this, we provide a full classification of finite subgroups of  $\mathrm{Aut}(\mathcal{D})$ .

# Realization problem of finite subgroups

- Nielsen asked (1923): Let  $G \subseteq \mathrm{MCG}(\Sigma)$  be a finite subgroup. Does there always exist a lifting  $G \subseteq \mathrm{Diff}(\Sigma)$ ?
- Kerckhoff (1983): Yes! Moreover, there exists a metric  $g$  such that  $G \subseteq \mathrm{Isom}(\Sigma, g)$ . Or equivalently,  $G$  fixes a point in  $\mathrm{Teich}(\Sigma)$ . (e.g.  $\mathrm{MCG}(T^2) = \mathrm{SL}(2, \mathbb{Z})$  acts on  $\mathrm{Teich}(T^2) = \mathbb{H}$ .)
- Farb–Looijenga (2021) also proved similar statements for K3 surfaces (under certain conditions), where  $g$  is replaced by complex structure or Ricci-flat metric on the K3 surface.
- F.–Lai (2023): Let  $\mathcal{D} = \mathcal{D}^b(X)$  be the derived category of a general K3 surface  $X$ . Then any finite subgroup  $G \subseteq \mathrm{Aut}(\mathcal{D})$  fixes a point in  $\mathrm{Stab}(\mathcal{D})$ . Using this, we provide a full classification of finite subgroups of  $\mathrm{Aut}(\mathcal{D})$ .



# Realization problem of finite subgroups

- Nielsen asked (1923): Let  $G \subseteq \mathrm{MCG}(\Sigma)$  be a finite subgroup. Does there always exist a lifting  $G \subseteq \mathrm{Diff}(\Sigma)$ ?
- Kerckhoff (1983): Yes! Moreover, there exists a metric  $g$  such that  $G \subseteq \mathrm{Isom}(\Sigma, g)$ . Or equivalently,  $G$  fixes a point in  $\mathrm{Teich}(\Sigma)$ . (e.g.  $\mathrm{MCG}(T^2) = \mathrm{SL}(2, \mathbb{Z})$  acts on  $\mathrm{Teich}(T^2) = \mathbb{H}$ .)
- Farb–Looijenga (2021) also proved similar statements for K3 surfaces (under certain conditions), where  $g$  is replaced by complex structure or Ricci-flat metric on the K3 surface.
- F.–Lai (2023): Let  $\mathcal{D} = \mathcal{D}^b(X)$  be the derived category of a general K3 surface  $X$ . Then any finite subgroup  $G \subseteq \mathrm{Aut}(\mathcal{D})$  fixes a point in  $\mathrm{Stab}(\mathcal{D})$ . Using this, we provide a full classification of finite subgroups of  $\mathrm{Aut}(\mathcal{D})$ .

# Poincaré translation numbers vs Shifting numbers

$$0 \rightarrow \mathbb{Z} \rightarrow \operatorname{Homeo}_{\mathbb{Z}}^+(\mathbb{R}) \rightarrow \operatorname{Homeo}^+(S^1) \rightarrow 1.$$

Poincaré translation number:  $\rho(f) := \lim_{n \rightarrow \infty} \frac{f^{(n)}(x_0) - x_0}{n}$ .

In the setting of triangulated categories, we have a central extension

$$0 \rightarrow \mathbb{Z} \rightarrow \operatorname{Aut}(\mathcal{D}) \rightarrow \operatorname{Aut}(\mathcal{D})/[1] \rightarrow 1.$$

Translation numbers	Shifting numbers
$f \in \operatorname{Homeo}_{\mathbb{Z}}^+(\mathbb{R})$	$F \in \operatorname{Aut}(\mathcal{D})$
$x_0 \in \mathbb{R}$	$G \in \mathcal{D}$
amount of translation	phases $\phi_{\sigma}^{\pm}: \operatorname{Ob}(\mathcal{D}) \rightarrow \mathbb{R}$
$f^{(n)}(x_0) - x_0$	$\phi_{\sigma}^{\pm}(F^n G) - \phi_{\sigma}^{\pm}(G)$
translation number	upper/lower shifting numbers

# Poincaré translation numbers vs Shifting numbers

$$0 \rightarrow \mathbb{Z} \rightarrow \operatorname{Homeo}_{\mathbb{Z}}^{+}(\mathbb{R}) \rightarrow \operatorname{Homeo}^{+}(S^1) \rightarrow 1.$$

Poincaré translation number:  $\rho(f) := \lim_{n \rightarrow \infty} \frac{f^{(n)}(x_0) - x_0}{n}.$

In the setting of triangulated categories, we have a central extension

$$0 \rightarrow \mathbb{Z} \rightarrow \operatorname{Aut}(\mathcal{D}) \rightarrow \operatorname{Aut}(\mathcal{D})/[1] \rightarrow 1.$$

Translation numbers	Shifting numbers
$f \in \operatorname{Homeo}_{\mathbb{Z}}^{+}(\mathbb{R})$	$F \in \operatorname{Aut}(\mathcal{D})$
$x_0 \in \mathbb{R}$	$G \in \mathcal{D}$
amount of translation	phases $\phi_{\sigma}^{\pm}: \operatorname{Ob}(\mathcal{D}) \rightarrow \mathbb{R}$
$f^{(n)}(x_0) - x_0$	$\phi_{\sigma}^{\pm}(F^n G) - \phi_{\sigma}^{\pm}(G)$
translation number	upper/lower shifting numbers

# Poincaré translation numbers vs Shifting numbers

$$0 \rightarrow \mathbb{Z} \rightarrow \operatorname{Homeo}_{\mathbb{Z}}^{+}(\mathbb{R}) \rightarrow \operatorname{Homeo}^{+}(S^1) \rightarrow 1.$$

Poincaré translation number:  $\rho(f) := \lim_{n \rightarrow \infty} \frac{f^{(n)}(x_0) - x_0}{n}$ .

In the setting of triangulated categories, we have a central extension

$$0 \rightarrow \mathbb{Z} \rightarrow \operatorname{Aut}(\mathcal{D}) \rightarrow \operatorname{Aut}(\mathcal{D})/[1] \rightarrow 1.$$

Translation numbers	Shifting numbers
$f \in \operatorname{Homeo}_{\mathbb{Z}}^{+}(\mathbb{R})$	$F \in \operatorname{Aut}(\mathcal{D})$
$x_0 \in \mathbb{R}$	$G \in \mathcal{D}$
amount of translation	phases $\phi_{\sigma}^{\pm}: \operatorname{Ob}(\mathcal{D}) \rightarrow \mathbb{R}$
$f^{(n)}(x_0) - x_0$	$\phi_{\sigma}^{\pm}(F^n G) - \phi_{\sigma}^{\pm}(G)$
translation number	upper/lower shifting numbers

# Poincaré translation numbers vs Shifting numbers

$$0 \rightarrow \mathbb{Z} \rightarrow \operatorname{Homeo}_{\mathbb{Z}}^+(\mathbb{R}) \rightarrow \operatorname{Homeo}^+(S^1) \rightarrow 1.$$

Poincaré translation number:  $\rho(f) := \lim_{n \rightarrow \infty} \frac{f^{(n)}(x_0) - x_0}{n}$ .

In the setting of triangulated categories, we have a central extension

$$0 \rightarrow \mathbb{Z} \rightarrow \operatorname{Aut}(\mathcal{D}) \rightarrow \operatorname{Aut}(\mathcal{D})/[1] \rightarrow 1.$$

Translation numbers	Shifting numbers
$f \in \operatorname{Homeo}_{\mathbb{Z}}^+(\mathbb{R})$	$F \in \operatorname{Aut}(\mathcal{D})$
$x_0 \in \mathbb{R}$	$G \in \mathcal{D}$
amount of translation	phases $\phi_{\sigma}^{\pm}: \operatorname{Ob}(\mathcal{D}) \rightarrow \mathbb{R}$
$f^{(n)}(x_0) - x_0$	$\phi_{\sigma}^{\pm}(F^n G) - \phi_{\sigma}^{\pm}(G)$
translation number	upper/lower shifting numbers

# Shifting numbers

## Theorem (F.–Filip, 2023)

- *The limit*

$$\tau^{\pm}(F) := \lim_{n \rightarrow \infty} \frac{\phi_{\sigma}^{\pm}(F^n G) - \phi_{\sigma}^{\pm}(G)}{n}$$

*always exists, and is independent of the choices of  $G$  and  $\sigma$ .*

- *The function*

$$h_t(F) := \lim_{n \rightarrow \infty} \frac{1}{n} \log \left( \sum_{k \in \mathbb{Z}} \dim \operatorname{Hom}(G, F^n G[k]) e^{-kt} \right),$$

*is a convex function in  $t$  satisfying:*

- $t \cdot \tau^+(F) \leq h_t(F) \leq h_0(F) + t \cdot \tau^+(F)$  for  $t \geq 0$ , and
- $t \cdot \tau^-(F) \leq h_t(F) \leq h_0(F) + t \cdot \tau^-(F)$  for  $t \leq 0$ .

## Theorem (F., 2023)

*Let  $X$  be an abelian variety. Then  $\tau = \tau^{\pm}: \operatorname{Aut}(\mathcal{D}^b(X)) \rightarrow \mathbb{R}$  is a quasimorphism.*

# Shifting numbers

## Theorem (F.–Filip, 2023)

- *The limit*

$$\tau^{\pm}(F) := \lim_{n \rightarrow \infty} \frac{\phi_{\sigma}^{\pm}(F^n G) - \phi_{\sigma}^{\pm}(G)}{n}$$

*always exists, and is independent of the choices of  $G$  and  $\sigma$ .*

- *The function*

$$h_t(F) := \lim_{n \rightarrow \infty} \frac{1}{n} \log \left( \sum_{k \in \mathbb{Z}} \dim \operatorname{Hom}(G, F^n G[k]) e^{-kt} \right),$$

*is a convex function in  $t$  satisfying:*

- $t \cdot \tau^+(F) \leq h_t(F) \leq h_0(F) + t \cdot \tau^+(F)$  for  $t \geq 0$ , and
- $t \cdot \tau^-(F) \leq h_t(F) \leq h_0(F) + t \cdot \tau^-(F)$  for  $t \leq 0$ .

## Theorem (F., 2023)

*Let  $X$  be an abelian variety. Then  $\tau = \tau^{\pm}: \operatorname{Aut}(\mathcal{D}^b(X)) \rightarrow \mathbb{R}$  is a quasimorphism.*

# Shifting numbers

## Theorem (F.–Filip, 2023)

- *The limit*

$$\tau^{\pm}(F) := \lim_{n \rightarrow \infty} \frac{\phi_{\sigma}^{\pm}(F^n G) - \phi_{\sigma}^{\pm}(G)}{n}$$

*always exists, and is independent of the choices of  $G$  and  $\sigma$ .*

- *The function*

$$h_t(F) := \lim_{n \rightarrow \infty} \frac{1}{n} \log \left( \sum_{k \in \mathbb{Z}} \dim \operatorname{Hom}(G, F^n G[k]) e^{-kt} \right),$$

*is a convex function in  $t$  satisfying:*

- ▶  $t \cdot \tau^+(F) \leq h_t(F) \leq h_0(F) + t \cdot \tau^+(F)$  for  $t \geq 0$ , and
- ▶  $t \cdot \tau^-(F) \leq h_t(F) \leq h_0(F) + t \cdot \tau^-(F)$  for  $t \leq 0$ .

## Theorem (F., 2023)

*Let  $X$  be an abelian variety. Then  $\tau = \tau^{\pm} : \operatorname{Aut}(\mathcal{D}^b(X)) \rightarrow \mathbb{R}$  is a quasimorphism.*



# Shifting numbers

## Theorem (F.–Filip, 2023)

- *The limit*

$$\tau^{\pm}(F) := \lim_{n \rightarrow \infty} \frac{\phi_{\sigma}^{\pm}(F^n G) - \phi_{\sigma}^{\pm}(G)}{n}$$

*always exists, and is independent of the choices of  $G$  and  $\sigma$ .*

- *The function*

$$h_t(F) := \lim_{n \rightarrow \infty} \frac{1}{n} \log \left( \sum_{k \in \mathbb{Z}} \dim \operatorname{Hom}(G, F^n G[k]) e^{-kt} \right),$$

*is a convex function in  $t$  satisfying:*

- ▶  $t \cdot \tau^+(F) \leq h_t(F) \leq h_0(F) + t \cdot \tau^+(F)$  for  $t \geq 0$ , and
- ▶  $t \cdot \tau^-(F) \leq h_t(F) \leq h_0(F) + t \cdot \tau^-(F)$  for  $t \leq 0$ .

## Theorem (F., 2023)

*Let  $X$  be an abelian variety. Then  $\tau = \tau^{\pm}: \operatorname{Aut}(\mathcal{D}^b(X)) \rightarrow \mathbb{R}$  is a quasimorphism.*

Thank you for your attention!

Reference:

- Dimitrov–Haiden–Katzarkov–Kontsevich. *Dynamical systems and categories*. arXiv:1307.8418.
- F.. *Entropy of an autoequivalence on Calabi-Yau manifolds*. MRL 2018.
- F.–Lai. *Nielsen realization problem for Bridgeland stability conditions on generic K3 surfaces*. arXiv:2302.12663.
- F.–Filip. *Asymptotic shifting numbers in triangulated categories*. Adv. Math 2023.
- F.. *Shifting numbers of abelian varieties via bounded t-structures*. arXiv:2306.13330.